

H₂O/H₂ 比に規定されるフォルステライト蒸発挙動 Evaporation of forsterite controlled by H₂O/H₂ in the ambient gas

橋 省吾¹, 瀧川 晶^{2*}
Shogo Tachibana¹, Aki Takigawa^{2*}

¹ 北海道大学大学院理学研究院, ² カーネギー研究所地磁気部門

¹Dept. Natural History Sci., Hokkaido Univ., ²Carnegie Inst. Washington, Dept. Terrestrial Magnetism

Evaporation experiments on forsterite have been revealed its evaporation kinetics in vacuum (free-evaporation regime) and in the presence of low-pressure hydrogen gas (hydrogen reaction-dominated regime). Tsuchiyama et al. (1999, GCA) proposed another evaporation regime called H₂O/H₂ buffer-dominated regime (HBD), where evaporation is controlled by the H₂O/H₂ ratio in the ambient gas. The HBD evaporation regime is important to consider the dust evolution in protoplanetary disks, but no experimental study has been done to investigate the evaporation kinetics. We have performed evaporation experiments on forsterite at low pressures with controlled H₂O/H₂ ratios, and have found that evaporation rates are controlled by the H₂O/H₂ ratio as proposed by Tsuchiyama et al. (1999).

キーワード: フォルステライト, 蒸発, 水蒸気, 速度論
Keywords: forsterite, evaporation, water vapor, kinetics