

震源断層周辺のヘリウム同位体比の分布：活断層の調査・評価技術の構築に向けて Geographic distribution of $^3\text{He}/^4\text{He}$ ratios along seismic source faults in Japan

梅田 浩司^{1*}, 浅森 浩一¹, 草野 友宏¹
Koji Umeda^{1*}, Koichi Asamori¹, Tomohiro Kusano¹

¹ 日本原子力研究開発機構 地層処分研究開発部門

¹ Geological Isolation Research and Development Directorate, Japan Atomic Energy Agency

It is well known that mantle degassing does not occur homogeneously over the Earth's surface. The elevated $^3\text{He}/^4\text{He}$ ratios found in volcanic regions and tectonically active areas are higher than the atmospheric values. This distribution is interpreted to indicate transfer of mantle volatiles into the crust by processes or mechanisms such as magmatic intrusion, continental underplating and lithospheric rifting. This study was undertaken to elucidate the geographic distribution of $^3\text{He}/^4\text{He}$ ratios around seismic source faults in Japan, using helium isotope data obtained from gas samples. Several case studies suggest that there is a significant trend of high ^3He emanations along the trace of active faults, resulting in leakage of mantle volatiles through crustal pathways (faults) due to more frequent development of higher permeability pathways and/or upwelling of mantle fluids through the ductile lower crust. From the viewpoint of site selection and implementation of a geological disposal facility, helium isotopes may be regarded as a tool for investigating and/or mapping concealed active faults with no surface expression.

Keywords: helium isotope, active fault