

2011年東北沖地震により誘発された紀伊半島沖の浅部低周波イベント Broadband features of the shallow low frequency events in Nankai trough, excited after the 2011 Tohoku-Oki earthquake

藤 亜希子¹, 尾鼻 浩一郎^{1*}, 高橋 成実¹, 深尾 良夫¹
Akiko To¹, Koichiro Obana^{1*}, Narumi Takahashi¹, Yoshio Fukao¹

¹ 海洋研究開発機構

¹ Japan Agency for Marine-Earth Science and Technology

Low frequency events are seismic events, which have longer duration and less energy radiation compared to regular earthquakes. The low frequency events detected in the shallow part of the Nankai trough (depth < 10 km), reported by previous studies, can roughly be divided into two groups depending on the observable frequency ranges of the signal, where the frequency ranges actually depend on the observed instrument.

The events of the first group are very low frequency earthquakes (VLFE), which were originally detected by broadband seismographs on-land (Ishihara et al., 2003; Ito & Obara, 2005), dominant in the frequency around 0.1-0.05 Hz. More recently a close-in observation was successfully made by temporally deployed broadband ocean-bottom seismometers (BBOBS), which revealed many intriguing features of the VLFEs (Sugioka et al., 2012). The events of the second group are low frequency tremors (LFT), which are recorded by OBSs equipped with 4.5 Hz short-period seismometer sited close to the source regions. They are dominant in the frequency range of 2-8 Hz with a lack of energy above 10 Hz (Obana & Kodaira, 2009). The classification between LFTs and VLFEs must be an important step toward estimating the physical process of the shallow low frequency events.

After the 2011 Mw9.0 Tohoku-oki earthquake, many shallow low frequency events were recorded at a cabled network of ocean bottom broadband stations (DONET) deployed in the northern part of Nankai trough. The characteristics of the events are similar to previously observed LFTs at the frequency range around 2-8 Hz. In addition, some of the events are accompanied by a lower frequency signal, clearly visible around 0.02-0.05 Hz, whose features are similar to those previously observed as VLFEs by Sugioka et al. (2012). One of such features of VLFEs is the ramp-type motion of the instrument-corrected seafloor displacement, which corresponds to a subsidence of up to 0.04 mm with a rise time of 10-20 s.

In order to examine whether the events accompanied by the 0.02-0.05 Hz signal are intrinsically different from those without the 0.02-0.05 Hz signal, the amplitudes of each event measured at 2-8 Hz and 0.02-0.05 Hz are compared. The comparison shows that the events without the 0.02-0.05 Hz signal tend to have lower amplitude in 2-8 Hz than those accompanied by the 0.02-0.05 Hz signal. The result indicates that there is no such event, which is intrinsically missing the 0.02-0.05 Hz components but has large amplitude in 2-8 Hz. In other words, the events without the 0.02-0.05 Hz signal are likely to be either smaller in size or occurred further away from the stations, compared to the events accompanied by the 0.02-0.05 Hz signal. Our dataset shows that the two types of low frequency events are likely the same phenomenon.

キーワード: 浅部低周波微動, 浅部低周波地震

Keywords: very long frequency earthquake, low frequency tremor