

広帯域MT探査による霧島火山群の3次元比抵抗構造

Three dimensional resistivity structure of Kirishima volcanoes inferred from anomalous magnetotelluric data

相澤 広記^{6*}, 小山 崇夫¹, 長谷 英彰¹, 上嶋 誠¹, 神田 径², 宇津木 充³, 吉村 令慧⁴, 山谷 祐介⁷, 橋本 武志⁵, 山崎 健一⁴, 小松 信太郎⁴, 渡邊 篤志¹, 小川 康雄²

Koki Aizawa^{6*}, Takao Koyama¹, Hideaki Hase¹, Makoto Uyeshima¹, Wataru Kanda², Mitsuru Utsugi³, Ryokei Yoshimura⁴, Yusuke Yamaya⁷, Takeshi Hashimoto⁵, Ken'ichi Yamazaki⁴, Shintaro Komatsu⁴, Atsushi Watanabe¹, Yasuo Ogawa²

¹ 東京大学地震研究所, ² 東京工業大学火山流体研究センター, ³ 京都大学火山研究センター, ⁴ 京都大学防災研究所, ⁵ 北海道大学地震火山研究観測センター, ⁶ 九州大学地震火山観測研究センター, ⁷ 産業技術総合研究所

¹Earthquake Research Institute, University of Tokyo, ²Volcanic Fluid Research Center, Tokyo Institute of Technology, ³Aso Volcanological Laboratory, Institute for Geothermal Sciences, Graduate School of Science, Kyoto, ⁴Disaster Prevention Research Institute, Kyoto University, ⁵Institute of Seismology and Volcanology, Faculty of Science, Hokkaido University, ⁶Institute of Seismology and Volcanology, Faculty of Sciences, Kyushu University, ⁷National Institute of Advanced Industrial Science and Technology

Broad-band magnetotelluric (MT) measurements were conducted on 2010?2011 around Shinmoe-dake volcano in the Kirishima volcanic group, Japan, where sub-Plinian eruptions took place three times on 26?27 January 2011. Combining with the previous MT data, it is found that the anomalous phase in excess of 90 degree is commonly observed at the northern part of the Kirishima volcanic group. Because the anomalous phase is not explained by 1-D or 2-D structure with isotropic resistivity blocks, 3-D inversions were conducted. By applying the small error bars on anomalous phase, we successfully estimated a 3-D resistivity structure that explains not only the usual data but also the anomalous phase data. The final model shows a eastward inclined and clockwise twisted pillar-like conductor that connects a deep-seated conductive body (at a depth greater than 10 km) to a shallow conductive layer at the central part of Kirishima volcanoes. By using the geophysical and petrological studies of the 2011 sub-Plinian eruptions, we infer that the pillar-like conductor represent the zone of hydrothermal aqueous fluids over 400 C, in which a magma pathway (interconnected melt) is partly and occasionally formed before magmatic eruptions. To the north of the deep conductor, earthquake swarms occurred on 1968?69, suggesting that these earthquakes were caused by volcanic fluids.