

## Chemical dynamics of snow in the Japanese Alps region

KURAMOTO, Takayuki<sup>1\*</sup> ; SASAKI, Akihiko<sup>1</sup> ; SUZUKI, Keisuke<sup>1</sup>

<sup>1</sup>IMS, Shinshu University

The Japanese Alps region is one of the heaviest snowy regions in Japan. In this area, winter precipitation is observed mainly two patterns such as winter monsoon pattern and low pressure pattern. Therefore, the chemical characteristics of the snowpack are different by snowfall types. In this study we aimed to clarify chemical dynamics of snowpack in the Japanese Alps region. We conducted a snow pit studies in the Japanese Alps region. We collected snow samples at the large flat place without obstruct by trees and the impact of human activities. The samples were melted, then pH, electric conductivity and major ions ( $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$ ,  $\text{Cl}^-$ ,  $\text{NO}_3^-$  and  $\text{SO}_4^{2-}$ ) were analyzed in clean room. The  $\text{Na}^+$  concentration correlates well with  $\text{Cl}^-$  concentration. These ions are considered to be sea-salt components. On the other hand,  $\text{SO}_4^{2-}$  concentrations included non-sea-salt components.

## Change for chemical component of rime ice in two decades

NAGAFUCHI, Osamu<sup>1\*</sup> ; YOKOTA, Kuriko<sup>2</sup> ; NAKAZAWA, Koyomi<sup>1</sup> ; HISHIDA, Naoko<sup>1</sup> ; IKEDA, Keisuke<sup>1</sup>

<sup>1</sup>the University of Shiga Prefecture, <sup>2</sup>Toyohashi University of Technology

Rime-ice and snow samples were collected at mountainous sites in Kyushu Island, Japan during from 1991 to 2014, and both soluble and insoluble substances in the melted rime-ice were analyzed by ion chromatography, inductively coupled plasma-mass spectrometry (ICP/MS) and analytical electron microscopy, in order to find the change of composition ratio of atmospheric pollutants cause by East Asian region. Although N/S ratio in rime-ice ranged from 0.1 to 0.3 in 1990's, recent year this ratio increased about 1.0. This phenomenon indicates that the composition of atmospheric pollutants changed during two decades in East Asian Continent.

## Variations in chemical composition of surface ice and meltwater on Svalbard glaciers

KONNO, Yudai<sup>1\*</sup> ; TAKEUCHI, Nozomu<sup>1</sup>

<sup>1</sup>Graduate School of Science, Chiba University

Snow and ice on glaciers usually contain small amounts of various soluble substances. These chemical substances were supplied through the atmosphere, rain, and snow. Chemical substances in glacier surface ice and melting water are important to understand atmospheric circulation, material cycle and the ecology of glacial microbes. We analyzed the major soluble ions of surface ice, melt water, and fresh snow for three glaciers (Austre Broggerbreen, Midtre Lovénbreen, and Pedersenbreen) in the northwestern part of Svalbard in the melting season of 2013.

The concentrations of all of the solutes were low generally, but the compositions varied among the samples. Based on the significant correlation on the concentration of each samples, the solutes could be classified into 3 groups: Group A ( $\text{Cl}^-$ ,  $\text{SO}_4^{2-}$ ,  $\text{Na}^+$ ), B ( $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$ ), and C ( $\text{NO}_3^-$ ,  $\text{NH}_4^+$ ,  $\text{K}^+$ ). They are likely to be derived from different sources. Group A solutes are probably sea salt origin, and Group B solutes are terrestrial dust origin, and Group C are anthropogenic or other unknown origin.

The altitudinal distribution of the concentration on Austre Broggerbreen showed that Group A (sea salt) solutes varied significantly although Group B (dust) solutes did not. Group A solutes were particularly higher in the higher area of the glacier. This variation can not be explained by distance from sea. It is probably due to supply from snow area remaining upper part of the glacier.

The chemical composition of the melting water was generally similar to those of the ice at the same site. However, some of the solutes ( $\text{K}^+$  and  $\text{SO}_4^{2-}$ ) were more abundant in meltwater than glacier ice at specific locations. This is probably due to supply from upper part of the glacier.

Keywords: Svalbard, glacier, chemical composition, sea salt

## Interannual variability of total SWE obtained by snow surveys in the Tuul river basin, Mongolia, from 2002 until 2013

SUGIURA, Konosuke<sup>1\*</sup> ; KADOTA, Tsutomu<sup>2</sup> ; IJIMA, Yoshihiro<sup>2</sup> ; KONYA, Keiko<sup>2</sup> ; ZHANG, Yinsheng<sup>3</sup> ; ISHIKAWA, Mamoru<sup>4</sup> ; YAMAZAKI, Takeshi<sup>5</sup> ; PUREVDAGVA, Khalzan<sup>6</sup> ; DAVAA, Gombo<sup>6</sup> ; OHATA, Tetsuo<sup>2</sup>

<sup>1</sup>University of Toyama/JAMSTEC, <sup>2</sup>JAMSTEC, <sup>3</sup>Institute of Tibetan Plateau Research, Chinese Academy of Science, <sup>4</sup>Hokkaido University, <sup>5</sup>Tohoku University, <sup>6</sup>Institute of Meteorology, Hydrology, and Environment, Mongolia

It is necessary for the southern limit of snow cover and the semi-arid region to be supplied with sustainable water. Mongolia is located in the southern limit and the semi-arid region. The capital, Ulaanbaatar, lies in a valley on the Tuul River. In order to investigate the actual conditions of snow water equivalent (SWE) in the upper Tuul River as water resources, the snow survey in the upper Tuul River has been carried out from 2002 to 2013. We have observed in each February when it will be the maximum amount of SWE. The snow water equivalent was estimated using a cylindrical snow sampler with 0.005-m<sup>2</sup> area. The total amount of SWE in the upper Tuul River basin was estimated using a global digital elevation model (DEM) with a horizontal grid spacing of 30 arc seconds (GTOP30) and a relationship between altitude and SWE. The main results obtained are shown below. The amount change of SWE in the upper Tuul River basin fluctuates over multiple years. The range of fluctuation was  $0.25 \pm 0.07 \text{ km}^3$ . It was not necessarily fluctuating in monotone. The steep rise of the amount of SWE in the upper Tuul River basin coincides with increasing air temperature and snowfall roughly.

Keywords: snow water equivalent, snow cover, water resources, water cycle, Mongolia

## Identifying the ice thickness of five perennial snow patches in the Tateyama Mountains based on GPR soundings

FUKUI, Kotaro<sup>1\*</sup> ; IIDA, Hajime<sup>1</sup>

<sup>1</sup>Tateyama Caldera Sabo Museum

We carried out ground penetrating radar (GPR) soundings in the Kuranosuke, the Hamaguri-yuki, the Tsurugisawa, the Chojiro and the Ikenotan-migimata perennial snow patches in the Tateyama Mountains, the northern Japanese Alps since 2012. The Kuranosuke and the Ikenotan-migimata perennial snow patches had large ice masses (>30 m in thickness). We had measured the surface flows of both ice masses since 2011. The maximum surface flows of the Ikenotan-migimata and the Kuranosuke perennial snow patches were about 2 m a<sup>-1</sup> and 0.14 m a<sup>-1</sup>, respectively. Thus, we regard the both snow patches as active glaciers.

The Hamaguri-yuki, the Tsurugisawa and the Chojiro perennial snow patch had thin ice masses (<20 m in thickness). It is possible that these ice masses are not flowing at the present time. Thus, we guess that these snow patches are glacierets rather than active glaciers.

Keywords: glacier, perennial snow patch, Mt. Tateyama, Mt. Tsurugi, GPR

## Studies on internal structure of active glacier in the Tateyama Mountains

IIDA, Hajime<sup>1\*</sup> ; FUKUI, Kotaro<sup>1</sup>

<sup>1</sup>Tateyama Caldera Sabo Museum

In 2013, we carried out 20 m depth boring in the Sannomado Glacier (2000 m above sea level), in the Tateyama Mountains, the northern Japanese Alps. The 20 m core was analyzed paying attention to the structure of firn and ice. The following observations were carried out;

- (i) stratigraphic observation of snow layer and glacier ice.
- (ii) measurement of density profile.
- (iii) measurement of grain shape and profile of grain size.
- (iv) observation of elongation of air bubbles.

The internal structure of the Sannomado Glacier was characterized by obvious boundary between firn and ice. At the depth of 5m, there was a distinct dirt layer formed in the last autumn. Above this dirt layer, firn transformed into ice abruptly and the density curve showed a discontinuity to  $850\text{kg/m}^3$ . The temperature of the snow and ice measured in the borehole was  $0\text{ }^\circ\text{C}$  throughout the layer. Spouting water was found in the borehole, indicating an aquifer in the glacier. From these results, such a rapid transformation process from snow to ice in this glacier will be discussed.

The grain size gradually increased with depth and elongation of air bubbles was shown below 15m which suggests internal flow of the glacier.

Keywords: glacier, perennial snow patch, Mt. Tateyama, Mt. Tsurugi, boring

## Contribution ratio of glacier discharge to the river water in Mongolian Altai

KONYA, Keiko<sup>1\*</sup> ; KADOTA, Tsutomu<sup>1</sup> ; DAVAA, Gombo<sup>2</sup> ; PURVDAGVA, Kalzan<sup>2</sup>

<sup>1</sup>JAMSTEC, <sup>2</sup>IMHE, Mongolia

The discharge from glaciers is an important theme for arid regions like Mongolia. The water from glaciers are accounted for the important water resources in Mongolia where little water is supplied from precipitation. It is to be revealed that how much water is available from the glaciers. In this study, we estimate how much water is available in the present states and in the future by analyzing water chemistry and water quantity.

We have measured the discharge, water temperature, electric conductivity (EC), pH, dD, d18O of the river at the beginning, middle and end of the melt season of the glacier. The contribution rate of the melt water to the river water were estimated by the two methods; A) discharge and EC, B) glacier melt rate.

The contribution rate was estimated to be 20-50 % of the flow. The future change of the contribution rate was estimated by method B with the temperature warming rate in the future estimated by the climate models in Mongolia. The results show that the river water is supposed to be increasing in next some decades and decreasing in next century. The snow melt water to the river is also need to be taken into account.

Keywords: glacier, glacier discharge, temperature warming, water resources, Mongolia, climate change

## Comparison of surrounding land features on the glacier terminal areas in the Himalayas derived from DEM

SUZUKI, Ryohei<sup>1\*</sup>

<sup>1</sup>none

Radiation field on the surface of mountain glaciers is reported to be controlled by the surrounding land features and does not distribute uniformly. Net radiation, which is obtained as a result of radiation budget, has been reported as an important factor to melt the ice surface of glaciers in the Himalayas by the previous studies. On the other hand, the lower areas of many glaciers in the south area of Himalayas are covered with debris that has variety of thermal properties, which corresponds to the one of the factor to cause the spatial variety of ice melt rates fields. Considering the surrounding land form features in the radiation field is probably required for estimating thermal properties on the debris-covered glaciers accurately. However, in-situ observation of those phenomena cannot be conducted mainly due to the difficulties of accessibility to the Himalayas. Thus, application of satellite remote sensing techniques is a powerful tool as an alternate method.

This study focuses on the Lunana region, Bhutan, which corresponds to the target area where in-situ observation has been carried out since 2002, as a case study to estimate the influence of surrounding land features around the lower glacier areas on their radiation field. I calculated a distribution of azimuth and zenith angles from each target point (pixel) on the lower glacier areas to each direction to the skylines with an approximate method to derive surrounding land feature; the value 1 corresponds to the full sky view and it decreases to 0 with the decrease of openness. My first result, which was derived from 8 directions for each pixel with the limit of about 4.5 km far from the point as a line of sight, shows it ranges from 0.7 to 0.9 regarding the three glaciers; Thorthormi, Lugge and Lugge II.

Because these values depend on the method to move the line of sight on the DEM, in other words, the method for processing the raster image as well as the accuracy of both elevations and horizontal positions, I would also like to present the other experimental results that were calculated with some different conditions to discuss the influence of the surrounding land form on the radiation field in the Himalayas.

Keywords: Digital Elevation Model (DEM), image processing, glacier melt rate, radiation budget, mountain glaciers, Himalayas

## Glacier lake and glacier lake outburst floods in Tien Shan and Ladakh Range

NARAMA, Chiyuki<sup>1\*</sup> ; KAZEHARE, Saiga<sup>1</sup> ; YAMAMOTO, Minako<sup>1</sup> ; UKITA, Jinro<sup>1</sup> ; IKEDA, Naho<sup>2</sup> ; TADONO, Takeo<sup>3</sup>

<sup>1</sup>Niigata University, Department of Environmental Science, <sup>2</sup>Tohoku University, Institute for Disaster Reconstruction and Re-generation Research, <sup>3</sup>JAXA

As a result of recent glacier melting, present glacier lakes develop rapidly at glacier fronts in Tien Shan Mountains, Central Asia and Ladakh Range, Indian Himalayas. Although glacier lakes in the Tien Shan and Ladakh Range are small, compared to those in the eastern Himalayas (Bhutan and eastern Nepal), several GLOFs in the past have caused fatalities and serious damage to infrastructure and crops. With recent changes in the development of glacier lakes in these mountain regions, floods are becoming an increasing threat to local residents. However, the current status of glacier lakes is poorly characterized in this region. In this study, we researched glacier lakes in the Tien Shan Mountains and Ladakh Range using high-resolution ALOS/PRISM-AVNIR-2 images taken in 2007-2010. In addition, we report the characteristic of glacier lake and glacier lake outburst floods (GLOFs) in Tien Shan, using investigation of the appearance of glacier lake and GLOFs in the past.

We examined about 1600 glacier lakes ( $>0.001 \text{ km}^2$ ) in the Tien Shan. Although glacier lakes are distributed throughout the Tien Shan Mountains, regional differences in their number and size are large. Larger glacier lakes are found in the Teskey and Ili-Kungoy regions, whereas most small glacier lakes occupy other mountain ranges (4000-5000 m asl), such as the Pskem, Talas, and Kyrgyz Ala-Too ranges. In seven mountainous regions, many present-day glacier lakes have appeared since the 1980s. Glaciers in these mountain regions are most glacier shrinkage area in Tien Shan over the 30 years, and these glacier lakes appeared after glacier shrinkage area. We report our results in detail in JpGU meeting.

Keywords: glacier lake, glacier lake outburst floods, small-size glacier lake region, Tien Shan, Ladakh Range

## Chemical Composition on the surface in the Urumqi No.1 glacier, Tien Shan, China

WAKABAYASHI, Kozue<sup>1</sup> ; TAKEUCHI, Nozomu<sup>1\*</sup> ; TANAKA, Sota<sup>1</sup> ; AMEMIYA, Shun<sup>1</sup>

<sup>1</sup>Graduate School of Science, Chiba University

Various chemical solutes are deposited in snow and ice on glaciers. Such solutes are usually washed out of glaciers during melting season. However, concentration and composition on snow and ice during melting season are little known. The solutes are important to understand microbes living on the glacier surface since chemical conditions affect their growth. In this study, we analyzed chemical compositions of ice surface in the melting season on Urumqi glacier No.1, Tien Shan Mountains in central Asia. Results showed that calcium ion was dominated more than 60% in all of the area on the glacier. This indicates that the chemical composition on the melting glacial surface is greatly affected by dust from desert in this region. Total concentration of nitrogen solutes (ammonium) was highest in the middle part of the glacier. Measurements of chlorophyll a concentration revealed that it was also maximal in the middle of the glacier. The altitudinal variation of solutes may affect the algal community and biomass on the glacier.

Keywords: chemical composition, glacier