

造礁性サンゴ骨格の結晶組織が示す aragonite と halite の共沈 New evidence for halite co-precipitation during coral calcification

甕 聡子^{1*}; 永井 隆哉¹; 川野 潤²; 渡邊 剛¹
MOTAI, Satoko^{1*}; NAGAI, Takaya¹; KAWANO, Jun²; WATANABE, Tsuyoshi¹

¹ 北海道大学大学院理学院, ² 北海道大学創成研究機構

¹Hokkaido University Faculty of Science, ²Creative Reseach Institution, Hokkaido University

In the last JpGU Meeting, we reported halite grains in coral skeleton through the observation of massive coral skeleton of *Porites lobata* by Analysis Transmission Electron Microscope (ATEM). Each halite grain typically shows a square shape and its grain size is around 80 nm. The spatial distribution of halite grains is inhomogeneous and seems to be independent on the arrangement of growth lines.

We observed new evidence that the halite grains in coral skeleton could precipitate during coral calcification. The electron diffraction patterns from some selected areas including both an aragonite and a halite grain show that there are special crystallographic orientation relationships between them. In consideration of misfit ratios between some selective bond lengths of halite and those of aragonite, crystallographic orientations of halite and aragonite seem to be a kind of hetero-epitaxial relationship.

This is the first observation for a primary precipitated mineral phase other than aragonite in coral skeletons. The halite phase in coral skeleton will provide a new perception for understanding the process of coral calcification.

Keywords: reef-building coral, calcification, biomineral, aragonite

サンゴのホウ素同位体組成から復元された20世紀半ば以降の北西太平洋熱帯域のpH変動 Ocean acidification in the tropical Northwest Pacific since the mid-20th century reconstructed from coral boron isotope

新城 竜一¹; 浅海 竜司^{1*}; Huang Kuo-Fang²; You Chen-Feng²; 井龍 康文³
SHINJO, Ryuichi¹; ASAMI, Ryuji^{1*}; HUANG, Kuo-fang²; YOU, Chen-feng²; IRYU, Yasufumi³

¹ 琉球大学・理, ²National Cheng Kung University, Taiwan, ³ 東北大学・院理
¹University of the Ryukyus, Japan, ²National Cheng Kung University, Taiwan, ³Tohoku University, Japan

Ocean acidification has been accelerating as a result of absorption of increasing anthropogenic CO₂ in the atmosphere emitted by fossil-fuel combustion and land-use practices since the Industrial Revolution, which can be resulting in decreased surface ocean pH and posing a critical threat to marine ecosystems (e.g., Sabine et al., 2004; Orr et al., 2005). By the end of the 21st century, predictions based on different scenarios indicate that ocean pH will decrease by 0.3-0.4 pH units. Only a few long-term continuous observations of sea surface pH have been derived from Station ALOHA off the Hawaiian Islands in the North Pacific, near the Bermuda Islands, and near the Canary Islands in the North Atlantic. A decreasing pH trend in the western North Pacific surface waters for 1983-2007 was estimated from the observational data of oceanic CO₂ partial pressure and related properties (Midorikawa et al., 2010). To elucidate the natural variability of ocean pH and assess the actual trend in ocean acidification more accurately, we must go further back in time. For these reasons, we rely on paleo-pH archives or other related parameters.

Massive corals, an informative archive of past ocean environments, precipitate annually banded calcium carbonate skeletons at a relatively rapid rate (about 1 cm per year), allowing for accurate chronological control and high-resolution sampling. Because of pH-dependent isotopic fractionation between the two dominant boron species in seawater, boron-isotopic systematics in marine carbonates provide a potential proxy for ocean pH in the past (e.g., Hemming and Hanson, 1992). Nevertheless, only two previous investigations provided boron-isotope time series from long-lived corals from the Great Barrier Reef in the South Pacific for the last 300 years. Unlike seawater temperature and salinity records (Asami et al., 2005; Felis et al., 2009), no coral-based reconstruction of long-term pH variation in the North Pacific has been reported.

Here, we generated an annually resolved 60-year-long (1940-1999 A.D.) record of seawater pH from boron isotope composition in a *Porites* coral collected in Guam Island, located in the Western Pacific Warm Pool which contains the highest annual sea surface waters and serves as a heat engine for the earth climate. The first long-term continuous boron isotope-pH proxy record in the North Pacific from the coral provides evidence of a slight ocean acidification trend (equivalent to 0.05-0.08 pH units for surface water) since the mid-20th century, although the critical factors that affect interannual variability remain unknown (Shinjo et al., 2013). From this perspective, the results of this study will provide improved constraints on global atmosphere-ocean interaction models and understanding of the future coral reef ecosystems.

キーワード: サンゴ骨格, ホウ素同位体組成, pH, 海洋酸性化, 北太平洋

Keywords: coral skeleton, boron isotope composition, pH, ocean acidification, North Pacific

有孔虫細胞軟組織の NanoSIMS による窒素同位体比マッピング 15N/14N mapping of the isotope labeling cultured foraminifera using ultra thin section

野牧 秀隆^{1*}; 石田 章純²; 白井 厚太郎²; 力石 嘉人¹; 植松 勝之³; 多米 晃裕³; 土屋 正史¹; 豊福 高志¹; 高畑 直人²; 佐野 有司²; 大河内 直彦¹

NOMAKI, Hidetaka^{1*}; ISHIDA, Akizumi²; SHIRAI, Kotaro²; CHIKARAISHI, Yoshito¹; UEMATSU, Katsuyuki³; TAME, Akihiro³; TSUCHIYA, Masashi¹; TOYOFUKU, Takashi¹; TAKAHATA, Naoto²; SANO, Yuji²; OHKOUCHI, Naohiko¹

¹ 海洋研究開発機構, ² 東京大学大気海洋研究所, ³ マリン・ワーク・ジャパン

¹JAMSTEC, ²AORI, University of Tokyo, ³Marine Works Japan

Shallow water benthic foraminifera, *Ammonia beccarii*, survive under anoxic conditions in conjunction with possible endobionts. Based on the amino acid analysis, those endobionts are expected to utilize nitrate pool in the foraminifera. However, nitrogen cycles in the foraminiferal cell and endobionts are still unclear. Here, we obtained two dimensional-nitrogen isotopic compositions of *A. beccarii* which had been incubated under oxic and anoxic conditions with ¹⁵N-labeled nitrate. After observing with transmission electron microscope to confirm cellular ultrastructure and endobiont distribution, same ultra thin section was examined for nitrogen isotopic composition analysis using secondary ion mass spectrometer. Nitrogen isotopic compositions were measured with spatial resolution better than 400 nm. ¹⁵N-enriched parts were found in certain structures in the cell, but not in the endobionts in this experiment.

キーワード: 底生有孔虫, 硝酸塩呼吸, 共生微生物, 高分解能二次イオン質量分析装置

Keywords: Benthic foraminifera, nitrate respiration, symbiotic microbe, NanoSIMS