

A Brief History of Collaborative Study on Equatorial MLT Dynamics using Meteor and MF Radars in Indonesia

A Brief History of Collaborative Study on Equatorial MLT Dynamics using Meteor and MF Radars in Indonesia

TSUDA, Toshitaka^{1*} ; DJAMALUDDIN, Thomas² ; YATINI, Clara² ; BUDIYONO, Afif² ; VINCENT, Robert³ ; REID, Iain³
TSUDA, Toshitaka^{1*} ; DJAMALUDDIN, Thomas² ; YATINI, Clara² ; BUDIYONO, Afif² ; VINCENT, Robert³ ; REID, Iain³

¹Research Institute for Sustainable Humanosphere (RISH), Kyoto University, ²Indonesian National Institute of Aeronautics and Space (LAPAN), ³Department of Physics, University of Adelaide

¹Research Institute for Sustainable Humanosphere (RISH), Kyoto University, ²Indonesian National Institute of Aeronautics and Space (LAPAN), ³Department of Physics, University of Adelaide

In the tropics active cumulus convection generates various atmospheric waves, such as Kelvin waves, planetary waves, tides, and gravity waves. The wave energy and momentum are transported upward through propagating of these waves. Wave-mean flow interactions are crucially important for understanding of dynamical processes in the equatorial atmosphere, including the formation of peculiar long-term variations such as quasi-biennial oscillation (QBO) and semi-annual oscillation (SAO) in both the stratosphere and the MLT (mesosphere and lower thermosphere) region (70-120 km).

We constructed a total of five meteor and medium frequency (MF) radars in Indonesia since 1992 under close collaboration between RISH, LAPAN and the University of Adelaide. The MLT radar network has been expanded in India, Central and Eastern Pacific, and China. These radars have clarified the behavior of atmosphere dynamics in the MLT region. This paper gives an overview of our collaborative studies as well as highlights of scientific achievements using the MLT radar network

キーワード: mesosphere and lower thermosphere, equatorial atmosphere, atmospheric waves, meteor radar, medium frequency radar, Indonesia

Keywords: mesosphere and lower thermosphere, equatorial atmosphere, atmospheric waves, meteor radar, medium frequency radar, Indonesia

Atmospheric Waves in the MLT: A Review Atmospheric Waves in the MLT: A Review

VINCENT, Robert^{1*}
VINCENT, Robert^{1*}

¹University of Adelaide
¹University of Adelaide

Through their efficient transfer of energy and momentum, atmospheric waves propagating up from the lower atmosphere play an important role in determining the structure of the Mesosphere/Lower Thermosphere (60-100 km). A wide range of wave types are involved, with periods ranging from minutes to days. Here we review developments in our understanding of wave coupling and impacts on the MLT, with an emphasis on developments in the past decade.

キーワード: MLT Dynamics, Gravity Waves, Atmospheric Tides, Planetary Waves
Keywords: MLT Dynamics, Gravity Waves, Atmospheric Tides, Planetary Waves

MLT 領域での nonmigrating tide のふるまいについて Behavior of non-migrating tides in the MLT region

三好 勉信^{1*}; 藤原 均²
MIYOSHI, Yasunobu^{1*}; FUJIWARA, Hitoshi²

¹九州大学, ²成蹊大学
¹Kyushu University, ²Seikei University

It is well established that non-migrating tides have significant amplitudes in the mesosphere and lower thermosphere (MLT). Using a general circulation model that contain the region from the ground surface to the upper thermosphere, behavior and excitation sources of non-migrating tides are examined. In this study, behaviors of the westward moving semidiurnal tide with zonal wavenumber 1 (SW1), the semidiurnal tide with zonal wavenumber 0 (S0) and the diurnal tide with zonal wavenumber 0 (D0) are examined in detail. There are two main sources for non-migrating tides. One is latent heat release due to the cumulus convection in the troposphere. The other is the nonlinear interaction between the migrating tide and the stationary planetary wave in the middle atmosphere. Our results indicate that the amplitudes of SW1, S0 and D0 are enhanced when the stationary planetary wave in the stratosphere and mesosphere is active. This means that SW1, S0 and D0 are mainly excited by the nonlinear interaction between the migrating tide and the stationary planetary wave. Furthermore, we discuss excitation sources of other non-migrating tides, such as the eastward moving diurnal tide with zonal wavenumber 3 (DE3) and the eastward moving semidiurnal tide with zonal wave number 2 (SE2).

キーワード: 大気潮汐波, 大気大循環モデル
Keywords: Tides, General Circulation Model

Long-term observations of MLT zonal wind variations in relation to stratospheric zonal winds over low-latitudes
Long-term observations of MLT zonal wind variations in relation to stratospheric zonal winds over low-latitudes

GRANDHI, Kishore kumar^{1*} ; KARANAM, Kishore kumar² ; SINGER, Werner¹ ; ZULICKE, Christoph¹ ; S, Gurubaran³ ; GERD, Baumgarten¹ ; RAMKUMAR, Geetha² ; S, Sathishkumar⁴ ; RAPP, Markus⁵
GRANDHI, Kishore kumar^{1*} ; KARANAM, Kishore kumar² ; SINGER, Werner¹ ; ZULICKE, Christoph¹ ; S, Gurubaran³ ; GERD, Baumgarten¹ ; RAMKUMAR, Geetha² ; S, Sathishkumar⁴ ; RAPP, Markus⁵

¹Leibniz Institute of Atmospheric Physics, University of Rostock, Kuhlungsborn, Germany, ²Space Physics Laboratory, Vikram Sarabhai Space Center, Trivandrum, India, ³Indian Institute of Geomagnetism, Navi Mumbai, India, ⁴Equatorial Geophysical Research Laboratory, Indian Institute of Geomagnetism, Tirunelveli, India, ⁵German Aerospace Center Institute of Atmospheric Physics (IPA), Oberpfaffenhofen, Wessling, Germany

¹Leibniz Institute of Atmospheric Physics, University of Rostock, Kuhlungsborn, Germany, ²Space Physics Laboratory, Vikram Sarabhai Space Center, Trivandrum, India, ³Indian Institute of Geomagnetism, Navi Mumbai, India, ⁴Equatorial Geophysical Research Laboratory, Indian Institute of Geomagnetism, Tirunelveli, India, ⁵German Aerospace Center Institute of Atmospheric Physics (IPA), Oberpfaffenhofen, Wessling, Germany

Long-term observations from medium-frequency and meteor radars (1993-2012) and rocket soundings (1979-1990 and 2002-2007) are used to study mesosphere lower thermosphere (MLT) zonal wind variations in relation to the stratospheric winds over Northern low-latitudes. The combined dataset provide a complete height profile of amplitude of semiannual oscillation (SAO) up to 100 km, with an exception around 75-80 km. The SAO signal has maxima around 50 km and 82 km and a minimum around 65 km. The MLT zonal winds show remarkable inter-annual variability during spring equinox and much less during fall equinox. Zonal wind mesospheric spring equinox enhancements (MSEE) appear with a periodicity of 2-3 years suggesting a modulation by the quasi-biennial oscillation, which we identified with the strength of stratospheric westward winds. Out of 20 years of observations, the stratospheric westward winds are strong during 11 years (non-MSEE) and weak during 9 years. Six of these years show large MLT winds (MSEE) and 3 years (1999, 2004 and 2006) show small MLT winds (missing-MSEEs). These unexpected small winds occur in years with global circulation anomalies as identified with strong sudden stratospheric warmings and an early spring transition of zonal winds, along with a minor enhancement in the tidal amplitudes.

キーワード: MLT winds, MSAO, Meteor radar, MF radar, QBO
Keywords: MLT winds, MSAO, Meteor radar, MF radar, QBO

The saturation of gravity waves traveling from the lower to the upper atmosphere observed by the MU radar and understood
The saturation of gravity waves traveling from the lower to the upper atmosphere observed by the MU radar and understood

加藤 進^{1*}; 津田 敏隆¹; 山本 衛¹; 中村 卓司²

KATO, Susumu^{1*}; TSUDA, Toshitaka¹; YAMAMOTO, Mamoru¹; NAKAMURA, Takuji²

¹Research Institute for Sustainable Humanosphere, Kyoto University, ²National Polar Research Institute)

¹Research Institute for Sustainable Humanosphere, Kyoto University, ²National Polar Research Institute)

The MU radar of Kyoto University was constructed in 1984. One of the main purpose of the radar construction is to observe atmospheric gravity waves particularly to find how gravity waves saturate in traveling from the lower atmosphere to the upper atmosphere. In the 1980s Matsuno, Geller and others put forwards an idea suggesting that the gravity wave saturation may release momentum for driving the mesosphere general circulation. Their idea is based on rocket and satellite global- observation of winds and temperature varying peculiarly with seasons in the mesosphere.

Our MU radar observation has been successful in proving the gravity wave momentum release to be in a good agreement with the required quantity for the mesosphere general circulation. Also our success of precise measurements of the saturated gravity wave power spectrum strongly supports to explain the gravity wave saturation idea in terms of a simple theory based on the linear or monochromatic gravity wave theory by Hines in 1960.

Our theory on the basis of our MU radar observation shows that the gravity wave saturation is attained for each gravity wave in amplitude reaching the phase speed due to a balance between the increasing amplification expected by the linear theory and non-linear braking effects. We can consider that the original gravity wave dispersion relation is maintained upon the saturation.

Gravity waves should experience a number of such saturations before reaching the thermosphere on the way.

Ducted Concentric Gravity Wave Observed by IMAP/VISI Associated with Super Typhoon Haiyan

Ducted Concentric Gravity Wave Observed by IMAP/VISI Associated with Super Typhoon Haiyan

PERWITASARI, Septi^{1*}; SAKANNOI, Takeshi¹; YAMAZAKI, Atsushi²; OTSUKA, Yuichi³; HOZUMI, Yuta⁴; AKIYA, Yusuke⁴; SAITO, Akinori⁴; SUZUKI, Shin³
PERWITASARI, Septi^{1*}; SAKANNOI, Takeshi¹; YAMAZAKI, Atsushi²; OTSUKA, Yuichi³; HOZUMI, Yuta⁴; AKIYA, Yusuke⁴; SAITO, Akinori⁴; SUZUKI, Shin³

¹PPARC, Tohoku University, ²JAXA/ISAS, ³STEL, Nagoya University, ⁴Geophysics Dept., Kyoto University

¹PPARC, Tohoku University, ²JAXA/ISAS, ³STEL, Nagoya University, ⁴Geophysics Dept., Kyoto University

Although the convection activity in the troposphere is generally accepted as one of important source of gravity waves in the mesosphere and lower thermosphere, however it is still uncertain how these waves can reach these regions and what types of waves are generated. For decades, the study of gravity waves has been classified into two categories; first is that the waves travel directly from the source and the second is that the waves are ducted or trapped. Many studies tried to explain both categories yet all studies focused on gravity waves produced by transient events. There were almost no observation reports of airglow emissions during a large storm and what type of gravity waves and typical wavelength can be produced from such event. To address this issue, a space-based observation is more preferable since it covers wider area. Until recently, IMAP/VISI is the only space-based instrument that capable of imaging gravity waves above the troposphere in the nadir direction. The Visible and near-Infrared Spectral Imager (VISI) of the IMAP mission was launched successfully on July 21, 2012 with H-IIB/HTV-3 and installed onto the International Space Station (ISS). IMAP/VISI is now operated in the night side hemisphere with a range of +/- 51 deg. GLAT. IMAP/VISI is measuring three different airglow emissions of OI at 630 nm, the OH Meinel band at 730 nm and the O₂ (762 nm) atmospheric band at 762 nm at an altitude of ~400 km with the typical spatial resolution of 16-50 km.

We found concentric gravity waves events in the southeastern part of Australia that was observed around 13-15 UT for 3 days from 6-8 November 2013 in O₂ (0-0) airglow emission by IMAP/VISI. The waves have horizontal wavelength vary from 80 – 210 km. By using the least squares method, the curvature of the waves was fitted to a perfect circle. The center of the wave was found to be around 155°E; -42°S with the radius varies from 400-1200 km. From the meteorological satellite, we cannot locate any convective source around the center of the wave. The nearby local convective source was located a few hundreds km to the south of the wave center and the rainfall rate was less than 10 mm/hr. Therefore, we rule out the possibility of local convective activity as the source of these waves. From the past studies, there were evidences that the gravity waves may be ducted and traveled a great distance away from a specific convective source (e.g. Nakamura et al., 1999; Walterscheid et al., 1999; Hecht et al., 2001). Their studies suggested that the gravity waves observed in Australia were originated from convective activity several thousands km north of Australia. During the observed events, the Typhoon Haiyan was underway. On November 6, the typhoon was categorized as 5 – equivalent of super typhoon and reached its peak on November 7 and then made a landfall in Philippine on November 8. In this study, we argue that the concentric gravity waves seen by IMAP/VISI could be generated by the intense convective activity associated with the Haiyan Typhoon event. Background wind data from TIDI (TIMED Doppler Interferometer) and MF Radar will be used to examine the plausibility for the formation of a ducted/trapped region that can explain the long distance propagation of these waves. The temperature profile from MSISE-90 model will also be used to examine the mesospheric inversion layer and if it's possible to get the data, we will also use the ground-based airglow imager data from Adelaide and Alice Spring.

キーワード: IMAP/VISI, O₂ (0-0), concentric gravity wave, ducted, typhoon Haiyan

Keywords: IMAP/VISI, O₂ (0-0), concentric gravity wave, ducted, typhoon Haiyan

The MF Radar Technique: a Review The MF Radar Technique: a Review

REID, Iain^{1*}

REID, Iain^{1*}

¹Department of Physics, University of Adelaide, ²ATRAD Pty Ltd

¹Department of Physics, University of Adelaide, ²ATRAD Pty Ltd

The Medium Frequency (MF) radar technique has been applied for more than five decades to measure winds and turbulence in the upper atmosphere in the region between 60 and 100 km during the day, and between 80 and 100 km at night. It is one of the few techniques able to provide winds reliably in the 60 to 80 km height region during the day. Although some care is needed in interpretation of the results, it remains a powerful and very useful technique. In this paper, we review the technique and highlight some recent recent results.

キーワード: Radar, Medium Frequency, Spaced Antenna, Mesosphere Lower Thermosphere, Winds, Turbulence
Keywords: Radar, Medium Frequency, Spaced Antenna, Mesosphere Lower Thermosphere, Winds, Turbulence

アラスカ及びトロムソ MF レーダーで観測された 中間圏重力波の日内変動 Tidal periodicity of mesospheric gravity waves observed with MF radar at Poker Flat, Alaska and at Tromsø, Norway

村山 泰啓^{1*}; 木下 武也¹; 川村 誠治¹; 野澤 悟徳²; ホール クリス³
MURAYAMA, Yasuhiro^{1*}; KINOSHITA, Takenari¹; KAWAMURA, Seiji¹; NOZAWA, Satonori²; HALL, Chris³

¹ 情報通信研究機構, ² 名古屋大学太陽地球環境研究所, ³ トロムソ大学

¹National Institute of Information and Communications Technology, ²Solar-Terrestrial Physics Laboratory, ³Tromsø University

中間圏における重力波と潮汐波、それらの相互作用に関する観測研究は、以前から様々な地点で行われてきた (e.g., Saskatoon, Canada (Manson et al. 1998), Rothera, Antarctica (Beldon and Mitchell, 2010)). アラスカではポーカーフラット、ノルウェーではトロムソに設置された MF レーダーにより、中間圏から下部熱圏における中性風速データが 1990 年代後半より蓄積されている。本研究では昨年引き続き、潮汐波と重力波の結合プロセスの理解を深めるため、10 年間 (1999 ~ 2008 年) の上記観測データを用いて、中間圏重力波と 12 時間及び 24 時間周期成分の日内および季節変動を調べた。まず始めに、トロムソ MF レーダー観測の水平風速 30 分間隔データ 5 日分の時系列に 8, 12, 24, 48 時間周期成分の調和フィッティングを行った。重力波はこれらの残差で 1~12 時間周期を持つ擾乱と定義した。上記手法を 30 分間隔ごとに適用し、12 時間及び 24 時間周期成分の振幅や位相の 5 日間移動平均値を計算した。得られた 12 時間及び 24 時間周期成分と重力波の運動エネルギーとの関係を調べた結果、今年の解析におけるポーカーフラット MF レーダー観測で 5~8 月に見られた 1~4 時間周期の重力波運動エネルギーの 2 つのピークは見られなかった。一方で 11~2 月にかけて、6UT 付近でエネルギーの増大が確認された。これは、24 時間周期成分の東風が最大、または 12 時間周期成分の東風から西風に変わる時刻に対応していた。続いて、ポーカーフラット MF レーダー観測で 2000 年 11 月 15 日から約 20 日間続いた東西風の半日成分の位相に重力波運動エネルギーがロックされる現象がトロムソでも見られるか確認した。その結果、同日から約 10 日間は同様の現象が見られたが、重力波運動エネルギーの位相が 90 度ずれて対応していた。

ポーカーフラットで見られた現象は、この時期以外の期間、高度においてはよく見られ、一ヶ月以上続く事例も確認された。今後は、それぞれの地点における潮汐波の太陽同期、非同期成分及び背景場に焦点をあて、重力波特性の違いを引き起こす要因を明らかにすしたいと考えている。

南極昭和基地および Davis 基地上空の中間圏下部熱圏領域における短周期潮汐波の振る舞い
Characteristics of Short Period Tidal Components in Antarctic MLT above Syowa and Davis

堤 雅基^{1*}; Murphy Damian²
TSUTSUMI, Masaki^{1*}; MURPHY, Damian²

¹ 国立極地研究所, ² 豪南極局

¹National Institute of Polar Research, Japan, ²Australian Antarctic Division

The behaviour of short period atmospheric tidal components in the Antarctic mesosphere and lower thermosphere is studied based on long term observations over Syowa (69.0S, 39.6E) and Davis (68.6S, 78.0E) stations. Semidiurnal tides in the Antarctic mesosphere and lower thermosphere have been extensively studied through the recently established Antarctic radar network [e.g., Murphy et al., 2006; 2008]. However, details of shorter period components such as terdiurnal and six-hour tides are less investigated and poorly known because of their smaller amplitudes compared to the semidiurnal and diurnal tides in the height region of conventional MF radar observations of around 70-90 km. These short period tides also fall in the frequency range of inertial gravity waves and are often hard to distinguish from these waves.

The characteristics of the terdiurnal tide above Davis and Syowa have been measured on a short-term to seasonal basis in the mesosphere and lower thermosphere using long-term simultaneous MF radar data at the two sites (1999-). The terdiurnal tide achieves moderate amplitudes in the winter at these heights but there are subtle differences between the two stations. These differences are explored further by differencing tide phasors in local time and checking the amplitude of the result on a seasonal basis. If the terdiurnal tide was made up entirely of migrating components, this difference would yield a zero-average amplitude. However, the observed non-zero values suggest that the terdiurnal tide at these latitudes contains strong non-migrating tidal components.

The Syowa MF radar has a great advantage over other MF radars in that it has been conducting simultaneous meteor wind measurements together with the conventional correlation based measurements, which enables wind observations in a very wide height range of 65-120km [Tsutsumi and Aso, 2005]. A clear enhancement in terdiurnal amplitudes is seen in early winter months of March-June. The amplitudes can reach 20 m/s around 110 km even in the composite plot made with more than 10 years of data. These amplitudes can be comparable or sometimes even larger than those of diurnal and semidiurnal tides, and indicate a possible significant role of short period tidal components in the polar lower thermosphere.

Keywords: Antarctic, mesosphere and lower thermosphere, short period atmospheric tidal waves

SMILES が捉えた日食時の中間圏オゾンの変動について The SMILES observations of mesospheric ozone during the solar eclipse

今井 弘二^{1*}; 秋吉 英治²; 高橋 けんし³; 山下 陽介²; 今村 隆史²; 鈴木 睦¹; 海老沢 研¹; 塩谷 雅人³
IMAI, Koji^{1*}; AKIYOSHI, Hideharu²; TAKAHASHI, Kenshi³; YAMASHITA, Yousuke²; IMAMURA, Takashi²; SUZUKI,
Makoto¹; EBISAWA, Ken¹; SHIOTANI, Masato³

¹ 宇宙航空研究開発機構 宇宙科学研究所, ² 国立環境研究所, ³ 京都大学 生存圏研究所

¹Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency, Sagami-hara, ²National Institute for Environmental Studies, ³Research Institute for Sustainable Humanosphere, Kyoto University

Solar eclipse temporally reduces the amount of solar radiation, providing an opportunity to verify the ozone photochemistry under changing solar radiation. During the longest annular solar eclipse in this millennium occurred on 15 January 2010, Superconducting Submillimeter-Wave Limb-Emission Sounder (SMILES) successfully captured increased ozone mostly in the mesosphere with a decrease in solar illuminations. The ozone increment shows altitude dependence in the mesosphere. Using an atmospheric chemistry box model, it is found that the dependence results from the difference in chemical reaction rates to the solar radiation change. The model also predicts the difference in the ozone concentration evolution between the sunlight decreasing and increasing phases, although SMILES observation does not resolve the difference.

キーワード: SMILES, オゾン, 中間圏

Keywords: SMILES, ozone, mesosphere

超高層大気研究への IUGONET データ解析システムの貢献 Contribution of the IUGONET data analysis system to upper atmospheric researches

新堀 淳樹^{1*}; 八木 学²; 田中 良昌³; 佐藤 由佳³; 谷田貝 亜紀代⁴; 梅村 宜生⁴; 上野 悟⁵; 小山 幸伸⁶; 阿部 修司⁷
SHINBORI, Atsuki^{1*}; YAGI, Manabu²; TANAKA, Yoshimasa³; SATO, Yuka³; YATAGAI, Akiyo⁴; UMEMURA, Norio⁴;
UENO, Satoru⁵; KOYAMA, Yukinobu⁶; ABE, Shuji⁷

¹ 京都大学生存圏研究所, ² 東北大学理学研究科 惑星プラズマ・大気研究センター, ³ 国立極地研究所, ⁴ 名古屋大学太陽地球環境研究所 ジオスペース研究センター, ⁵ 京都大学大学院理学研究科附属天文台, ⁶ 京都大学大学院理学研究科附属地球磁気世界資料解析センター, ⁷ 九州大学 国際宇宙天気科学・教育センター

¹Research Institute for Sustainable Humanosphere (RISH), Kyoto University, ²Planetary Plasma and Atmospheric Research Center, Graduate School of Science, Tohoku University, ³National Institute of Polar Research, ⁴Nagoya University Solar Terrestrial Environment Laboratory Geospace Research Center, ⁵Kwasan & Hida Observatories, School of Science, Kyoto University, ⁶Data Analysis Center for Geomagnetism and Space Magnetism Graduate School of Science, Kyoto University, ⁷International Center for Space Weather Science and Education, Japan

Various kinds of atmospheric disturbances and long-term variation as seen in several parameters (temperature, mean wind etc.) in the upper atmosphere (mesosphere, thermosphere and ionosphere) is caused by energy input from solar radiation, momenta and energies from the lower atmosphere (stratosphere and troposphere) via atmospheric waves, and chemical reaction. Such atmospheric phenomena observed by ground-based and satellite instruments are the result of such complicated processes. In order to investigate the mechanisms of the atmospheric disturbances and long-term variations in the upper atmosphere, which may be affected by solar activities and global warming, researchers need to conduct comprehensive analyses with various kinds of long-term observation data that have been continued by means of a global network of MST (Mesosphere-Stratosphere-Troposphere) radars, optical sensors, radiosondes, etc. The IUGONET (Inter-university Upper atmosphere Global Observation NETWORK) project initiated in 2009 aims at the establishment of a cross-reference system for various kinds of ground-based observation data. The IUGONET participants consist of five universities/institutes: the National Institute of Polar Research (NIPR), Tohoku University, Nagoya University, Kyoto University, and Kyushu University. We have developed a metadata database (MDB) of ground-based observations and IUGONET data analysis software (UDAS) in order to provide researchers in a wide range of disciplines with a seamless data environment to link databases spread across the IUGONET institutions. In particular, the MDB and UDAS will be of great help in data acquisition and integrated analyses to understand the dynamics on the mesosphere-lower thermosphere (MLT) throughout the Sun-Earth system. Therefore, the IUGONET MDB and UDAS will greatly contribute to upper atmospheric researches on the basis of integrated analysis of various kinds of long-term observation data covering a wide region from both the poles to the equator. In this talk, we introduce a brief overview of the IUGONET project and an application of the IUGONET products for upper atmospheric researches.

キーワード: 超高層大気, 地上観測データ, ユーゴネット, メタデータ検索システム, 統合解析ツール

Keywords: Upper atmosphere, Ground-based observation data, IUGONET, metadata search system, IUGONET data analysis tool

Vertical and lateral wave coupling observed with network of MLT/MST Radars over Indian region

Vertical and lateral wave coupling observed with network of MLT/MST Radars over Indian region

M Venkat Ratnam^{1*}; S. Eswariah²; N. Venkateswara rao¹; S. Vijayabhaskar rao²; K. Kishore kumar³; S. Sathish kumar⁴; S. Gurubaran⁴

M, Venkat ratnam^{1*}; S., Eswariah²; N., Venkateswara rao¹; S., Vijayabhaskar rao²; K., Kishore kumar³; S., Sathish kumar⁴; S., Gurubaran⁴

¹National Atmospheric Research Laboratory, ²Sri Venkateswara University, ³Space Physical Laboratory, ⁴Indian Institute of Geomagnetism

¹National Atmospheric Research Laboratory, ²Sri Venkateswara University, ³Space Physical Laboratory, ⁴Indian Institute of Geomagnetism

It is well known that gravity waves and tides play an important role in delineating the middle atmospheric structure and dynamics. There have been several studies in recent years, using different measurement techniques, to understand significant roles played by gravity waves and tides in the lower, middle and upper atmospheres. However, only a few studies addressed this problem with simultaneous observations of all the three regions. Moreover, no efforts have been made so far to understand the lateral forcing of these waves and tides since such a study needs a network of radars located nearby which was missing. With the establishment of an advanced meteor radar at Sri Venkateswara University, Tirupati (13.63oN, 79.4oE), India, and up gradation of MF radar at Kolhapur (16.8oN, 74.2oE) together with MST radar at Gadanki (13.5oN, 79.2oE), Meteor radar at Thumba (8.5oN, 77oE) and MF radar at Tirunalveli (8.7oN, 77.8oE) forms a unique network of radars in the tropical region. Importantly, all these radars are located within 1000 km distance. Accordingly, this network is suitable to study the lower atmospheric forcing and its impacts on middle and upper atmospheric structure and dynamics. For the present study, all these radars were simultaneously operated for a few days in September 2013. These observations show the presence of short period gravity waves and tides (diurnal, semi-diurnal and ter-diurnal) at all locations. Large day-to-day variability in gravity waves and tides is observed within a station and among different stations providing insight on lateral coupling. Phase propagations of the three tidal components at different stations is used to further understand the lateral coupling. Using simultaneous MST radar, Rayleigh lidar and SVU meteor radar (which are nearly co-located), lower atmospheric forcing and its impacts on the mesosphere and lower thermosphere are investigated. This study showed need for long-term measurements, with simultaneous operation of all the above mentioned network of radars, to effectively address the problem of vertical and latitudinal wave forcing.

キーワード: Coupling, Meteor/MF radars, Tropical MLT region

Keywords: Coupling, Meteor/MF radars, Tropical MLT region