

About drift, oscillations and steps of the center of mass of the Moon About drift, oscillations and steps of the center of mass of the Moon

BARKIN, Yury^{1*} ; HANADA, Hideo²
BARKIN, Yury^{1*} ; HANADA, Hideo²

¹Sternberg Astronomical Institute, Moscow, Russia, ²National Astronomical Observatory of Japan, Mizusawa, Japan
¹Sternberg Astronomical Institute, Moscow, Russia, ²National Astronomical Observatory of Japan, Mizusawa, Japan

We have previously predicted and studied a step (abrupt) shift of the center of mass of the Earth in 1997 - 1998 years relatively to the mantle (Zotov, Barkin, Lubushin, 2009). In accordance with the basic provisions of the geodynamic model of excitation of planets and satellites shells (Barkin, 2002) we expected and we expect similar displacements of the centers of mass for other bodies in the solar system (for Mercury, Moon, Sun, Titan, Mars, etc.). Moreover, according to our hypothesis these abrupt geodynamic phenomena for solar system bodies are synchronous (Barkin, 2000) and, in particular, it should appear in 1997-1998. On the Earth, the similar jumps in 1997-1998 were observed in almost all planetary processes (Barkin, 2009). In the case of the Moon similar jump of center of mass obtains a confirmation in the data of laser observations and accounts for a specified period of time 1997-1998.

The jump (step) in the center of mass of the Moon in 1997 on data of laser ranging of reflectors on the lunar surface. On the basis of current laser measurements of distances to reflectors mounted on the Moon the preliminary estimates of the parameters of drift, oscillations and jump of the center of mass of the Moon were obtained. Their dynamic interpretation on the base of a geodynamic model of forced relative oscillations of the shells of planets and satellites has been done (Barkin, 2002). In the paper of G.A. Krasinskii (2003) from the analysis of lunar laser range measurements (or rather their residual differences compared with the theoretical celestial-mechanical design values of ranges) an abrupt (step) changes (in 1997 - 1998) in the coordinates of reflectors on the very substantial distances of about 15 -25 cm in selenographic coordinate system of the epoch have been discovered. Since jumps of coordinates for all four observed reflectors were quite close, it is natural to assume that the jump occurred in the position of the center of mass of the Moon by about 25-35 cm relatively to the lunar crust (in direction toward the Earth). Extremely important here is the fact that the jumps occurred in 1997-1998, as it was predicted by the theory of the unified geodynamic synchronous rhythms in variations of the activity of natural processes on the bodies of the solar system (Barkin, 2000). For the mean values of displacements of reflectors the following values were obtained (in meters): -0.15 +/- 0.04 m (offset along x coordinate - from the Earth), 0.23 +/- 0.07 m (offset on y - east), - 0.23 +/- 0.07 m (offset along z - to the north). Thus in 1997, the center of mass of the Moon abruptly shifted to a geographical point on the lunar surface with coordinates 40.00 N, 32.10 W approximately on distance in 0.36 +/- 0.11 m. According to the Krasinskii work (2003) we have identified trends in the changes of distances to reflectors and their abrupt changes before 1997 and after 1998, with rates of about 0.036 ns / year (before the jump) and at a rate of 0.128 ns / year (after the jump). If we consider only the drift relatively to the axis x, then estimates the drift velocities decrease: 0.98 cm / year - until 1997 and 1.47 cm / year - since 1998. It is expected to perform a spectral analysis of the residual differences of distances in order to identify their cyclic variations (with lunar months periods and with multiple periods).

キーワード: center of mass of the Moon, jumps and trends of center of mass of the Moon, LLR data
Keywords: center of mass of the Moon, jumps and trends of center of mass of the Moon, LLR data

Interpretation of unexplained secular changes of the lunar orbit Interpretation of unexplained secular changes of the lunar orbit

BARKIN, Yury^{1*}
BARKIN, Yury^{1*}

¹Sternberg Astronomical Institute, Moscow, Russia

¹Sternberg Astronomical Institute, Moscow, Russia

Unexplained secular effects in the orbital motion of the Moon are consequences of the observed phenomenon of remove of the center of mass of the Moon relatively to its mantle and crust toward the back-side. An explanation of anomalous part of secular variation in the longitude of the Moon and in the eccentricity of the lunar orbit has been obtained.

Unexplained secular variation of the eccentricity of the lunar orbit. In the works of James Williams and his colleagues showed that the observed rate of secular change of the eccentricity of the orbit of the Moon in 2.3×10^{-11} 1/year can not be explained within the framework of the classical model of the tides. Earth tides give only a fraction of the value specified in 1.3×10^{-11} 1/year and lunar tides result even effect with the opposite sign and give part of the acceleration in -0.6×10^{-11} 1/year. Remains unexplained an anomalous part of the secular change in the eccentricity (1.6 ± 0.4) $\times 10^{-11}$ 1/year. This value corresponds to abnormal changes in the distances to the perigee and apogee at 6 mm / year. "Abnormal speed distances to the perigee and apogee of the lunar orbit is up to 6 mm / year and its cause is unknown" (Williams J.,2006).

Tidal acceleration and evolution of the Moon's orbit. Laser ranging method proved to be very sensitive to the tidal acceleration of the Moon. Tides on the Earth dominate in the transfer of angular momentum, and energy in the orbital motion, in particular in the removal of the Moon from the Earth. Tidal effects on the Moon are separable from the effects of Earth tides in laser range measurements to the Moon (Chapront et al., 2002; Williams et al., 2009). Full tidal acceleration in the mean orbital longitude (due to the tides of the Earth and the Moon) is estimated at $-25.85''$ 1/cy², corresponding to the removal of the Moon from the Earth at a speed of 3.81 cm / year (Williams et al., 2009). The rate of secular variation of the eccentricity of the lunar orbit $e = (9 \pm 3) \times 10^{-12}$ 1/year also detected on the basis of long laser observations over a period of 38.7 years (March 16, 1970 - November 22, 2008) (Williams, Boggs, 2009). The basis of dynamical studies makes a precision lunar ephemeris DE421, taking into account all of Newtonian and Einsteinian effects. The authors believe that the study of the evolution of the lunar orbit is an important and surprisingly difficult task. Lunar laser ranging provides the numerical values for the two sources of dissipation on the Earth and the Moon.

Possible secular drift of the center of mass of the Moon relative to its crust and mantle toward the back side and an explanation of the anomalies of the orbital motion. In this report we give some first estimations of the possible rate of the secular drift of the Moon center of mass with respect to its crust and mantle in the 10 - 15 mm / year toward the back-side of the satellite. This secular drift of the center of mass of the Moon should be considered by the studying of the orbital motion of the Moon on laser-based observations. Namely, to add to the value obtained by laser observations. The result will be an estimate of the secular increasing of semi-major axis is the center of mass of the Moon. It should be expected that this will obtain the interpretation and explanation of the unexplained part of the secular acceleration of the Moon orbit and the anomalous part of the secular variation of the eccentricity of the lunar orbit, identified according to the perennial laser observations of the Moon. An anomalous part of the orbital acceleration (unexplained) of the Moon is $0.7''$ / cy², and the anomalous part of the secular variation of the eccentricity is characterized by rate in 1.23×10^{-11} 1/year (Williams et al., 2011). Found offset - drift of the center of mass of the Moon (12 - 15 mm / year) is explained by the mechanism of excitation and the relative displacements of the shells of the Moon (solid core, liquid core, mantle) (Barkin, 2002).

キーワード: anomalous secular variation of the eccentricity of the lunar, tidal and non-tidal acceleration of the Moon, the center of mass of the Moon drift

Keywords: anomalous secular variation of the eccentricity of the lunar, tidal and non-tidal acceleration of the Moon, the center of mass of the Moon drift

アポロ月震計データと最新の測月データから推定される月深部構造 Deep interior structure of the Moon inferred from Apollo seismic data and the latest se- lenodetic data

松本 晃治^{1*}; 山田 竜平¹; 菊池 冬彦¹; 鎌田 俊一²; 岩田 隆浩³; 石原 吉明³; 花田 英夫¹; 佐々木 晶⁴
MATSUMOTO, Koji^{1*}; YAMADA, Ryuhei¹; KIKUCHI, Fuyuhiko¹; KAMATA, Shunichi²; IWATA, Takahiro³; ISHIHARA,
Yoshiaki³; HANADA, Hideo¹; SASAKI, Sho⁴

¹ 国立天文台 RISE 月惑星探査権糖質, ² カリフォルニア大学サンタクルーズ校, ³ 宇宙航空研究開発機構, ⁴ 大阪大学
¹RISE Project Office, NAOJ, ²University of California Santa Cruz, ³JAXA, ⁴Osaka University

Internal structure and composition of the Moon provide important clue and constraints on theories for how the Moon formed and evolved. The Apollo seismic network has contributed to the internal structure modeling. Efforts have been made to detect the lunar core from the noisy Apollo data (e.g., [1], [2]), but there is scant information about the structure below the deepest moonquakes at about 1000 km depth. On the other hand, there have been geodetic studies to infer the deep structure of the Moon. For example, LLR (Lunar Laser Ranging) data analyses detected a displacement of the lunar pole of rotation, indicating that dissipation is acting on the rotation arising from a fluid core [3]. Bayesian inversion using geodetic data weakly suggests a fluid core and partial melt in the lower mantle region [4]. Further improvements in determining the second-degree gravity coefficients and the Love numbers will help us to better constrain the lunar internal structure.

Recent analyses of GRAIL data have achieved the improved k_2 accuracy; JPL solution is 0.02405 ± 0.00018 [5], and GSFC solution is 0.02427 ± 0.00026 [6]. The two solutions are consistent with each other within their error bounds, and the accuracy of k_2 is now about 1 %. By introducing the improved gravity coefficients and k_2 from GRAIL mission, the updated LLR data analysis has also resulted in a better h_2 determination. Such accurately-determined Love numbers will contribute to constrain the structure of the lunar deep interior, such as the radius of the possible liquid core. It is difficult, however, to tightly constrain the internal structure from the geodetic data only because there are trade-offs among the structures of crust, mantle, and core. The combination of the Apollo seismic data and the geodetic data therefore afford the key to better determination of the lunar interior structure. We included geodetic data of the mass, the mean moment of inertia, the Love numbers h_2 and k_2 , and 262 P and S travel time data in the analysis.

Markov Chain Monte Carlo (MCMC) method is used to infer the model parameters. When we used a five-layer model consisting of crust, upper-mantle, mid-mantle, lower-mantle, and core, the core radius is estimated to be 483 ± 22 km, and the core density values tend to be sampled around the assumed lower limit of 3600 kg/m^3 . However, the inferred core radius is significantly larger than the magnetic constraint from SELENE data [7] which predicts the upper bound of the core radius to be 400 km. This discrepancy might be attributed to a possible low velocity layer above the core-mantle boundary which was not included in the five-layer model. We will discuss the results when such a low velocity layer is taken into account.

- [1] Weber et al. (2011), *Science*, 331, 309-312, doi:10.1126/science.1199375
- [2] Garcia et al. (2011), *PEPI*, doi:10.1016/j.pepi.2011.06.015
- [3] Williams et al. (2001), *JGR*, 106, E11, 27,933-27,968
- [4] Khan and Mosegaard (2005), *GRL*, 32, L22203, doi:10.1029/2005GL023985
- [5] Konopliv et al (2013), *JGR*, 118, doi:10.1002/jgre.20097
- [6] Taken from the PDS label of GRAIL Derived Data Products
- [7] Simizu et al. (2013), *Icarus*, 222, doi:10.1016/j.icarus.2012.10.029

キーワード: 月, 内部構造, 重力場, 潮汐ラブ数, グレイル

Keywords: Moon, internal structure, gravity field, tidal Love number, GRAIL

月面での回転観測：可能性と問題点 Observations of lunar rotation on the Moon: possibility and problems.

花田 英夫^{1*}; 鶴田 誠逸¹; 浅利 一善¹; 千葉 皓太²; 稲葉 健太²; 船崎 健一²; 佐藤 淳²; 谷口 英夫²; 加藤 大雅²; 菊池 護²; 荒木 博志¹; 野田 寛大¹; 鹿島 伸悟¹
HANADA, Hideo^{1*}; TSURUTA, Seiitsu¹; ASARI, Kazuyoshi¹; CHIBA, Kouta²; INABA, Kenta²; FUNAZAKI, Ken-ichi²; SATOH, Atsushi²; TANIGUCHI, Hideo²; KATO, Hiromasa²; KIKUCHI, Mamoru²; ARAKI, Hiroshi¹; NODA, Hiroto¹; KASHIMA, Shingo¹

¹ 国立天文台 RISE 月惑星探査検討室, ² 岩手大学工学部

¹RISE Project, National Astronomical Observatory, ²Faculty of Engineering, Iwate University

The lunar rotation is one of the essential and basic target of selenodetic observations for investigation of the interior of the Moon as well as those of gravity fields, and high accuracy of the observations have a potential to detect signals related to the structure of lunar deep interior including the core. We have developed a small telescope like a PZT (Photographic Zenith Tube) for observations of Lunar rotation with the target accuracy of 1 milli-seconds of arc (1 mas)[1]. Theoretical investigation shows that observations by the telescope in the polar area of the Moon will open great possibilities for determining the libration in inclination ρ and node $I\sigma$ with the accuracy much better than before, although the determination of the libration in longitude will not be very well. It also showed that the determination error in the libration angles will not exceed $\sqrt{2\varepsilon}$, where ε is the positioning error of stars and is regarded as 1 milli-seconds of arc [2].

There are several technical problems to be solved in the development of the telescope. Effect of large temperature change is one of the most serious problem for such a precise observation, and we can loosen thermal condition by about ten times by introducing a diffraction lens compared with the case not introducing it. It is possible, on the other hand, that the vibrations of the mercury surface caused by the ground vibrations lead to fluctuations of star positions on CCD as large as 1 second of arc judging from laboratory experiments. The amplitude of the fluctuations depend on the amplitude of the ground vibrations and the depth of mercury pool. We can reduce the effect of the vibrations by making the mercury pool shallow down to the minimum depth. In the case of the mercury pool of 64mm diameter, the depth of 0.5mm is the best according to our experience [3]. It is important to keep the proper period of the mercury pool away from the period of ground vibrations in order to avoid the resonance. It is also effective to lengthen the integration time, and it can improve the reliability of the mean value of the center of a star image by statistical procedure.

We have already made a bread board model (BBM) and we will observe the deflection of the vertical on the ground by using the BBM for the time being in order to evaluate the characteristics of the total system of the telescope.

References

- [1] Hanada, H. et al., Development of a digital zenith telescope for advanced astrometry, Science China, 55, 723-732, 2012.
- [2] Petrova, N. and H. Hanada, Computer simulation of observations of stars from the Moon using the polar Zenith Telescope of the Japanese Project ILOM, Solar Sys. Res., 47, 504-517, 2013.
- [3] Tsuruta, S. et al., Stellar imaging experiment using a mercury pool as a ground test of the telescope for In-situ Lunar Orientation Measurements(ILOM), Proc. 14th Space Science Symposium, 2014.

キーワード: 回転, 月, 望遠鏡, PZT, 物理ひょう動

Keywords: rotation, moon, telescope, PZT, librations

THE EARTH ORIENTATION PARAMETERS AND THE VARIATION OF THE SECOND ZONAL HARMONIC OF THE GEOPOTENTIAL THE EARTH ORIENTATION PARAMETERS AND THE VARIATION OF THE SECOND ZONAL HARMONIC OF THE GEOPOTENTIAL

BARKIN, Mikhail^{1*} ; FILIPPOVA, Alexandra¹ ; NOVIKOVA, Daria¹ ; PEREPELKIN, Vadim¹
BARKIN, Mikhail^{1*} ; FILIPPOVA, Alexandra¹ ; NOVIKOVA, Daria¹ ; PEREPELKIN, Vadim¹

¹Moscow Aviation Institute, Moscow, Russia

¹Moscow Aviation Institute, Moscow, Russia

The study of the time variations of the geopotential as a result of the rotary-oscillatory processes of the Earth motion is of a significant natural-sciences and practical interest. Oscillations of the Earth's inertia tensor components depend on many factors, among them the mechanical and physical parameters of the planet, the motions of tide-forming bodies, and the observed large-scale phenomena in nature. Time-dependent variations of these and other factors (regular and irregular oscillations, stochastic fluctuations, secular variations) affect the Earth rotary-oscillatory processes and the rotational parameters of the planet. The dynamic processes of the Earth orientation parameters (EOP) in turn have an effect on its figure and lead to the fluctuations of the gravitation field. Observed variations of the EOP, the variations of the Earth's gravitational field and oscillations in the large-scale geophysical events appear to be in a considerable correlation.

An amplitude-frequency analysis of the rotary-oscillatory Earth motion under the action of gravitational-tidal perturbing torques from the Sun and the Moon is carried out using the classical mechanics' methods. The simulation results of the oscillatory process in the motion of the Earth pole and the variations of the second zonal harmonic of the geopotential are studied. Based on the dynamic Euler-Liouville equations expressions for amplitude and phase of the Earth pole oscillations are obtained. A comparison of the spectral power densities of the time series between the Earth pole coordinates and the variations of the geopotential is carried out. A functional dependence of the aforementioned component of the geopotential from the amplitude and phase of the Earth's pole oscillatory process is shown.

キーワード: the rotary-oscillatory processes, secular variations, stochastic fluctuations, geopotential

Keywords: the rotary-oscillatory processes, secular variations, stochastic fluctuations, geopotential

The free and forced librations of the Moon with liquid shell and solid core The free and forced librations of the Moon with liquid shell and solid core

BARKIN, Yury^{1*}; HANADA, Hideo²; MATSUMOTO, Koji²; BARKIN, Mikhail³
BARKIN, Yury^{1*}; HANADA, Hideo²; MATSUMOTO, Koji²; BARKIN, Mikhail³

¹Sternberg Astronomical Institute, Moscow, Russia, ²National Astronomical Observatory of Japan, Mizusawa, Japan, ³Moscow Aviation Institute, Moscow, Russia

¹Sternberg Astronomical Institute, Moscow, Russia, ²National Astronomical Observatory of Japan, Mizusawa, Japan, ³Moscow Aviation Institute, Moscow, Russia

In report we present our results of the study of lunar physical libration of the Moon on the base of its two and three layers models. On the base of analytical solution for two layers model (the Moon with liquid core) and empirical theory of the Moon's rotation (Rambaux, Williams, 2011), we have identified period, amplitude, and the initial phase of the fourth mode of free libration of the Moon, caused by liquid ellipsoidal core. Preliminary results of studies of three-layers model of physical librations of the Moon have been obtained on the base of some simplified approach for the problem of rotation of the Moon with liquid and rigid cores. The plans for future studies of the Moon rotation are discussed.

The modern view of internal structure of the Moon planet takes into account a complex two- or three-layer model. In our work the analytical theory of lunar physical libration based on its two-layer model consisting of a non-spherical solid mantle and of the ellipsoidal liquid core has been developed. The Moon moves on high-accurate perturbed orbit in the gravitational field of the Earth and other celestial bodies. On the base of two layers model of the Moon we have fulfilled systematic studies of the Moon physical librations. And in first we have presented a solution of the problem in components of vector of angular velocity of the Moon. An analytical presentation of LOD of the Moon with high accuracy in form of trigonometric series has here the progressive value. In first we have determined the fourth mode of free libration of the Moon caused by the influence of the liquid core oscillations of pole axis of rotation of the Moon (its mantle), with a long period in 205.7 yr, with an amplitude of $0''$ 0395 and the initial phase of -134° (for the initial epoch 2000.0). This oscillation reflects the so-called phenomenon of free oscillation of the liquid core. The estimates for the dynamic (meridional) oblatenesses of the ellipsoidal liquid core of the Moon: 0.000442 and 0.000283 have been obtained. These fundamental parameters of geodynamics of the Moon could be determined only on the base of data of observations. Earlier the attempts to determine the period of free core nutation undertaken. Our results were obtained by comparing of the developed analytical theory of lunar physical libration with empirical theory libration of the Moon, constructed on the basis of laser observations in last about 40 years (Rambaux, Williams, 2011).

Preliminary results of studies of three-layers model of physical librations of the Moon have been obtained on the base of some simplified approach for the problem of rotation of the Moon with liquid and rigid cores. We have analyzed the Cassini's motion of the decoupled solid core and its librations in longitude to compare with the Moon motions. On the base of Getino, Ferrandiz et al. approach we give estimations of the periods of free librations of this system. We have constructed differential equations of rotational motion of three layers Moon from positions of the Hamiltonian formalism with application of Andoyer's and Poincaré's variables. Now we construct analytical theory of rotation of the Moon system consisting from the non-spherical mantle, ellipsoidal liquid core and solid core.

キーワード: Moon rotation, free libration, liquid core, solid core

Keywords: Moon rotation, free libration, liquid core, solid core

地球の大規模な真の極移動の速度の粘性構造依存性：地球マントル内部の低粘性領域が緩和モードの時定数に及ぼす潜在的影響の評価 Viscosity structure dependence of large-scale polar wander rate of the Earth: A potential impact of a low-viscosity zone

原田 雄司^{1*}; 肖 竜¹
HARADA, Yuji^{1*}; XIAO, Long¹

¹ 中国地質大学地球科学学院行星科学研究所
¹China University of Geosciences

地球の真の極移動の速度に与えるマントル内部の低粘性層の影響に関しては、それ程には多くないが幾つかの研究が行なわれている。種々の地球物理学的制約から、地球マントル、特に最上部や最下部において著しく粘性の低い層が存在する、という可能性が指摘されている。このような極端な粘性の差異は極移動にも或る程度の影響を及ぼす、という事が上述の先行研究で調べられている。

但し、これらの研究で取り扱われていたのは比較的小規模な極移動である。それらのモデル計算は主として後氷期回復による質量再分配に伴う極移動を想定していた。この場合、極移動の典型的な時間スケールは地質学的時間スケールと比べて短く、概ねマントルの粘弾性変形の特徴的時間スケールを反映する。そのような極移動の振幅は大きくても数度程度であり、線形近似された極運動方程式を適用するのが定石である。その一方、今の所、数十度程度の大規模な極移動に対する低粘性層の潜在的影響について議論された研究例は無い。

このような地球内部における粘性の不均質性の影響を検討する事は、観測量から導かれた極移動のシナリオ、特にその時間変化の妥当性を定量的に評価する上で重要であると考えられる。実際の地球の太古の極移動は古地磁気を初めとする地質学的状況証拠から推定されているが、この種の筋書きは上記のようなモデル計算に基づいて初めて物理的に解釈され得る。そのような変遷が起こり得る力学的条件を考察する事によって、当時の粘性構造を制約する観点において有益な情報が得られると期待される。こうした議論をする為には前述のような物理的に特殊な層の効果も重視すべきかもしれない。

そこで本研究では、地球マントル内部における低粘性層の存在が極移動の時間スケールに及ぼす影響について定量的評価を試みた。特に地球内部の力学的応答を特徴付ける粘弾性ラブ数のモデル計算を実施し、ラブ数に含まれる複数の緩和モードの強度と時間スケールの粘性構造に対する依存性について調査した。更にそれらの緩和モードの構造依存性を極移動速度の構造依存性と比較した。尚、数値計算の都合上、緩和モードの算出に際しては地球の層構造を簡略化し、かつ非圧縮性を仮定した。

本計算においては、極運動方程式を非線形のまま積分可能とする準流体近似を適用した。その根拠は、ここで取り扱うような大規模な極移動の場合には線形近似を使えないからである。この準流体近似の適用範囲に従い、ここではアセノスフェアの粘弾性変形の特徴的時間スケールよりも遅い荷重の進化を仮定した。

上述の計算の結果、極移動の時間スケールはほぼ最長期の緩和モードにのみ依存している事が分かった。ここで特筆すべき点は、この極移動において支配的な緩和モードの強度が粘弾性ラブ数全体に占める比率は、実は余り大きくないという事である。それは換言すれば、潮汐変形の振幅がより卓越する他のモードは、極移動の時間スケールに対しては殆んど影響しない事を意味する。これは一見すると奇妙な結果に感じられるかもしれない。

このような依存性を示す理由は、前述した最長期のモードの時間スケールだけが他のモードと比較して桁違いに長いからである。緩和の時定数が長いモードは、たとえ振幅が小さくても長期的に自転軸を安定化する作用を有する。逆に振幅が大きくても時定数が短いモードでは速やかに緩和する為、自転の長期的安定性に対する寄与は小さい。

この結果を踏まえるならば、極移動速度の構造依存性も概ね、この最長期のモードの緩和時間の依存性のみ反映していると言える。実際には、たとえマントル内部の低粘性層を含まない内部構造を仮定しても、このモードの影響は依然として支配的である。そして低粘性層が存在する場合、その粘性が低くなると、それに伴ってこの最長モードの時間スケールも短くなる。最も短い場合では四割にも満たない。但し粘性が或る値よりも低くなると、このモードの時間スケールは一定の値に漸近する。この傾向の理由は、この層の粘性が低過ぎる為、十分に長い時間スケールでは実質的に粘弾性体ではなく流体として振る舞うからである。

以上の計算結果から得られる結論は、一般に地球内部の低粘性層の存在は大規模極移動の時間スケールを短くする事、そしてその影響は主に最長期の緩和モードの時間スケールの変化に起因する事である。確かに地球の大規模極移動の時間スケールの内部構造依存性に関しては既に既存の研究でも議論されている。しかし以前の研究ではマントルの粘性構造が単純化され、その中の低粘性層の影響について検討されていなかった。かつ極移動速度を強く支配している要因も明確には指摘されていなかった。それに対して今回の研究では、この層の影響を明示的に含む極移動の時間スケールを見積もり、不均一な粘性構造が大規模極移動に与える影響が無視出来ない事を示した。

PPS03-P02

会場:3 階ポスター会場

時間:4 月 29 日 18:15-19:30

キーワード: 真の極移動, 地球, マントル, 低粘性層, 緩和モード, 時定数

Keywords: true polar wander, the Earth, mantle, low-viscosity layer, relaxation mode, time constant

Effects of global geodynamics in a series of astrometry observations of latitude at Poltava

KHALYAVINA, Lydmila¹ ; BARKIN, Yury^{2*}
KHALYAVINA, Lydmila¹ ; BARKIN, Yury^{2*}

¹Gravimetry Observatory, Poltava, Ukraine, ²Sternberg Astronomical Institute, Moscow, Russia

¹Gravimetry Observatory, Poltava, Ukraine, ²Sternberg Astronomical Institute, Moscow, Russia

Diverse geodynamic phenomena observed in the modern era, received a convincing explanation in the framework of the northern drift of Earth's core. Model proposed and developed by Yuri Barkin relative to the set of ancient geodynamic processes: the secular drift of the Earth's pole, non-tidal acceleration of the Earth's rotation, secular change of gravity, the evolution of the earth's figure, plate tectonics, the formation of specific geological structures, etc.

The North drift of the core generates mass redistribution of the Earth and leads to changes in the gravitational field. Since astrometry instruments have as a reference axis direction of the local plumb line, then this process should be displayed in the slow position changes no polar zenith Observatory. It is shown that for locations in the northern hemisphere, the north drift of the core causes the displacement of local plumb in a southerly direction. Is the picture of long-term changes in the direction of gravity (NST) in the meridian of Poltava for the period 1962 - 2013 based on long-term observations of latitude prismatic astrolabe taking into account: 1) high-precision catalogs (HIPPARCOS, ARIHIP, Tycho-2), 2) improved model of the pole C01 IERS; 3) the theory of the precession-nutation IAU2000; 4) plate tectonics (NUVEL1A-NNR). The resulting long-period changes in NST can be represented as the sum of three components: a linear trend with velocity $\sim -0.0003''/\text{yr}$, the nonlinear part of the trend, consisting of two branches (descending - in 1962 to 1996. And rising - in 1998 and 2010.), which can be regarded as a fragment of a wave with period $T \sim 50$ years and amplitude $A \sim 0.02''$; quasi cyclic part with 11 - year period, close to the main period of index of solar activity period and amplitude $< 0.01''$.

The linear part of the translational displacement means the plumb line to the south of Poltava, which is consistent with the above Barkin's model. The observed velocity of motion of zenith corresponds to moving the center of mass of the Earth in a northerly direction at the velocity in 1.4 cm/yr. Found that the nonlinear part of the trend and the 11-year cyclicity in the shifts of plummet quite clearly reproduce the form of low-frequency polar latitude variations at Poltava derived from model C01 (EOP IERS). Actually observed amplitude of long-period oscillations of latitude caused by pole motion, in 2 times higher than the calculated amplitude. The non-linear part of the trend is the projection on the Poltava meridian of the Markowitz wave.

It is shown that both low-frequency cycles are negatively correlated with the corresponding components of the index of solar activity. The most probable mechanism of solar activity influence on the motion of the pole is the North Atlantic Oscillation. An increase in the amplitude of low-frequency polar displacements of Poltava zenith in astrometric observations requires a special study. One from possible explanations - the influence of the features of the geological structure in the vicinity of Poltava, which is located in the center of the so-called rift Poltava site.

キーワード: plate tectonics, secular change of gravity, Markowitz wave

Keywords: plate tectonics, secular change of gravity, Markowitz wave

Minerals detection on Mars from Mars Reconnaissance Orbiter (MRO) CRISM data Minerals detection on Mars from Mars Reconnaissance Orbiter (MRO) CRISM data

Jin Shuanggen¹ ; Barkin Yury^{2*}
JIN, Shuanggen¹ ; BARKIN, Yury^{2*}

¹ 上海天文台, ² モスクワ大学

¹Shanghai Astronomical Observatory, Chinese Academy of Science, ²Sternberg Astronomical Institute, Moscow State University

Martian mineral detection and mapping can provide important information and constraints on Martian aqueous history, which can be used to assess the potential habitability of Mars. Degrees of addressing the key question for Martian aqueous alteration are dictated by the depth and extent of grasping the Martian hydrous mineral. Therefore, it is important to know detailed minerals and chemical induction of the existence of water on the Martian surface at past or present. In-situ observations of the Martian rovers, such as Spirit, Opportunity and Curiosity have provided the mineralogical analysis of Martian surface, but restricting in a limited areas. Compact Reconnaissance Imaging Spectrometer for Mars (CRISM) aboard the Mars Reconnaissance Orbiter (MRO) with enhanced spectral resolution can provide more information at spatial and time scale. In this paper, CRISM near-infrared spectral data are used to identify mineral classes and distribution at Martian Gale region, including kaolinite, chlorites, smectite, jarosite, northupite and salts. The detection of northupite that is indicative of evaporation in Gale region suggests that the Gale region has experienced the climate change from moist condition with mineral dissolution to dryer climate with water evaporation.

キーワード: Martian minerals, Mars Reconnaissance Orbiter, CRISM
Keywords: Martian minerals, Mars Reconnaissance Orbiter, CRISM