

## The Van, Turkey Earthquake of October 2011: Seismicity, Mechanism and its Aftershocks

KALAFAT, Dogan<sup>1\*</sup>  
KALAFAT, Dogan<sup>1\*</sup>

<sup>1</sup>Bogazici University Kandilli Observatory and ERI Cengelkoy, Istanbul Turkey

<sup>1</sup>Bogazici University Kandilli Observatory and ERI Cengelkoy, Istanbul Turkey

On 23 October 2011, a strong earthquake ( $M_w=7.1$ ) occurred east of Van Lake. The earthquake destroyed damage along the Van Fault Zone. Generally, it is caused significant damage in the city of Van with Ercis town, as well as in many villages. The epicenter of the main shock was located in Tabanlı Village between Van city and Ercis town. Shortly afterwards the November 9, 2011 earthquake ( $M_w=5.6$ ) occurred southeastern part of Van Lake, Edremit town area. The main shock and second shocks caused significant damage and deaths of 644 people.

After the main shock 11 important earthquakes ( $5.0 \leq M \leq 6.0$ ) were occurred in the region which has hypo-central distances of 8-38 km. from the main shock location. The main shock triggered mass movement, spreading, and local liquefaction. The important earthquakes and the aftershocks distribution showed that the E-W and NE-SW oriented fault segments caused the earthquake activities. The distribution of the aftershocks supported the presence a rupture of approximately  $70 \pm 10$  km. Aftershocks occurred within an area of approximately 2300 km<sup>2</sup>.


Fault mechanism solution has supported 23 October 2011 Van Earthquake was a reverse fault with a northward dipping fault plane. The fault plane solutions of about 225 important earthquakes ( $M \geq 4.0$ ) were calculated using MT inversion solution technique by this study.

In addition, the stress tensor analysis was completed using the focal mechanism solutions. The stress tensor values of the region were calculated using the azimuth and plunge pairs of P and T axes for 225 earthquakes. The maximum principal stress (P-compressional) of the main shock is aligned in an N-S (NNW/SSE) direction and the tensional axis (T-dilatation) is aligned in an E-W (ENE-WSW) direction. The results of the focal mechanism and stress analysis solutions show that earthquakes have occurred on the reverse faulting and the seismic activity has been continuing under the compressional regime in the region. The Van earthquake activity initiated and caused an increase in seismic activity of the region. This may be explained by the triggering of small faults by the Van Earthquake. The Van earthquake is a good example of compressional deformation and the activity of blind reverse faulting.

This study was supported by Bogazici University Research Projects Commission under SRP/BAP project No. 6671.

キーワード: Van Lake, aftershocks distribution, focal mechanism, stress analysis, reverse faulting

Keywords: Van Lake, aftershocks distribution, focal mechanism, stress analysis, reverse faulting



## Microstructural observation of quartz and K-feldspar in quartzo-feldspathic granulite in Sri Lanka Microstructural observation of quartz and K-feldspar in quartzo-feldspathic granulite in Sri Lanka

Athurupana Bhatthiya Madhura Bandara<sup>1\*</sup>; 福田 惇一<sup>1</sup>; 武藤 潤<sup>1</sup>; 長濱 裕幸<sup>1</sup>  
ATHURUPANA, Bhatthiya madhura bandara<sup>1\*</sup>; FUKUDA, Jun-ichi<sup>1</sup>; MUTO, Jun<sup>1</sup>; NAGAHAMA, Hiroyuki<sup>1</sup>

<sup>1</sup>Department of Earth Science, Tohoku University, Japan

<sup>1</sup>Department of Earth Science, Tohoku University, Japan

Sri Lankan basement has been considered as high grade terrains which have suffered poly phase deformation and also upper amphibolite to granulite facies metamorphism during Pan-African amalgamation. In order to illuminate the rheological properties of felsic granulite in deep crustal level associated with high grade metamorphism, microstructural and petrographic observations were conducted.

A quartzofeldspathic gneiss sample (Quartz~40%, K-feldspar~55% and plagioclase <5%) contains highly elongated ribbon quartz which demarcate lineation, in K-feldspar matrix. The sample was collected from the hinge zone of kilometer scale large upright cusped antiform between two large synforms named Dumbara and Huluganga in Sri Lanka. The lineation as maximum elongation direction is N35°W/horizontal. Thin sections were made parallel and perpendicular to the lineation for petrographic and microstructural observations. Based on energy dispersive X-ray spectroscopic (EDS) measurement, most of K-feldspars in the matrix are sanidine [(K<sub>0.59</sub>Na<sub>0.41</sub>)AlSi<sub>3</sub>O<sub>8</sub>] composition. Plagioclase is albite (An 0.1) composition and occurs in clusters and exsolution lamellas in K-feldspar. The grain size of sanidine is from ~50 μm to 2 mm. Large sanidine porphyroclasts are partly recrystallized and show core and mantle structures. Grain boundaries of sanidine are sub polygonal to amoeboid shapes with increasing grain size, while plagioclase grains in clusters show perfect polygonal boundaries with grain size ranging from 50 to 200 μm. Around the plagioclase clusters, scapolite reaction corona exists with crosscutting elongated ribbon quartz. Sericite presents in the same region as very low temperature alteration of plagioclase and scapolite. Most importantly ribbon quartz grains are boudinaged in K-feldspar matrix. Any sign of undulatory extinction or dynamic recrystallization is absent in both boudinaged and large ribbon quartz. Measured average axial ratio of ribbon quartz grains is 21:7:1 with respect to X, Y and Z directions.

The crystallographic orientations of both quartz and sanidine were measured by electron backscattered diffraction (EBSD) method, including large ribbon quartz and boudinaged quartz and also matrix sanidine. The lattice preferred orientation (LPO) patterns of sanidine manifest (010)<001> as a dominant slip system with a minor activation of (010)<100> slip system. The quartz LPO indicates the activations of prism<a> and rhomb<a> slip systems.

According to the presence of K-feldspar as sanidine and their slip systems, we can point out that the sample has prevailed high grade conditions (~550-800°C and ~0.4-0.8 GPa) during deformation [e.g. Menegon et al. 2008]. Scapolitization which occurs at granulite facies could be indicative of peak metamorphism. Also, reaction coronas of scapolite crosscutting ribbon quartz can be interpreted as strong deformation prior to the peak metamorphism. Exsolution of K-feldspar indicates the post dated cooling relative to the strong deformation and the peak metamorphism.

### Reference:

Menegon, L., Pennacchioni G., Spiess R., 2008. Dissolution-precipitation creep of K-feldspar in mid-crustal granite mylonites. *Journal of Structural Geology* 30(5): 565-579

キーワード: Microstructures, Quartz, K-feldspar, Quartz micro boudins

Keywords: Microstructures, Quartz, K-feldspar, Quartz micro boudins

## Multiple events of metamorphism in lenses of eclogite within marbles of Maykhan Tsakhir Formation, Lake Zone, Mongolia

### Multiple events of metamorphism in lenses of eclogite within marbles of Maykhan Tsakhir Formation, Lake Zone, Mongolia

JAVKHLAN, Otgonkhuu<sup>1\*</sup>; TAKASU, Akira<sup>1</sup>; BAT-ULZII, Dash<sup>2</sup>; KABIR, Md fazle<sup>1</sup>  
JAVKHLAN, Otgonkhuu<sup>1\*</sup>; TAKASU, Akira<sup>1</sup>; BAT-ULZII, Dash<sup>2</sup>; KABIR, Md fazle<sup>1</sup>

<sup>1</sup>Department of Geoscience, Shimane University, Japan, <sup>2</sup>School of Geology and Petroleum, Mongolian University of Science and Technology, Mongolia

<sup>1</sup>Department of Geoscience, Shimane University, Japan, <sup>2</sup>School of Geology and Petroleum, Mongolian University of Science and Technology, Mongolia

The eclogite-bearing Alag Khadny metamorphic complex in the Lake Zone, SW Mongolia located in the central part of the Central Asian Orogenic Belt, consist mainly of orthogneisses which interleaving with marbles including lenses of garnet-chloritoid schists of Maykhan Tsakhir Formation. Eclogites have two modes of occurrence, i.e. lenses and boudins *eclogite-1* in orthogneisses and *eclogite-2* in marbles. Thermocalc calculations for the peak eclogite facies metamorphism for *eclogite-1* with the assemblage of Grt + Omp + Brs ± Ph ± Ep yielded 570-630 °C and 22-25 kbar (Javkhan et al., 2013a). In contrast, pressure conditions of the garnet-chloritoid schists (10-11 kbar) are distinctly lower than those of the eclogite-1, whereas temperatures (560-590 °C) are similar (Javkhan et al., 2013b).

*Eclogite-2* in marbles consists of small grains of garnet (<0.1 mm) and omphacite with minor amounts of amphibole, epidote, paragonite, plagioclase, chlorite, calcite, biotite, quartz, titanite and rutile. The matrix of *eclogite-2* shows a pseudomorphous texture, where small grains of garnet crowd cemented by titanite forming isomorphous round shape. Some of cores of garnet grain contain relics of garnet ( $X_{Ca}=0.32-0.42$ ;  $X_{Mg}=0.06-0.08$ ) indicating previous mineral were larger porphyroblastic garnet. In addition, small grains of omphacite forming rectangular prismatic nature surrounded by garnet grains.

Garnet grains have compositionally zoning with core ( $X_{Ca}=0.08-0.20$ ;  $X_{Mg}=0.10-0.16$ ), mantle ( $X_{Ca}=0.24-0.39$ ;  $X_{Mg}=0.08-0.17$ ), rim ( $X_{Ca}=0.22-0.26$ ;  $X_{Mg}=0.18-0.23$ ) and outer-rim ( $X_{Ca}=0.20-0.22$ ;  $X_{Mg}=0.12-0.18$ ). Few omphacites preserved their core ( $X_{Jd}=0.27-0.31$ ;  $Fe_2O_3=1.34-2.22$  wt%) whereas most of grains have compositional heterogeneity with  $X_{Jd}$  from 0.34 to 0.48 ( $Fe_2O_3=0.04-2.31$ ) and locally with rims of higher  $Fe_2O_3$  (3.79 wt%)-bearing omphacite ( $X_{Jd}=0.32$ ). Omphacites partially replaced by symplectites of Pl (An=15-17), amphibole (Ed, Act, Mg-hbl) and Ep. Three types of amphibole are distinguished by their texture, amphibole (Amp1) [zoned with Act ( $X_{Mg}=0.79-0.81$ ) core, Brs ( $X_{Mg}=0.54-0.69$ ) mantle and rims with Ts, Mg-Trm and Prg in compositions] coexisting with Grt and Omp, poikiloblastic barroisitic amphibole (Amp2) ( $X_{Mg}=0.65-0.75$ ) containing eclogitic minerals of Grt and Omp with their symplectitic assemblage and finally actinolitic amphiboles (Amp3) partially replacing Omp and Grt.

Based on the textures we distinguished two metamorphic events, i.e. eclogite facies metamorphism and poikiloblastic barroisitic amphibole metamorphism. The peak eclogite facies metamorphism characterized by assemblages of Grt (mantle) + Omp + Amp1 (Brs) + Ep + Pg + Rt. Thermocalc calculation yielded  $487 \pm 46$  °C and  $19.7 \pm 2.1$  kbar (sigfit=1.80). Thermocalc calculation of Grt (rim) + Omp (rim) + Amp1 (Mg-Trm) + Ep + Pl yielded  $666 \pm 45$  °C and  $13.7 \pm 1.6$  kbar (sigfit = 2.11) suggesting a decompression stage after the eclogitic metamorphism. The poikiloblastic barroisitic Amp2 shows decreasing Si (7.01-6.69 pfu) and increasing NaB (0.61-0.70 pfu) from core to rim, suggesting that the Amp2 grew after the peak eclogite facies metamorphism, and probably during the second prograde metamorphic event. Approximate P-T conditions of the poikiloblastic barroisitic Amp2 are estimated as 5-7 kbar at c. 450 °C.

<sup>40</sup>Ar/<sup>39</sup>Ar muscovite ages for eclogites ( $543 \pm 3.9$  Ma) in marbles (probably eclogite-2) and the garnet-chloritoid schists ( $537 \pm 2.7$  Ma) were determined (Stipska et al. 2010). K-Ar ages for eclogite-1 [ $603 \pm 15$  Ma,  $602 \pm 15$  Ma (Amp) and  $612 \pm 15$  Ma (Ph)] within orthogneisses have been obtained (Javkhan et al., 2014). These ages are interpreted as the exhumation ages for the eclogites (-1 and -2) and the garnet-chloritoid schists.

The peak temperature conditions of eclogite-2 considerably lower than eclogite-1 whereas the pressure conditions are similar. The peak P-T conditions garnet-chloritoid schists are correlated with the poikiloblastic Amp2 metamorphism of the eclogite-2.

キーワード: eclogite-2, pseudomorphous texture, garnet-chloritoid schists, Maykhan Tsakhir Formation, Lake Zone, SW Mongolia

Keywords: eclogite-2, pseudomorphous texture, garnet-chloritoid schists, Maykhan Tsakhir Formation, Lake Zone, SW Mongolia

---

SCG08-03

会場:311

時間:4月29日 09:45-10:00

lia

## チェコ共和国ボヘミア山塊のザクロ石橄欖岩が経験した多重上昇履歴について Multiple exhumation episodes recorded in orogenic garnet peridotites from the Bohemian Massif (Czech Republic)

苗村 康輔<sup>1\*</sup>; スフォイッカ マルティン<sup>2</sup>; アッカーマン ルカーシュ<sup>2</sup>; 清水 以知子<sup>1</sup>; 平島 崇男<sup>3</sup>  
NAEMURA, Kosuke<sup>1\*</sup>; SVOJTKA, Martin<sup>2</sup>; ACKERMAN, Lukas<sup>2</sup>; SHIMIZU, Ichiko<sup>1</sup>; HIRAJIMA, Takao<sup>3</sup>

<sup>1</sup> 東京大学大学院理学系研究科地球惑星科学専攻, <sup>2</sup> チェコ共和国地質科学アカデミー, <sup>3</sup> 京都大学大学院理学研究科地質学鉱物学教室

<sup>1</sup>Department of Earth and Planetary Science, Graduate School of Science, The University of Tokyo, <sup>2</sup>Geologicky ustav Akademie, Czech Republic, <sup>3</sup>Department of Geology & Mineralogy, Faculty of Science, Kyoto University

Presence of garnet peridotites in *HP/UHP* metamorphic terranes is taken as evidence for interaction between crust and mantle during orogeny. In order to constrain the timing of interaction, *P-T* paths for both peridotites and crustal rocks have been constrained, which demonstrated that there are significant gaps between peak pressure of peridotites and host continental crust. In this contribution, we will show an evidence for multiple exhumations recorded by clinopyroxene (*Cpx*) megacryst discovered in the garnet peridotite from the Bohemian Massif, and will constrain the timing of crust-mantle interaction in the light of the new data.

The Gföhl Unit experienced the highest metamorphic grade in the Moldanubian zone of the Bohemian Massif, and it mainly consists of quartz-feldspathic garnet-kyanite granulite with peak condition at 2.2-2.3 GPa and 1000 °C (Vrana et al., 2013, *J. Geosci.* **58**, 347-378), although Kotková et al. (2011, *Geology* **39**, 667-670) recently found diamond and coesite from the Saxony-type granulite. Gföhl granulite occurs as tens km-sized isolated blocks and contains garnet peridotites whose peak conditions were estimated as 2-6 GPa and 850-1350 °C (Medaris et al., 2005, *Lithos* **82**, 1-23). We studied *P-T* path for the garnet peridotite at Lom pod Libínem quarry in the Prachatice granulite massif in the south Bohemia. Lom pod Libínem (*LPL*) peridotite generally displays granoblastic texture consisting of mm-sized (0.1-5.0 mm in diameter) garnet, pyroxenes, olivine, and most garnet grains are transformed to kelyphite and are replaced by phlogopite. *LPL* peridotite includes a lot of cm-size *Cpx* megacrysts. Among them, the largest megacryst (3×5 cm) shows a strong chemical zoning consisting of three zones (Fig. 1), namely, core, mantle and rim: (1) the "pale-green core" is poor in Ca-Tschemak (CaTs, ~6 mol.%) and rich in Enstatite (En, ~9 mol.%) (components after Simakov, 2008, *Lithos* **106**, 125-136) and includes phlogopite and orthopyroxene, (2) the "mantle" is lower in CaTs. (4-5 mol.%) and En. (4 mol.%), and includes hornblende, chlorite, apatite, titanite, andradite, olivine, and celsian, and (3) the "rim", rich in CaTs. (7-9 mol.%) and En. (9 mol.%), includes olivine, phlogopite, and hornblende, respectively. The mm-size *Cpx* has identical composition to the megacryst-rim, and (4) the smaller *Cpx* is richer in CaTs. (12 mol.%) and poorer in En. (7 mol.%). Four mineral stages can be identified: **Stage 1** is defined by megacryst-core coexisting with phlogopite, orthopyroxene. Assuming the co-existence with garnet, the equilibrium condition was estimated at ~4 GPa and 1000 °C by use of the single *Cpx* geothermobarometer (Nimis & Taylor, 2000, *Contrib. Mineral. Petrol.* **139**, 541-554), **Stage 2** is defined by the "mantle" coexisting with hornblende, chlorite, orthopyroxene, and andradite, which were equilibrated at ~700 °C, *P*<2.5 GPa. **Stage 3** is defined by the core of mm-size pyroxenes, garnet and olivine that were equilibrated at ~3.0 GPa and 1000 °C in the garnet lherzolite facies. **Stage 4** is defined by the matrix spinel lherzolite assemblage equilibrated at *T*~800 °C at 1-2 GPa.

We envisaged the following juxtaposition mechanism: *LPL* peridotite originally came from the upper mantle (4 GPa) that exhumed to the depth of <2.5 GPa and was partially transformed to chlorite peridotite. Assuming that *LPL* peridotite was entrained in the crust at Stage 2, peridotite and host continental crust could have shared the Variscan *UHP* metamorphism at 3 GPa (Stage 3), followed by final exhumation. Although the peak *P-T* condition of Stage 3 is still higher than that of country granulite (<2.3 GPa), this could reflect different degree of retrogression during the final exhumation. Our study suggests some orogenic peridotites were exotically derived from the *UHP* mantle (>4 GPa), where continental crust have not subducted more than 3 GPa. This will give an another solution to the observed pressure gaps between orogenic peridotites and host continental crust.

キーワード: 造山型橄欖岩, 超高压変成岩, 地殻マントル相互作用, ボヘミア山塊, 多重上昇履歴

Keywords: orogenic peridotite, UHP metamorphism, crust-mantle interaction, Bohemian Massif, multiple exhumations

SCG08-04

会場:311

時間:4月29日 10:00-10:20




Fig. 1. Cpx megacryst (*Al map*) from the Lom pod Libinem peridotite and its compositions along the line.

## Modelling of the Phase Relations in High- and Ultrahigh-pressure Metabasic Rocks Modelling of the Phase Relations in High- and Ultrahigh-pressure Metabasic Rocks

WEI, Chunjing<sup>1\*</sup>  
WEI, Chunjing<sup>1\*</sup>

<sup>1</sup>School of Earth and Space Sciences, Peking University

<sup>1</sup>School of Earth and Space Sciences, Peking University

Pseudosections calculated with THERMOCALC predict that for glaucophane-lawsonite eclogite facies conditions (500-600 °C and 18-28 kbar), MORB compositions in the NCKMnFMASHO system will contain glaucophane, garnet, omphacite, lawsonite, phengite and quartz, with chlorite at lower temperature and talc at higher temperature. In these assemblages, the pyrope contents (X<sub>py</sub>) in garnet is mostly controlled by variations in temperature, grossular content (X<sub>gr</sub>) is strongly controlled by pressure, and the silica content (Si-) in phengite increases linearly with pressure. As the P-T conditions for these given isopleths are only subtly affected by common variations in bulk-rock compositions, the P-T pseudosections potentially present a robust geothermobarometric method for natural glaucophane-bearing eclogites. The maximum X<sub>py</sub> content may define the temperature peak (T<sub>max</sub>) and the minimum X<sub>gr</sub> content constrains the pressure peak (P<sub>max</sub>) conditions. An isothermal decompression of these lawsonite-bearing assemblages would result in epidote-bearing assemblages through dehydration reactions such as lawsonite + omphacite = glaucophane + epidote + H<sub>2</sub>O, releasing a large amount of bound fluid. Thus, most natural HP epidote eclogites may have experienced a metamorphic stage of lawsonite stability.

Under low-T UHP conditions (>28 kb, 550-650 °C), basic rocks are predicted to contain garnet, omphacite, lawsonite, phengite, coesite and talc. In this assemblage, the X<sub>py</sub> contents steadily increase as temperature rises and the Si-in phengite increases linearly with pressure. However, the X<sub>gr</sub> content is very sensitive as pressure changes, showing slowly decrease as pressure rises. The peak P-T conditions for low-T UHP eclogites can be determined using the isopleths of maximum X<sub>py</sub> and Si-in phengite in P-T pseudosections. An isothermal decompression of these low-T UHP eclogites at temperature i.e. 600 °C would result in disappearance of lawsonite and talc in the peak stage, but appearance of glaucophane, epidote and kyanite, forming the mineral assemblages involving garnet + omphacite + glaucophane + epidote ? kyanite + quartz/coesite + phengite commonly observed. Moreover, garnet in the low-T UHP eclogites is characteristic of growth zoning with its rims containing lower X<sub>gr</sub> and higher X<sub>py</sub> contents.

Under Medium-T UHP conditions (>28 kb and >650 °C), basic rocks are predicted commonly to contain garnet + omphacite + lawsonite + phengite + coesite. In this assemblage, the X<sub>py</sub> in garnet mostly depends on bulk compositions, whereas the X<sub>gr</sub> in garnet and the Si-contents in phengite regularly increase, respectively, as temperature and as pressure rise, and thus, can provide robust thermobarometric constraints. Decompression of the eclogites with lawsonite in the peak stage is inferred to be dominated by lawsonite dehydration, resulting in increase in the mode of anhydrous minerals, or further eclogitization, and formation of epidote porphyroblasts and kyanite-bearing quartz veins in eclogite. As lawsonite dehydration can facilitate evolution of assemblages under fluid-present conditions, the UHP eclogites with lawsonite are hard to memorize their real peak P-T conditions.

キーワード: HP-UHP eclogites, pseudosection, geothermobarometer, basic rock

Keywords: HP-UHP eclogites, pseudosection, geothermobarometer, basic rock

## DEHYDRATED FLUID AND SEISMIC DEFORMATION IN DEEP SUBDUCTION ZONE ?constraints from lawsonite eclogite and olivine-opx spinifex DEHYDRATED FLUID AND SEISMIC DEFORMATION IN DEEP SUBDUCTION ZONE ?constraints from lawsonite eclogite and olivine-opx spinifex

岡本 和明<sup>1\*</sup>  
OKAMOTO, Kazuaki<sup>1\*</sup>

<sup>1</sup> 埼玉大学, <sup>2</sup> 東京学芸大学連合大学院  
<sup>1</sup>Saitama University, <sup>2</sup>Joint Graduate school, Tokyo Gakugei University

**Introduction:** It has been considered that there is a correlation between the double seismic zone and metamorphic dehydration reaction in deep slab. The location of the upper limits of the upper seismic plane correspond to metamorphic facies boundary where H<sub>2</sub>O contents change in subducting crust; numerous earthquakes from 60 to 110 km depths in the lawsonite-blueschist facies, many earthquakes in the lower crust of the slab from 110 to 150 km depths in the lawsonite-amphibole eclogite facies and few earthquakes in the lawsonite eclogite facies. It was considered that the dry eclogite is distributed in the area where there is few earthquake [1]. However, ultrahigh pressure experiments and thermodynamic calculation are both demonstrating that the lawsonite eclogite is stable [2] in the area. In order to unravel relation between dehydration and seismic deformation, we have investigated dehydration process of natural metamorphic rocks recording very cold geo-thermal history in the crust and lithosphere in the slab.

**Lawsonite eclogite (Alpine Corsica):** Alpine Corsica is the best representative field area exhibiting lawsonite eclogite and blueschist as a coherent high pressure, low temperature metamorphic belt. Ophiolite sequence (oceanic plate structure) is also well-preserved, and the pillow structure is clearly recognized in the lawsonite eclogite. Recent petrological researches have revealed that both blueschist (rims of the pillow) and lawsonite eclogite (core of the pillow) are stable in the same pressure and temperature condition [3] because chemical variation including water content creates both lawsonite-amphibole eclogite and lawsonite eclogite in different portion of subducted crust. We carefully observed microtexture of the lawsonite eclogite and blueschist and have found that omphacite vein and lawsonite vein in mylonitized blueschist matrix.

**Olivine-opx spinifex in serpentinite (Cerro del Almirez, Nevado complex):** Metamorphic olivine after antigorite has been described in Italian Alps and also from the Mt. Shiraga, Japan. However, the olivine was formed with talc and fluid by antigorite breakdown reaction in pressures lower than 1.5 GPa. Spinifex olivine with opx in the Cerro del Almirez, is the product at pressures ( $P > 1.5$  GPa) relevant to the lower seismic plane beneath Northeast Japan. In Cerro del Almirez, olivine-opx rocks underlie antigorite schist by a contact [4]. In the olivine-opx rocks, the blade-like, elongated olivine and opx were grown, representing spinifex texture. It clearly indicates the presence of large amount of water facilitate crystallization of elongated olivine with opx.

**Discussion and conclusion:** In the lawsonite eclogite in the Alpine Corsica, blueschist and lawsonite coexist together reflecting chemical difference in pillowed structure or lithology. Omphacite and lawsonite veins are observed along the shear band in mylonitized blueschist. It suggests that decomposition of glaucophane caused hydro-fracturing and precipitate omphacite and lawsonite vein. Garnet was grown statically close to the vein.

Olivine-opx spinifex in the serpentinite, Cerro del Almirez, were probably recrystallized in the presence of large amount of water. The estimated dehydration reaction has a negative P-T slope at pressures higher than 1.5 GPa. The reaction is volume reducing reaction and the olivine-opx spinifex texture was formed under volume reducing reaction.

**References:** [1] S. Kita et al., *Geophys. Res. Lett.*, 33, doi:10.1029/2006GL028239. (2006). [2] K. Okamoto et al., *Geochemistry*, 46 205-215 (2013). [3] A. Vitale Brovarone et al., *J. Met. Geol.*, 29, 583-600, (2011). [4] V. Trommsdorff et al., *CMP*, 132, 139-148 (1998).

**キーワード:** 二重深発地震面, 上面地震帯, ローソン石エクロジャイト, 下面地震帯, 変成かんらん石、輝石スピニフェック  
す, 脱水脈

**Keywords:** double-seismic zone, upper seismic plane, Lawsonite eclogite, lower seismic plane, olivine-opx spinifex, dehydrated vein

## Ultrahigh-pressure eclogites: paleo-environment indicators Ultrahigh-pressure eclogites: paleo-environment indicators

UR REHMAN, Hafiz<sup>1\*</sup>

UR REHMAN, Hafiz<sup>1\*</sup>

<sup>1</sup>Department of Earth and Environmental Sciences, Graduate School of Science and Engineering, Kagoshim

<sup>1</sup>Department of Earth and Environmental Sciences, Graduate School of Science and Engineering, Kagoshim

Ultrahigh-pressure (UHP) eclogites generally form by the metamorphism of mafic lithologies (gabbros or basalts) at depths greater than 90 km (minimum stability field of coesite) indicating P-T conditions of >2.7 GPa and 600-800 °C. At such conditions most minerals reequilibrate their chemical elements (e.g. major and trace and even isotopes) and new minerals crystallize or grow at the expense of other minerals formed during the magmatic crystallization. Some chemical elements, considered as relatively less mobile or immobile (e.g. Sm, Nd, Lu, Hf), are widely used for the extraction of past records the rocks have evolved through. Besides those elements, oxygen, the major component of silicates and oxides, impart important information related to protolith formation of rocks and their metamorphism. In general, most basaltic rocks show a narrow range of  $\delta^{18}\text{O}$  (ca.  $+5.7 \pm 0.3$  ‰ relative VSMOW) and values for altered basalts vary from 0 to +12 ‰.

In this paper, I discuss the origin of low or negative  $\delta^{18}\text{O}$  values; recently we found in the Himalayan UHP eclogites of Kaghan Valley and explain the mechanism how these low values were acquired. These eclogites are formed during Eocene by the India-Asia collision and their protoliths were the Panjal Trap basalts which were emplaced in Permian when Indian Plate was part of Gondwana. The  $\delta^{18}\text{O}$  values are as low as  $-2.25$  ‰ in the fresh parts of eclogites and increase towards more positive in the retrogressed or amphibolitized parts. The unusually low  $\delta^{18}\text{O}$  values in eclogites are interpreted to have resulted from the hydrothermal alteration of the protoliths by meteoric water interaction when Greater India was still at southern high latitudes (>65 degrees S) during the Permian indicating glacial paleo-environment. These low  $\delta^{18}\text{O}$  values were frozen-in in the protolith rocks and did not change during subduction-related UHP metamorphism. However retrogressive process, due to infiltration of  $^{18}\text{O}$ -rich fluids during exhumation, shifted these values towards more positive range.

キーワード: Ultrahigh-pressure eclogites, Himalaya, Oxygen isotope, Paleo-environment

Keywords: Ultrahigh-pressure eclogites, Himalaya, Oxygen isotope, Paleo-environment

## 第2大陸の役割 Role of the second continent

河合 研志<sup>1\*</sup>; 市川 浩樹<sup>2</sup>; 山本 伸次<sup>4</sup>; 土屋 卓久<sup>2</sup>; 丸山 茂徳<sup>3</sup>  
KAWAI, Kenji<sup>1\*</sup>; ICHIKAWA, Hiroki<sup>2</sup>; YAMAMOTO, Shinji<sup>4</sup>; TSUCHIYA, Taku<sup>2</sup>; MARUYAMA, Shigenori<sup>3</sup>

<sup>1</sup> 東京工業大学大学院理工学研究科地球惑星科学専攻, <sup>2</sup> 愛媛大・GRC, <sup>3</sup> 東京工業大学地球生命研究所, <sup>4</sup> 東京大学大学院総合文化研究科広域科学専攻

<sup>1</sup>Department of Earth and Planetary Sciences, Tokyo Institute of Technology, <sup>2</sup>Geodynamics Research Center, Ehime University, <sup>3</sup>Earth-Life Science Institute, Tokyo Institute of Technology, <sup>4</sup>Department of Earth and Astronomy, Graduate School of Arts and Sciences, The University of Tokyo

It has been thought that granitic crust, having been formed on the surface, must have survived through the Earth's evolution because of its buoyancy. Recent geological studies have suggested that a significant amount of crustal material has been lost from the surface due to delamination, continental collision, and subduction at oceanic?continental convergent margins (von Huene and Scholl 1991; Yamamoto et al. 2009; Ichikawa et al. 2013a). If so, then the subducted crustal materials are expected to be trapped in the mid-mantle due to the density difference from peridotitic materials induced by the phase transition from coesite to stishovite (Kawai et al. 2013). In order to study the effect of the subducted granitic materials floating around the mantle transition zone, we conducted two-dimensional numerical experiments of mantle convection incorporating a continental drift with a heat source placed around the bottom of the mantle transition zone. We found that the addition of heat source in the mantle transition zone considerably enhances the onset of upwelling plumes in the upper mantle, which further reduces the time scale of continental drift. The heat source also causes massive mechanical mixing, especially in the upper mantle. The results suggest that the heat source floating around the mantle transition zone can be a possible candidate for inducing the supercontinent cycle (Ichikawa et al. 2013b).

キーワード: 第2大陸, ウィルソンサイクル, 花崗岩, 構造浸食  
Keywords: Second continent, Wilson's cycle, Granite, Tectonic erosion

## Where had the primordial continent gone? Where had the primordial continent gone?

丸山 茂徳<sup>1\*</sup>; 河合 研志<sup>2</sup>; 土屋 卓久<sup>3</sup>  
MARUYAMA, Shigenori<sup>1\*</sup>; KAWAI, Kenji<sup>2</sup>; TSUCHIYA, Taku<sup>3</sup>

<sup>1</sup> 東京工業大学地球生命研究所, <sup>2</sup> 東京工業大学, <sup>3</sup> 愛媛大学  
<sup>1</sup>ELSI Tokyo Institute of Technology, <sup>2</sup>Tokyo Institute of Technology, <sup>3</sup>Ehime University

There are no Hadean rocks on the Earth's surface. This indicates (1) there was no continents on the primordial Earth, or (2) continents were present in the Hadean but lost afterwards. It is well-known that Moon surface was covered by anorthositic continental crust with KREEP basalts with ca. 50-60km thickness. Those rocks are the fractionated final residues of magma ocean after the giant impact at 4.56Ga.

The Earth must have been completely melted if giant impact was correct, but due to 6 times larger gravity, the thickness of anorthositic continents must be 21 km. Moreover, during the consolidation of magma ocean, bulk of anorthositic blobs could be transformed into (1) zoisite+kyanite+quartz, (2) grossular+kyanite+quartz, (3) grossular +kyanite+coesite, (4) grossular +kyanite+stishovite, in 300km depth.

Absence of TTG rocks in the Hadean and nearly absent in early Archean less than 5% among the surface of present continents must be caused by extensive tectonic erosion by subducting slabs which are present even today at trench. Therefore, it is wrong to believe that low-density granite must have accumulated on the surface once formed on the surface of the Earth.

Density calculation of anorthosite, MORB, harzburgite, and pyrolite using first principles calculation showed that anorthosite was heavier if it convected at depth in mantle transition zone. If it turns into lower mantle, it becomes to be heaviest among those as mentioned above, suggesting the stability field at D" layer on the bottom of mantle which is right above CMB.

However, it depends on the geothermal gradient on which phase change of Al<sub>2</sub>O<sub>3</sub> occurs to determine whether or not the heaviest among all rocks at CMB. Depending on cooling the CMB, meta-anorthosite could have been buoyant particularly after the mantle overturn at 2.7-2.6Ga.

Before the overturn, if the basal magma ocean was present in the Archean, the lost primordial continents must have been melted into basal magma ocean where anorthosite-KREEP basalts mixed with FeO-enriched primordial basal magma ocean. The possible bulk chemistry can be estimated, though qualitatively; it must be super-enriched in anorthositic component. It can be expressed in Ca-Pv, Mg-Pv, Ferro-periclase as major components.

When, mantle overturn occurred at 2.7-2.6Ga, low-temperature materials of upper mantle must have cooled down the basal magma ocean to consolidate to crystallize Ca-Pv rock, bi-mineral Ca-Pv + Mg-Pv rock, and the tri-mineral Ca-Pv + Mg-Pv + FM periclase rock. According to the formation of these rocks, those three rocks would have been behaved differently, although depends on not only density but also mass.

Assuming the enough volume of Ca-Pv rocks, those rocks may be floating in mid-depth of lower mantle.

## 沈み込み起源の超高压クロミタイト：西彼杵変成岩の例 Subduction origin for UHP chromitite from the Nishisonogi metamorphic rocks, Western Kyushu, Japan

西山 忠男<sup>1\*</sup>; 森部 陽介<sup>1</sup>; 石丸 聡子<sup>1</sup>; 荒井 章司<sup>2</sup>; 森 康<sup>3</sup>; 重野 未来<sup>3</sup>

NISHIYAMA, Tadao<sup>1\*</sup>; MORIBE, Yousuke<sup>1</sup>; ISHIMARU, Satoko<sup>1</sup>; ARAI, Shoji<sup>2</sup>; MORI, Yasushi<sup>3</sup>; SHIGENO, Miki<sup>3</sup>

<sup>1</sup> 熊本大学自然科学研究科, <sup>2</sup> 金沢大学地球科学教室, <sup>3</sup> 北九州市立いのちのたび博物館

<sup>1</sup>Graduate School of Science and Technology, Kumamoto University, <sup>2</sup>Department of Earth Sciences, Kanazawa University,

<sup>3</sup>Kitakyushu Museum of Natural History and Human History

Ultrahigh-pressure (UHP) chromitite from the Luobsa Ophiolite in non UHP terrane has been an enigma because of its peculiar occurrence. We newly found a UHP chromitite from serpentinite in the Nishisonogi metamorphic rocks (NMR), a member of the Nagasaki Metamorphic Rocks, in Western Kyushu, following our finding<sup>1</sup> of it from the Higo Metamorphic Rocks (HMR), Central Kyushu. The UHP chromitite from NMR documents well a fluid-chromite interaction, showing partial graphitization of microdiamond. Such a fluid-chromite interaction is not observed in HMR chromitite. The NMR are high P/T (epidote-glaucophane schist subfacies) metamorphic rocks of Cretaceous in age, mainly consisting of pelitic and psammitic schists intercalating with minor basic schists<sup>2</sup>. The peak metamorphic condition of the crystalline schists is estimated as 1.4 GPa and 520 °C by an assemblage of garnet with inclusions of chloritoid and omphacite, glaucophane, paragonite, and phengite in a garnet galucophanite<sup>3</sup>. Serpentinite and serpentinite melanges occur as elongated bodies or lenses concordant with schistosity trending N-S of the country schists<sup>1</sup>. Jadeitite and omphacitite occur as tectonic blocks in the serpentinite melange, showing the peak condition of 1.5 GPa and 500 °C by coexistence of jadeite and quartz<sup>4</sup>. Microdiamond-bearing chromitite was found from serpentinite in a melange at Ooseto Town, Saikai City. Chromitite occurs as a thin layer several cm thick and meter-size long in a serpentinite with numerous magnesite (or ankerite) veins. The layer is strongly deformed to show a schlieren-like structure. The serpentinite consists of fine-grained antigorite with no relics of olivine and pyroxenes. The chromitite consists of an aggregate of rounded and fractured chromite crystals with small amounts of talc and magnesite as a matrix and veins. Microdiamond occurs as aligned crystals in narrow zones ranging from a few μm to several tens of μm in chromite. Chromite is zoned, consisting of Mg-rich core ( $\text{Mg}_{0.33}\text{Fe}^{2+}_{0.65}\text{Mn}_{0.03}$ )( $\text{Cr}_{0.84}\text{Al}_{0.12}\text{Fe}^{3+}_{0.04}$ )<sub>2</sub>O<sub>4</sub> and Fe-rich rim ( $\text{Mg}_{0.06}\text{Fe}^{2+}_{0.89}\text{Zn}_{0.02}\text{Mn}_{0.03}$ )( $\text{Cr}_{0.85}\text{Al}_{0.12}\text{Fe}^{3+}_{0.04}$ )<sub>2</sub>O<sub>4</sub>. The microdiamond-bearing zones are conspicuously richer in Fe<sub>2</sub>O<sub>3</sub> [ $\text{Mg}_{0.03}\text{Fe}^{2+}_{0.94}\text{Mn}_{0.04}\text{Zn}_{0.01}$ ]( $\text{Cr}_{0.67}\text{Ti}_{0.01}\text{Fe}^{3+}_{0.31}$ )<sub>2</sub>O<sub>4</sub>], observed as a brighter zone in a BSE image, than other part of chromite in the same grain. The network-like distribution of the zones clearly indicates fluid infiltration associated with the following exchange reaction of trivalent cations between chromite and the fluid:  $\text{Cr}^{3+} + \text{Al}^{3+}(\text{in chromite}) = \text{Fe}^{3+}(\text{in fluid})$

Microdiamond occurs either as polyhedral or as platy crystals, 1 to several μm across. Identification of diamond was carried out with an energy dispersive X-ray spectroscopy (EDS) analysis (carbon peak) and Raman spectroscopy with a He-Ne laser. We observed a broad Raman peak at 1331 cm<sup>-1</sup>, which is comparable to the peak (1332 cm<sup>-1</sup>) characteristic of diamond. Graphite peak at about 1600 cm<sup>-1</sup> is also observed, showing partial graphitization of microdiamond. These lines of evidence shows that the fluid infiltration may have occurred after inclusion of microdiamond.

It is quite astonishing that microdiamond is preserved in such a completely serpentinitized ultramafic rock. Chromite can be a good container of microdiamond to prevent graphitization during geologically long duration of exhumation and serpentinitization. Our finding suggests the subduction origin of UHP chromitite from NMR rather than mantle migration origin<sup>5</sup> in the case of the Luobusa Ophiolite.

### References

1. Nishiyama, T., Shiosaki, D., Eguchi, H., and Yoshiasa, A. JpGU Meeting, S-MP46 (2014); 2. Nishiyama, T. Mem. Geol. Soc. Japan 33, 237-257 (1989); 3. Moribe, Y. unpublished Mc thesis, Kumamoto University (2014); 4. Shigeno, M., Mori, Y., Shimada, K., and Nishiyama, T. Eur. J. Mineral., 24, 289-311 (2012); 5. Arai, S. J. Mineral. Petrol. Sci., 105, 280-285 (2010)

キーワード: マイクロダイヤモンド, 超高压クロミタイト, 沈み込み帯, 西彼杵変成岩, 超高压変成岩

Keywords: microdiamond, UHP chromitite, subduction zone, Nishisonogi metamorphic rocks, ultrahigh-pressure metamorphic rocks