

Vp/Vs ratio in the southernmost Japan Basin and its transition area, Japan Sea deduced from the seismic survey

SATO, Takeshi^{1*} ; NO, Tetsuo¹ ; KODAIRA, Shuichi¹ ; TAKAHASHI, Narumi¹ ; KANEDA, Yoshiyuki¹

¹JAMSTEC

The Japan Sea is one of very well studied back-arc basins in the northwestern Pacific. To clarify the formation process of the back-arc basin in the Japan Sea, many seismic surveys using ocean bottom seismographs (OBSs) and control sources have been conducted in the sea. The Japan Basin, which located in the northern to eastern Japan Sea, has an oceanic crust formed by seafloor spreading (Hirata et al., 1992; No et al., submitted). On the other hand, the ocean-continent transition area between the Japan Basin and the continental shelf in the eastern margin of this sea may have a thick oceanic crust (No et al., submitted). However, it is unknown the origin and the nature of this thick oceanic crust, due to the lack of the information about lithology in the transition area. To understand the origin and the nature of this thick oceanic crust, it is necessary to obtain the information lithology in the crust of the transition and the basin areas. For this study, we will present the Vp/Vs ratio of the crust from the southernmost Japan Basin to its transition area.

From the southernmost Japan Basin to the continental shelf off the west of Aomori and the northern Oga Peninsula, seismic surveys using OBSs and an air-gun array were undertaken. In vertical record sections of several OBSs, not only the first arrived phases but also later phases reflected from interfaces in the crust and uppermost mantle are visible. Moreover, in horizontal record sections of several OBSs, converted phases from P- to S-waves are apparent. In this study, we have obtained the S-wave velocity structure using travel times of these converted phases. Then, we have obtained the Vp/Vs ratio in the crust from the southernmost Japan Basin to its transition area using the obtained P- and S-wave velocity structures.

In the southernmost Japan Basin off west of Aomori, the Vp/Vs ratio in the sedimentary layer of shows 4 to 8 and has a lateral variation. The Vp/Vs ratios in the crustal upper and lower parts show around 1.85 and 1.8, respectively. On the other hand, in the transition area, the Vp/Vs ratio in the crustal upper part is similar to that in the southernmost Japan basin. This Vp/Vs ratio may show that the nature of the whole crust in the basin area and of the crustal upper part in the marginal area has an oceanic origin. Therefore, the crusts in southernmost Japan Basin and in its transition area are suggested as an oceanic crust and a thick oceanic crust, respectively. The oceanic crust formed by the opening of the Japan Sea may extend to the transition area of the southernmost Japan Basin.

Crustal structure study of the Sea of Japan: Recent results and future perspectives

NO, Tetsuo^{1*} ; SATO, Takeshi¹ ; KODAIRA, Shuichi¹ ; ISHIYAMA, Tatsuya² ; SATO, Hiroshi² ; TAKAHASHI, Narumi¹ ; OIKAWA, Nobutaka¹ ; KANEDA, Yoshiyuki¹

¹JAMSTEC, ²ERI, Univ. of Tokyo

In recent years, crustal structure study of the Sea of Japan has advanced. Various new seismic data have been obtained in the Sea of Japan; for example, the two ship seismic surveys study (Sato et al. 2007) and onshore-offshore seismic survey (Earthquake Research Institute, 2013) conducted by the University of Tokyo, and 2D/3D seismic reflection survey conducted by R/V SHIGEN (JOGMEC, 2013).

From 2007 to 2012, we conducted marine seismic surveys using the multichannel seismic reflection system and ocean bottom seismometers; the surveys covered the area between the Japanese coast of the Sea of Japan and the Yamato and Japan Basins. Based on the results, the crustal structure of the eastern margin of the Sea of Japan was classified into three types: island arc crust, thick oceanic crust, and oceanic crust (Sato et al. 2014; No et al. submitted). In addition, our studies found that the contractive deformation zones of the eastern margin of the Sea of Japan are associated with the crustal structure distribution. Further, seismic data suggests that the crustal structure in the south (off Yamagata to Niigata) differs from that of the north (off Akita to Nishi-tsugaru). These differences are critical in understanding the relation between the spatial distributions of the seismogenic and contractive deformation zones (JAMSTEC, 2013). These results can contribute to the review of long-term evaluations of earthquake occurrence potentials and the discussion of the seismogenic study in the eastern margin of the Sea of Japan.

In 2013, new projects have been observing and studying earthquakes and tsunamis in the Sea of Japan. *Integrated Research Project on Seis* scheduled in 2014 in order to conduct a seismic survey in the blind areas of the existing observations, which are in the southwest region of the Sea of Japan and off western Hokkaido. The addition of new observation data will advance the study of crustal structure in the Sea of Japan. In this study, we aim to improve the accuracy of the position and size of the source faults in the Sea of Japan. In addition, we investigate the relation among tectonic history, crustal structure, and factors that form source faults in the Sea of Japan. In the future, we aim to clarify the seismogenic zone of the Sea of Japan using new seismic data.

Keywords: the Sea of Japan, crustal structure, MCS, OBS

Results of 2013 Off-Joetsu and Hokuriku survey for the integrated research project on seismic and tsunami hazards around

SATO, Hiroshi^{1*} ; ISHIYAMA, Tatsuya¹ ; SHIRAIISHI, Kazuya² ; ABE, Susumu² ; KATO, Naoko¹ ; KURASHIMO, Eiji¹ ; TAKEDA, Tetsuya³

¹Earthquake Research Institute, Univ. Tokyo, ²JGI. Inc., ³NIED

An obvious convergent plate boundary cannot be recognized in the Sea of Japan, and convergence accommodates in defused wide area of the back arc. To estimate Tsunami and seismic hazards along the coastal area of Sea of Japan, more detailed survey to identify source faults are needed. A new research project funded by MEXT named "the integrated research project on seismic and tsunami hazards around the Sea of Japan" began in FY 2013. To obtain the information of source faults, we performed deep seismic reflection profiling off-Joetsu and Hokuriku area in the central part of Honshu, Japan. We used two vessels; a gun-ship with 3020 cu. inch air-gun and a cable-ship with a 2-km-long, streamer cable with 156 channels and 480 cu. inch air-gun. Common-mid point reflection data were acquired along 9 seismic lines with total 715 km in length. The seismic profiles portray the structure of failed rift basins, such as Toyama trough and Sado strait, bounded by rift axis reverse faults with rift axis vergence, which represents reactivation of boundary faults between mafic intrusion and pre-rift basement. Noto Peninsula is marked by syn-rift normal faults and their reactivation by shortening deformation. The back arc side of the SW-Japan arc experienced NS trending shortening deformation in the latest Miocene. From the Noto peninsula, undeformed Pliocene sediments covers folded Miocene. Some normal faults reactivated as active strike-slip and reverse faults. The survey results contributed to construct source faults models of Tsunami and seismic hazards estimation.

Keywords: Sea of Japan, source fault, crustal structure, seismic reflection profiling, Off-Joetsu, off-Hokuriku

Lithospheric Structure of the Hidaka Collision Zone, Hokkaido, from Reanalysis of 1998-2000 Hokkaido Transect Data IV

IWASAKI, Takaya^{1*}; TSUMURA, Noriko²; ITO, Taniao³; SATO, Hiroshi¹; KURASHIMO, Eiji¹; HIRATA, Naoshi¹; ARITA, Kazunori⁴; NODA, Katsuya⁵; FUJIWARA, Akira⁵; ABE, Susumu⁵; KIKUCHI, Shinsuke⁶; SUZUKI, Kazuko⁷

¹Earthquake Research Institute, the University of Tokyo, ²Chiba University, ³Teikyo Heisei University, ⁴Hokkaido University, ⁵JGI. Inc., ⁶JAPEX, ⁷Schlumberger Ltd.

The Hidaka region in the central part of Hokkaido Island, Japan, is known as an arc-arc collision zone ongoing from the middle Miocene. In 2012, we started reinterpretation for a series of seismic reflection/refraction surveys from 1994 to 2000 in this collision zone. In this analysis, we used integrated and sophisticated processing and analysis techniques, including CRS/MDRS method for seismic reflection data and refraction tomography both very dense arrival time data from both the reflection and refraction/wide-angle reflection data. The most important finding so far obtained is a clear image of the NE Japan Arc subducting eastward under the northern part of the collision zone. However, the following problems are remained unsolved.

(1) Shallow structure beneath the Hidaka Collision zone is still unsolved. Particularly, the structure just east of the Hidaka Main Thrust is not sufficiently evaluated from our seismic data.

(2) Delamination of the Hidaka crust as in the southern part of this collision zone is not unclear. Our CRS/MDRS processing for the reflection data provided no positive evidence for the delamination.

(3) Deeper collision structure of the NE Japan Arc and the Kuril Arc is still not constrained. It is necessary to elucidate the subducting structure of the NE Japan Arc from amplitude data as well as travel time data.

In this paper, we focus the items (1) and (3) from seismic refraction/wide-angle reflection approach. Previous refraction tomography elucidated a thick (4-5 km) undulated sediments in the hinterland, the outcrop of crystalline crust beneath the Hidaka Metamorphic Belt with higher V_p and V_p/V_s and an enormously thick (>8-10 km) sedimentary package beneath the foreland. In order to obtain the more reliable structure model, we intensively revised the travel time data obtained both from seismic reflection/wide-angle reflection line and reflection lines. The seismic tomography using these revised data sets indicate a clearer high velocity (>6.1 km/s) anomaly just east of the HMT. We also recognized some wide-angle reflections around 5-10 km depth beneath the HMT, from which we expect to determine the finer structure at the collision front. Our present analysis indicates the wide-angle reflection data sample a part of the lower crust of the subducting NE Japan Arc beneath the fold-and-thrust belt. According to the preliminary result, its velocity is ranging from 6.5-7.0 km/s. By combining the amplitude analysis, we expect to estimate the more reliable Moho depth of the NE Japan Arc than in the previous analyses.

Keywords: active source seismic experiment, collision, arc, crustal structure, lithosphere

Seismic crustal structure beneath the southeastern part of northeast Japan by dense seismic array observation

KURASHIMO, Eiji^{1*} ; SATO, Hiroshi¹ ; ISHIYAMA, Tatsuya¹ ; HIGASHINAKA, Motonori² ; ABE, Susumu² ; IWASAKI, Takaya¹

¹ERI, Univ. Tokyo, ²JGI, Inc.

The 2011 Tohoku-Oki Earthquake (Mw9.0), that occurred on the Japan Trench off the eastern shore of northern Honshu, Japan, generated enormous crustal deformations. Seismic activity in northeastern Japan increased significantly after the 2011 Tohoku-Oki Earthquake. Detailed crustal structure and deep geometry of the active fault is important to constrain the process of earthquake occurrence. Active and passive seismic experiments were conducted to obtain a structural image beneath the southeastern part of NE Japan (Sato et al., 2013). The geometry of the active faults have been revealed by seismic reflection profiling (Sato et al., 2013). Natural earthquake data set is useful to obtain a deep structural image. Forty portable seismographs were deployed along a 70-km-long line between Souma and Takahata during the period from August 16, 2012 to December 24, 2012. Each seismograph consisted of a 1-Hz 3-component seismometer and off-line data recorder (Shinohara et al., 1997). Waveforms were continuously recorded at a sampling rate of 200 Hz. In the area of the present study, deep seismic reflection profiling was conducted using vibrators (Sato et al., 2013). The off-line recorders observed the controlled seismic signals as well as natural earthquakes. During the seismic array observation, the JMA located 2956 earthquakes in a latitude range of 37.2-38.5 N and a longitude range of 139.6-141.3 E. We selected 200 earthquakes, all of which occurred near the survey line. In order to obtain a high-resolution velocity model, a well-controlled hypocenter is essential. Due to this, we combined the seismic array data with permanent seismic station data. The arrival times for the first P- and S waves obtained from local earthquakes and Vibrator shots were used in a joint inversion for earthquake locations and three-dimensional Vp and Vp/Vs structures, using the iterative damped least-squares algorithm, simul2000 (Thurber and Eberhart-Phillips, 1999). Permanent seismic stations observed the controlled seismic signals as well as natural earthquakes. We added the arrival time data of these controlled sources into the dataset to improve the shallow velocity structure. The depth section of Vp structure along the survey line shows that lateral variation of the Vp value at a shallow depth. This lateral variation correlates with the surface geology along the profile.

Keywords: dense seismic array observation, seismic tomography, the 2011 Tohoku-Oki Earthquake

Shallow geologic structure around the northern part of the Futaba Fault, northeast Japan, based on gravity survey

KOSHIYA, Shin^{1*} ; TERUI, Kyoko² ; YONEZAWA, Kenta¹ ; SATO, Hiroshi³ ; KATO, Naoko³ ; ABE, Susumu⁴ ; HIGASHINAKA, Motonori⁴

¹Faculty of Engineering, Iwate Univ., ²Graduate School of Engineering, Iwate Univ., ³ERI, Univ. of Tokyo, ⁴JGI, Inc.

The Futaba Fault, bounding the eastern margin of the Abukuma Mountains, is known as a left lateral fault in the Cretaceous and Paleogene period with a remarkable fracture zone of a few hundreds meter width. It trends NNW-SSE and divides into two branches between which the Wareyama horst develops. During early to middle Miocene, E-W extensional stress field caused large normal displacement along the western fault to form a half graben filled with sediments including breccia. In late Miocene, it had been a right lateral fault. In present, the eastern fault is active, along which left lateral offsets with western upheaval ingredient are geo-morphologically observed. Thus the Futaba Fault has experienced the complicated history of development. In this study, we modeled two dimensional shallow geological structure across the faults mainly based on gravity survey. The gravity survey was conducted across the faults with a G-type gravity meter (G827; LaCoste and Romberg Inc.) along two E-W survey lines, one of which is ca. 12 km long, (line 1), and the other of which is ca. 13km long (line 2). Each interval of observation sites is about 200 m. The elevation of observation sites was surveyed with a electric level and a RTK-GPS. Acquired gravity data was processed to obtain Bouguer anomaly mostly according to the methodology of Geological Survey of Japan, AIST (2004). We assumed that the density for Bouguer and terrain corrections were 2.2 g/cm³. In each survey line, Bouguer anomalies after trend correction show the highest value around the Wareyama horst consisting of pre-Paleogene basement rocks and a few maxima in the western side of the horst. We assume four layers in our model, which have densities of 2.00 g/cm³ (layer 1), 2.2 g/cm³ (layer 2), 2.55 g/cm³ (layer 3), and 2.67 g/cm³ (layer 4), respectively. The interpretation of the model is as follows. Layer 1 is correlated to the surface covers and Pliocene sedimentary rocks, layer 2 lower to middle Miocene sedimentary rocks, layer 3 Miocene breccia and layer 4 basement rocks. We will discuss the shallow structure across the faults in detail.

Keywords: Futaba fault, gravity anomaly, active fault

Structural characters of active faults, crustal architecture, and permanent deformation of the Hokuriku region

ISHIYAMA, Tatsuya^{1*} ; SATO, Hiroshi¹ ; KATO, Naoko¹ ; TAKEDA, Tetsuya² ; KURASHIMO, Eiji¹

¹Eathquake Research Institute, University of Tokyo, ²NIED

We discuss in this study about characters of crustal architectures around the Toyama trough revealed by new seismic reflection and refraction profiles and seismic tomography, and active structures based on Neogene geology and tectonic geomorphology. As revealed by onshore offshore deep seismic reflection profiling across the Toyama trough funded by MEXT named as The Integrated Research Project on Seismic and Tsunami Hazards around the Sea of Japan since 2013, crustal architectures across the Toyama trough is characterized by three domains: (1) crustal thrust wedge comprising the northwestern flanks of the Hida Mountains, (2) Neogene sedimentary basin near the axis of the Toyama trough, and (3) reactivated normal faults as thrust (or obliquely slipping) faults beneath the Noto peninsula, comprising structural higher domain west of the Toyama trough. These structural patterns and permanent, late Quaternary crustal deformation recorded by tectonic geomorphology are quite similar to adjacent Neogene sedimentary basins in the backarc failed rifts in the Sea of Japan, including northern Fossa Magna, Niigata, and Akita.

Phase changes and temperature of the subducted crust of Philippine Sea slab beneath Kanto, Japan

ISHIKAWA, Masahiro^{1*} ; NAKAGAWA, Shigeki² ; SAKAI, Shin'ichi² ; HIRATA, Naoshi² ; SATO, Hiroshi² ; KASAHARA, Keiji³

¹Graduate School of Environment Information Sciences, Yokohama National University, ²Earthquake Research Institute, The University of Tokyo, ³Association for the Development of Earthquake Prediction

The Philippine Sea plate subducts beneath the Greater Tokyo Area of Japan. Devastating M8-class earthquakes occurred on the upper surface of the Philippine Sea plate, examples of which are the Genroku earthquake of 1703 (magnitude M8.0) and the Kanto earthquake of 1923 (M7.9). A M7 or greater (M7) earthquake in this region may occur either on the upper surface or intra slab of Philippine Sea plate. To evaluate seismic hazard in the Greater Tokyo Area of Japan we need to clarify the lithological properties of Philippine Sea slab. This study presents an interpretation of the crustal and mantle structure of the Philippine Sea slab beneath Kanto based on recent MeSO-net seismic tomography data. The seismic tomography reveals that P wave velocity of the subducted crust of the Philippine Sea slab increases stepwise at 30 km and 40 km depths beneath the Kanto area. The cause of these two stepwise increases in P wave velocity of subducted crust is expected to correspond to metamorphic phase changes. Mineralogical assemblages of forearc basalt composition of the Izu arc was calculated by Theriak-Domino software, and the phase diagram shows that phase changes to garnet amphibolite and eclogite can account for these two stepwise increase in P wave velocity of the subducted crust of the Philippine Sea slab.

Keywords: slab, phase change, slab temperature, Kanto, Philippine Sea Plate, crust

Geologic structure in and around the Beppu Bay estimated by gravity analysis

YAMAKITA, Satoshi^{1*} ; KUDO, Takeshi² ; ITO, Tanio³

¹University of Miyazaki, ²Chubu University, ³Teikyo-Heisei University

Seismic profiling in the Beppu Bay and the Bungo Strait performed by Kyoto University and JGI inc. from 1988 to 1990 (Yusa et al., 1992) raised new progresses of studies on the geologic structure of the Median Tectonic Line (MTL) in Kyushu and the development of the Beppu Bay Sedimentary Basin (BSB) accompanying it (Yamakita et al., 1995; Ito et al., 1996). The P-wave velocity assumed by Yusa et al. (1992) for the rocks in Ryoke Belt, however, was too low for granitic and metamorphic rocks constituting this Belt, and it is likely that the dip of the MTL beneath it was underestimated. Besides, the structure of the basin in the innermost part of the Beppu Bay has remained uncertain because this part is located in the terminal part of the seismic lines. On the other hand, there are plenty of gravity data in and around the Beppu Bay (Yusa et al., 1992; GSJ, 2000; Gravity Research Group in Southwest Japan, 2001). It can be expected that these gravity data clarify the structure of BSB, combined with correct seismic profiles. Fortunately, a profile reprocessed with re-estimated P-wave velocity along the Bungo straight (J-line) was presented last year (Abe et al., 2013). Using this reprocessed profile and gravity data, we tried to determine the subsurface structure along the G-line of Yusa et al. (1992), trending N70E, 35 km long, from on-land area, across the Asamigawa fault (AF) and the Beppu Bay Central Fault (BCF), to the mouth of the Beppu Bay (Fig. A), occupied by Sanbagawa metamorphic rocks (Sm, $\rho=3.0\text{g/cm}^3$), Ryoke granitic and metamorphic rocks (Rk, $\rho=2.8\text{g/cm}^3$), lower (Bl, $\rho=2.6\text{g/cm}^3$) and upper (Bu, $\rho=2.4\text{g/cm}^3$) sediments of BSB. Assuming only the depth and form of the MTL estimated from the reprocessed profile of J-line and geologic constraints, and the position of the AF on the surface, we determine other subsurface structures to fit with the gravity data through trial and error (Fig. B). In this profile, the upper surface of the Ryoke basement almost coincides with that in the Yusa et al. (1992)'s profile. This fact suggests high reliability of this profile. We concluded the structure of BSB from this and Yusa et al.(1992)'s profiles as follows.

1. The innermost part of the BSB was formed by two listric normal fault systems, NE-dipping Asamigawa Fault System (AFS) and SW-dipping Beppu Bay Central Fault System (BCFS). Both systems formed roll-over structure in the sediments of BSB.
2. The AFS consists of three faults (I, II, III), which converge to the MTL. It is uncertain whether the fault AFS-II reaches the uppermost part of the sediments. The total amount of the vertical displacement of ASF may be up to 3000m, although it depends on the thickness of sediment in the SW-side on AF and the amount of erosion of Ryoke basement
3. Two faults BCFS-I and BCFS-II vertically displaced the bottom of the Bu in 250m and 150m respectively and reach the uppermost part of it, although it is difficult to recognize in this figure because of its highly reduced scale. Both, however, did not displace the bottom of the Bl (= the upper surface of the Ryoke basement).
4. BSB is inferred to be formed and developed by eastward movements of the hanging wall (Ryoke basement), strike-slip on the MTL and downward on the AF. The BCFS was secondarily formed in eastward moving sediments.


These results are almost concordant with the model of a strike-slip basin proposed by Yamakita and Ito (1999), which assume a not vertical but moderately dipping strike-slip fault with a releasing bend. This model ignored the effects of secondary listric normal faults in sediments, dipping toward the main oblique normal fault forming the oblique ramp, but they contributed to the development of BSB to some extent. Their effects, however, are rather small as indicated by the fact that they did not displace the basement.

Keywords: Gravity analysis, Beppu Bay, MTL, Asamigawa Fault, Beppu Bay Central Fault

SCG68-09

Room:502

Time:April 30 11:15-11:30



Structure and evolution of the lower crust constrained from alkaline basalts and xenoliths in southwest Japan

IGATA, Eri^{1*} ; OMORI, Soichi² ; IWAMORI, Hikaru³

¹Department of Earth and Planetary Sciences, Tokyo institute of technology, ²The open university of Japan, ³Japan Agency for Marine-Earth and Technology

The continental crust is unique to the Earth and, in spite of its small mass, is geochemically an important reservoir, concentrating about half of the radiogenic heat producing elements. In order to understand the differentiation history of the Earth, it is essential to decipher how and when the continental crust has been formed.

In contrast to the upper continental crust, which has been well surveyed with direct means, the lower continental crust is largely unknown in terms of composition, mineralogy and age distribution of formation. In this study, we investigate the xenoliths that have been derived possibly from the lower crust, in terms of petrology (mineral assemblages and their composition, modal abundances and bulk composition) and geochronology (zircon U-Pb age dating of both the xenoliths and the host basalt). Study area is located on the Kibi Plateau in southwest Japan. The xenoliths are classified into 4 types by petrography and EPMA analysis; Type 1, pyroxenite; Type 2, eclogitic gabbro; Type 3, anorthosite; Type 4, kyanite/garnet-bearing felsic granulite. Pseudosection and mineral stability analyses were performed by a thermodynamic program "Perplex". These analyses have revealed that pyroxenite is stable beneath the Moho and its seismic velocity estimated from pseudosection ranges from 7.23 to 7.65 km/s. Therefore pyroxenite corresponds to the olivine-pyroxenite layer under the Moho. Gabbro was formed at 8-10 kbar and 873-940 K and its velocity is higher than pyroxenite. Granulite is stable at 7.5- kbar and about 1000 K and its velocity is very low. In spite of the density gap between the eclogitic gabbro and granulite (3900 and 2740 kg/m³), the analysis suggests that the felsic granulite exhibits a higher equilibration pressure and may even underlay the eclogitic gabbro. Moreover, composition of the lower crust becomes more felsic than previously thought.

Ages of the eclogitic gabbro xenolith and the host basalt coincide, showing 70 Ma. It is argued that the subducted sediments (now appear as felsic granulites) were subducted and underplated to the bottom of the lower crust during the last 30 million years or so. If such a mechanism operates worldwide, then the continental crust may have an intermediate to felsic composition even without a hypothetical process of lower crustal delimitation.

The authors thank T. Hirata, K. Tani, JI. Kimura and Q. Chang for U-Pb zircon age dating.

Three-dimensional seismic velocity structure beneath East Asia using adjoint tomography

MIYOSHI, Takayuki^{1*} ; OBAYASHI, Masayuki¹ ; TONO, Yoko¹ ; TSUBOI, Seiji¹

¹JAMSTEC

East Asia is the complicated region where tectonic plates meet. In many studies, travel-time tomography based on ray theory has clarified three-dimensional (3D) velocity structure of the Earth. On the other hand, the recent studies show waveform inversion based on wave theory can construct realistic 3D structure (e.g. Obayashi et al., 2010 SSJ; Miyoshi et al., 2013 SSJ). In the present study, we have inferred 3D heterogeneous structure precisely beneath the eastern Asia region by using adjoint tomography method. We selected 161 earthquakes ($M > 5.5$, half duration < 5 second) occurred in the region based on Global CMT catalog. Displacement seismograms used in this study were recorded at broadband seismic stations in the region. The average number of stations used in inversion is about 180. Theoretical waveforms were calculated using the spectral-element method (Komatitsch and Tromp, 2001). We used GAP-P2 mantle tomography model (Obayashi et al., 2009) as an initial 3D model of inversion. Both observed and theoretical waveforms were filtered between 12.5 and 100 second to extract time windows of P- and S-waves, and between 30 and 150 second to extract time windows of surface waves. We applied adjoint method (Liu and Tromp, 2006) for calculating the misfit kernel, which is related to velocities, and performed inversion by using the steepest descent method. The parallel computing of theoretical waveforms and misfit kernels were used 256 CPU cores of supercomputers, such as K computer at Riken. The computing time was required 0.1 million CPU hours in each iteration. We have iterated four times on inversion. The VR value was improved about 10% by using the revised model. The V_p and V_s of improved model showed a few percent slower than the initial model. The ratios of the velocity perturbation show slightly large value than the initial model at a depth of 100 km in a wide area of the eastern Asia region. Acknowledgements: We used F-net seismograms of the National Research Institute for Earth Science and Disaster Prevention. This study was supported by the strategic Programs for Innovative Research "Field 3" Advanced prediction Researches for Natural Disaster Prevention and Reduction.

Keywords: adjoint method, tomography, velocity structure, East Asia

Seismological evidence for a transition from "hydrous" oceanic crust to typical oceanic crust in the Lau back-arc basin

ARAI, Ryuta^{1*}; DUNN, Robert¹

¹University of Hawaii

The Lau back-arc basin, associated with subduction of the Pacific plate beneath the Indo-Australian plate at the Tonga Trench, provides a superb study area to understand the interaction between plate subduction and back-arc spreading: Subducting oceanic lithosphere induces mantle corner flow within the mantle wedge above the subducting plate and releases a large amount of water and other elements into this wedge, producing heterogeneous chemical compositions and fluid gradients beneath the back-arc basin. While petrological studies suggest that the heterogeneity in the mantle source composition, mainly caused by slab-derived fluids, plays an important role for melt supply to the back-arc ridges, variations in thickness and internal structure of crust formed along back-arc ridges are poorly documented. On the basis of seismic tomography analyses, we present a structural model of crust formed along the Eastern Lau Spreading Center within the Lau back-arc basin as evidence for a transition from a "hydrous" type of oceanic crust to a more typical oceanic crust. The seismic data indicate that as the back-arc spreading center moved away from the active arc, the crust thinned from 8-9 km to ~7 km, the lower crust changed from high P wave velocity values (7.2-7.4 km/s) to typical values for oceanic crust (7.0-7.2 km/s), and the upper-crustal volcanic layer changed from a thick low-velocity layer to a thinner layer with more typical wave speeds. The seismic results, in combination with other geophysical and geochemical data, suggest that crustal formation along the ELSC is strongly controlled by the influence of slab water: When a spreading center is near the active arc, water from the downgoing slab is entrained in the melting zone beneath the back-arc ridges where it enhances melting. Thereafter, the water enhances crustal differentiation within sub-ridge magma chambers. This creates an anomalous "hydrous" form of oceanic crust with a thick felsic volcanic layer and a mafic/ultramafic lower crust - features that are not typically observed in crust formed at mid-ocean ridges. The Lau basin has a zoned structure with an abrupt transition from this type of oceanic crust to more typical oceanic crust, which resulted from a rapid change in the influence of slab water as the ridge moved away from the arc. The unique geodynamic setting of the Lau basin, such as proximity of the back-arc ridges to the volcanic arc (<100 km), the relatively low subduction angle of the slab (~45 degrees), and the fast subduction rate at the Tonga trench (>20 cm/yr), probably operate to effectively deliver slab-derived water far beyond the volcanic arc to the back-arc ridges and produce this "hydrous" oceanic crust in the back-arc basin. The abundance and high rate of production of the "hydrous" crust suggests that such crust may make up a significant proportion of the arc-like crust that forms continents.

Keywords: Back-arc basin, Crustal differentiation, Oceanic crust, Slab water, Seismic tomography, Eastern Lau Spreading Center

Tectonic province of the northern Fossa Magna region based on the crustal movement and seismic activity

IMAI, Yuki^{1*} ; TAKEUCHI, Akira²

¹Graduate school of Science and Engineering for Education, University of Toyama, ²Graduate school of Science and Engineering for Research, University of Toyama

The northern Fossa Magna region is located in an area where the Niigata-Kobe tectonic zone (Sagiya *et al.*, 2000) and geological strain concentration zone along the eastern margin of the Japan Sea (Okamura, 2002) is duplicated. In the past, inland crustal earthquakes, such as the Zenkoji earthquake in 1847 (M7.4) and the Niigata-ken Chuetsu earthquake in 2004 (M6.8), have occurred in this region. In order to discuss the current tectonics of the northern Fossa Magna region that has active crustal movement and seismic activity, it is necessary to clarify the characteristics of "tectonic province". The purpose of this paper is to reveal a detailed three-dimensional tectonic province model from the crustal movement and seismic activity of this region in the recent years. In order to clarify the characteristics of crustal movement and seismic activity, we have analyzed the GEONET observation data (from October 2007 to March 2011) using the GAMIT 10.4 software, and made the E-W cross-sectional view of the JMA hypocenter data.

The horizontal strain distribution for three and a half years just before the 2011 Tohoku-Oki earthquake shows that strain concentration zone with NW-SE directional contraction extends from the Niigata plain to the Matsumoto basin continuously. Moreover, the eastern margin of this zone corresponds roughly to the position of the Shibata-Koide tectonic line (Yamashita, 1970) running NNE-SSW direction in the eastern margin of the Niigata plain. The strain rate in the Echigo mountain range is smaller than in the Niigata plain. Takeuchi (1999) showed tectonic province based on the activity and characteristics of active faults. The strain distribution revealed from GPS data corresponds approximately to active faults provinces. The large and small strain region corresponds approximately to the reverse fault province (Shin'etsu Niigata sedimentary basin) and strike-slip faults province (Central upheaval zone and Echigo mountain range) respectively. Focusing on the depth distribution of the seismogenic layer in the E-W cross-section, the depth of the lower limit of seismogenic layer is shallow ($D = 10-15$ km) in the strike-slip province but is deeper ($D = 20-30$ km) in the reverse fault province. The seismogenic layer is located beneath the low P-wave velocity zone corresponding to the thick sediments layer in the sedimentary basin.

According to the above results, there is obvious spatial variation of the depth of seismogenic layer and strain distribution at the boundary of the sedimentary basin and Central upheaval zone. It is conceivable that two different tectonic provinces are adjacent along the tectonic boundary where characteristics of the crustal activity are changing greatly. The moderate-large crustal earthquakes around the northern Fossa Magna, such as the Zenkoji earthquake in 1847, the Niigata-ken Chuetsu earthquake in 2004 and the Nagano-ken Hokubu earthquake in 2011, have occurred on or near the tectonic boundary. Stress concentration is likely to occur due to large changing of the physical properties in the tectonic province boundary, and a large crustal earthquake tend to occur at the tectonic province boundary than at the inside province.

Keywords: tectonic province, northern Fossa Magna, crustal movement, seismic activity, seismogenic layer

Results of 2013 Off-Kanazawa and Noto peninsula survey for the integrated research project on seismic and tsunami hazard

SATO, Hiroshi^{1*} ; ISHIYAMA, Tatsuya¹ ; SHIRAIISHI, Kazuya² ; ABE, Susumu¹ ; KATO, Naoko¹ ; IWASAKI, Takaya¹

¹Earthquake Research Institute, Univ. Tokyo, ²JGI. Inc.

To estimate Tsunami and seismic hazards along the coastal area of Sea of Japan, more detailed survey to identify source faults are needed. A new research project funded by MEXT named "the integrated research project on seismic and tsunami hazards around the Sea of Japan" began in FY 2013. To obtain the information of source faults, we performed deep seismic reflection profiling off-Kanazawa and Noto area in the central part of Honshu, Japan. The source faults were estimated together with the results of seismic sections in the epicentral area of the 2007 Noto peninsula earthquake (Sato et al., 2007, BERI). We used two vessels; a gun-ship with 3020 cu. inch air-gun and a cable-ship with a 2-km-long, streamer cable with 156 channels and 480 cu. inch air-gun. Common-mid point reflection data were acquired along 4 seismic lines with total 245 km in length. The seismic profiles portray the reactivation of normal faults, which formed during syn-rift periods, associated with the opening of the Sea of Japan. 2007 Noto peninsula earthquake occurred by the oblique motion on source fault dipping 60 degrees, which is favorable normal faulting. The back arc side of the SW-Japan arc experienced NS trending shortening deformation in the latest Miocene. From the Noto peninsula to the west, undeformed Pliocene sediments covers folded Miocene. Some normal faults reactivated as active strike-slip and reverse faults in Quaternary. The survey results contributed to construct source faults models of Tsunami and seismic hazards estimation.

Keywords: Sea of Japan, source fault, crustal structure, seismic reflection profiling, Off-Kanazawa, Off-Noto Peninsula

Results of 2013 Off-Joetsu survey for the research project on seismic and tsunami hazards around the Sea of Japan

KATO, Naoko^{1*} ; SATO, Hiroshi¹ ; ISHIYAMA, Tatsuya¹ ; SHIRAIISHI, Kazuya² ; ABE, Susumu² ; KURASHIMO, Eiji¹

¹Earthquake Research Institute, Univ. of Tokyo, ²JGI. Inc.

To estimate Tsunami and seismic hazards along the coastal area of Sea of Japan, more detailed survey to identify source faults are needed. A new research project funded by MEXT named "the integrated research project on seismic and tsunami hazards around the Sea of Japan" began in FY 2013. To obtain the information of source faults, we performed deep seismic reflection profiling off-Joetsu area in the central part of Honshu, Japan. The seismic lines were located in the offshore extension previous onshore seismic lines forming onshore-offshore integrated seismic lines. We used two vessels; a gun-ship with 3020 cu. inch air-gun and a cable-ship with a 2-km-long, streamer cable with 156 channels and 480 cu. inch air-gun. Common-mid point reflection data were acquired along 3 seismic lines. Two offshore seismic lines are connected to the onshore seismic sections. The survey area consists of stretched continental crust, such as Noto peninsula and Sado island, and failed rift area with large amount of large mafic intrusive rocks, such as Sado strait and Toyama trough. Stretched continental area is marked by densely distributed syn-rift normal faults. On the other hand, in the Sado strait and Toyama trough, fault-related folds were developed, which show small amount of vertical displacement. Along the boundary between continental crust area and oceanic crust, thrusts with rift axis vergent well imaged by seismic reflection profiles. The survey results contributed to construct source faults models of Tsunami and seismic hazards estimation.

Onshore offshore, deep seismic survey across the Toyama trough

ISHIYAMA, Tatsuya^{1*} ; KATO, Naoko¹ ; SATO, Hiroshi¹ ; SHIRAIISHI, Kazuya² ; ABE, Susumu² ; TAKEDA, Tetsuya³ ; KURASHIMO, Eiji¹

¹Earthquake Research Institute, University of Tokyo, ²R&D Department, JGI, Inc, ³NIED

To estimate Tsunami and seismic hazards along the coastal area of Sea of Japan, more detailed survey to identify source faults are needed. A new research project funded by MEXT named “ the integrated research project on seismic and tsunami hazards around the Sea of Japan ” began in FY 2013. To obtain the information of source faults, we performed onshore offshore deep seismic reflection profiling across the Toyama trough. The seismic line extends from Unazuki at the foot hill of the Hida mountains to the shoreline for 15 km and connected with bay cable of 3 km in length. Seismic signals produced by four vibroseis trucks were recorded by onshore receivers and bay cable. For offshore, we used two vessels; a gun-ship with 3020 cu. inch air-gun and a cable-ship with a 2-km-long, streamer cable with 156 channels and 480 cu. inch air-gun. The P-wave velocity profile by refraction tomography, suggests that the upper surface of Vp 5 km/sec is located 5 km below the mean sea level at the Toyama trough. Vertical offset of Vp 5 km/sec layer is about 8 km. Trough fill sediments beneath the Kurobe alluvial fan show northward dipping. Beneath the apex of this fan, velocity profile and reflection profile suggest the existence of south dipping thrust at 4 km in depth. This fault extends northward as a blind thrust. The seismic section suggests the reverse fault at the northern edge of Toyama trough. Based on the distribution of 5e coastal terraces along the southern part of Noto peninsula, the reverse fault played a significant role for the uplift of Noto peninsula. Such basin structure is analogous to the Niigata sedimentary basin, and northern Fossa magna basin. The survey results contributed to construct source faults models of Tsunami and seismic hazards estimation.

Structure analysis of the Ryukyu arc by the receiver function

ARASHIRO, Yasuhisa^{1*} ; NAKAMURA, Mamoru¹

¹Graduate School of Science,Ryukyu University

The Ryukyu arc have converted plate boundary and back arc basin. The volcanic front in Tokara Islands is the main volcanism. Moreover, The activity of shaped Trough is supposed (Kimura, 1985). In addition, by the survey of igneous activity, the Okinawa Trough have upper flow mantle. Analyzed by receiver function in the Ryukyu arc, McCormack et al., (2013) ware anisotropy structure of slab beneath F-net station.

The Ryukyu arc have converted plate boundary and back arc basin. The volcanic front in Tokara Islands is the main volcanism. Moreover, The activity of shaped Trough is supposed (Kimura, 1985). In addition, by the survey of igneous activity, the Okinawa Trough have upper flow mantle. Analyzed by receiver function in the Ryukyu arc, McCormack *et al.*, (2013) ware anisotropy structure of slab beneath F-net station.

However, it was unknown how changed the structure in the slab and wedge mantle structure changed in the subducting direction. Therefore we clarified I sank by making a receiver functional analysis section at right angles to an errand, the trench including a seismometer record in addition to F-net broadband seismometer record in a short period of the Japan Meteorological Agency,(JMA) and how a Slavic angle changed into the direction.

In the receiver function analysis, we use 8 of short-period seismograph by JMA and 3 of broadband seismometers by NIED F-NET established in central Ryukyu . An analysis period is 2002 to 2013. and used 113 remote earthquake events more than M6.0 for analysis.

In receiver functional analysis, the discontinuity imaging depth is as same as JMA , in Okinawa-honto beneath 40km.

Keywords: receiver function, Ryukyu arc, mantle wedge