

SIT03-01

会場:418

時間:4月29日 09:00-09:15

## Effect of the upper mantle structure on the Moho geometry Effect of the upper mantle structure on the Moho geometry

TENZER, Robert<sup>1\*</sup> ; CHEN, Wenjin<sup>1</sup>  
TENZER, Robert<sup>1\*</sup> ; CHEN, Wenjin<sup>1</sup>

<sup>1</sup>School of Geodesy and Geomatics, Wuhan University

<sup>1</sup>School of Geodesy and Geomatics, Wuhan University

We investigate the effect of the lateral density structure within the upper (most) mantle on the Moho geometry. The gravimetric forward and inverse modeling methods are applied to determine the Moho depths using the gravity data corrected for major known anomalous density structures within the Earth crust. In our numerical experiment we compute and compare the Moho geometry determined using uniform and laterally varying models of the Moho density contrast. The laterally varying model of the Moho density contrast incorporates the information on the upper mantle lateral density structure taken from the CRUST1.0 global crustal model. For the uniform density contrast model, the constant value of the Moho density contrast is determined based on minimizing the spatial correlation between the gravity data and the Moho geometry. Except for the upper (most) mantle, the deeper heterogeneous mantle density structures including the core-mantle boundary zone are not taken into consideration due to the absence of a reliable 3-D density model of the whole mantle. The numerical results revealed that the consideration of the upper mantle density structure improves the fit of the gravimetric solution with the seismic Moho model.

キーワード: crust, density, gravity, mantle, Moho  
Keywords: crust, density, gravity, mantle, Moho

SIT03-02

会場:418

時間:4月29日 09:15-09:30

## 剪断変形による部分溶融ペリドタイトの電気伝導度異方性 Conductivity anisotropy of partial molten peridotite under shear deformation

Zhang Baohua<sup>1</sup>; 芳野 極<sup>1\*</sup>; 山崎 大輔<sup>1</sup>  
ZHANG, Baohua<sup>1</sup>; YOSHINO, Takashi<sup>1\*</sup>; YAMAZAKI, Daisuke<sup>1</sup>

<sup>1</sup> 岡山大学地球物質科学研究センター

<sup>1</sup>Institute for Study of the Earth's Interior, Okayama Univ.

Recent ocean bottom magnetotelluric investigations have revealed a high-conductivity layer (HCL) with high anisotropy characterized by higher conductivity values in the direction parallel to the plate motion beneath the southern East Pacific Rise (Evans et al., 2005) and beneath the edge of the Cocos plate at the Middle America trench offshore of Nicaragua (Naif et al., 2013). These geophysical observations have been attributed to either hydration (water) of mantle minerals or the presence of partial melt. Currently, aligned partial melt has been regarded as the most preferable candidate for explaining the conductivity anisotropy because of the implausibility of proton conduction (Yoshino et al., 2006).

In this study, we report development of the conductivity anisotropy of partial molten peridotite in three directions parallel and normal to shear on the shear plane, and perpendicular to the shear plane as a function of time and shear strain. Starting samples were pre-synthesized partial molten peridotite (Fe-free and Fe-bearing systems), showing homogeneous melt distribution. The Fe-free and Fe-bearing partially molten peridotite samples were deformed in simple shear geometry at 1 GPa and 1523 and 1723 K, respectively, in a DIA-type apparatus with uniaxial deformation facility. Conductivity of the partially molten peridotite parallel to the shear direction was initially identical to that normal to shear. However, shear-parallel conductivity increased by more than one order of magnitude after the initiation of shear by piston advancement. Shear-parallel conductivity then stayed constant for the duration of the experimental run. On the other hand, conductivity normal to the shear direction on the shear plane remained constant, whereas conductivity perpendicular to the shear plane decreased gradually after initiation of shear and finally close to that of olivine. Conductivity difference between parallel and normal to shear direction reached one order, which is equivalent to that observed beneath asthenosphere. In contrast, such anisotropic behavior was not found in the melt-free samples, suggesting that development of the conductivity anisotropy was generated under shear stress.

Microstructure of the deformed partial molten peridotite shows partial melt tends to preferentially locate grain boundaries parallel to shear direction, and forms continuously thin melt layer sub-parallel to the shear direction, whereas apparently isolated distribution was observed on the section perpendicular to the shear direction. The resultant melt morphology can be approximated by tube like geometry parallel to the shear direction. This observation suggests that the development of conductivity anisotropy is caused by the realignment of partial melt (forming tube-like melt) parallel to shear direction in the silicate matrix.

In conclusion, the high anisotropy of conductivity in the direction of plate motion can be well explained by anisotropic interconnection of melt in partially molten rocks at the top of asthenosphere, but not hydration of nominally anhydrous minerals. Therefore, our results provide the direct experimental evidence for supporting these geophysically observed high-conductivity anisotropy at the LAB and verify the validity of partial melting hypothesis (Yoshino et al., 2006; Naif et al., 2013).

### References

- Evans, R. L. et al. Geophysical evidence from the MELT area for compositional controls on oceanic plates. *Nature* 437, 249?252 (2005).  
Naif, S., Key, K., Constable, S. & Evans, R. L. Melt-rich channel observed at the lithosphere?asthenosphere boundary. *Nature* 495, 356?359 (2013).  
Yoshino, T., Matsuzaki, T., Yamashita, S. & Katsura, T. Hydrous olivine unable to account for conductivity anomaly at the top of the asthenosphere. *Nature* 443, 973?976 (2006).

キーワード: 部分溶融, アセノスフェア, 電気伝導度, 上部マントル, 異方性, 剪断変形

Keywords: partial melting, asthenosphere, electrical conductivity, upper mantle, anisotropy, shear deformation

SIT03-03

会場:418

時間:4月29日 09:30-09:45

## 2点間波形解析による北米大陸の表面波位相速度マップの復元 Surface-wave phase velocity maps of North America with inter-station waveform analysis

浜田 広太<sup>1\*</sup>; 吉澤 和範<sup>2</sup>  
HAMADA, Kouta<sup>1\*</sup>; YOSHIZAWA, Kazunori<sup>2</sup>

<sup>1</sup> 北海道大学大学院理学院, <sup>2</sup> 北海道大学大学院理学研究院

<sup>1</sup>Graduate School of Science, Hokkaido University, <sup>2</sup>Faculty of Science, Hokkaido University

The western United States encompasses a variety of tectonic features, including regions with east-west extension, volcanic areas and relatively stable cratonic regions.

In the last decade, the Transportable Array (USArray) has been installed throughout the U.S., and these waveform data have facilitated a variety of tomographic studies in this region using body and surface waves, and ambient noise analysis making the most of the high-density seismic network.

In this study, we have developed a new method of fully non-linear waveform fitting to measure inter-station phase velocities, using the Neighborhood Algorithm (NA) as a global optimizer. This algorithm searches for model parameters to fit two observed waveforms on a common great-circle path by perturbing the phase term of the fundamental-mode Love and Rayleigh waves. We have employed the reliability parameter, which represents how well the waveforms at two stations can be fitted in a time-frequency domain. This parameter is used as a data selection criterion for the subsequent step of phase velocity mapping.

The method has been applied to observed waveform data of the USArray from 2007 to 2010, and we could collect a large-number of phase speed data (over 45000 for Rayleigh and 15000 for Love) in a period range from 30 and 200 seconds, at short distances less than 1000 km. The phase velocity models for Rayleigh and Love waves indicate good correlation on large scales with the recent tomographic maps derived from different approaches for inter-station phase velocity measurements (Foster et al., 2013); e.g., significant slow velocity anomaly in volcanic regions in western United States and extremely fast anomaly in the cratonic region in the longer period range, which implies the robustness of such tectonic features as well as the validity of our new measurement technique. The current method can be expanded for the measurements of inter-station higher-mode phase velocities, which will be of great help in enhancing the vertical resolution of the 3-D shear wave models.

Keywords: surface wave, phase velocity, tomography, North America

SIT03-04

会場:418

時間:4月29日 09:45-10:00

## Diffusion to dislocation creep transition in the upper mantle inferred from silicon grain boundary diffusion rates

## Diffusion to dislocation creep transition in the upper mantle inferred from silicon grain boundary diffusion rates

FEI, Hongzhan<sup>1\*</sup>; KATSURA, Tomoo<sup>2</sup>; KOIZUMI, Sanae<sup>3</sup>; SAKAMOTO, Naoya<sup>4</sup>; HASHIGUCHI, Minako<sup>5</sup>; YURIMOTO, Hisayoshi<sup>5</sup>; YAMAZAKI, Daisuke<sup>1</sup>

FEI, Hongzhan<sup>1\*</sup>; KATSURA, Tomoo<sup>2</sup>; KOIZUMI, Sanae<sup>3</sup>; SAKAMOTO, Naoya<sup>4</sup>; HASHIGUCHI, Minako<sup>5</sup>; YURIMOTO, Hisayoshi<sup>5</sup>; YAMAZAKI, Daisuke<sup>1</sup>

<sup>1</sup>Institute for Study of the Earth's Interior, Okayama University, Japan, <sup>2</sup>Bayerisches Geoinstitut, University of Bayreuth, Germany, <sup>3</sup>Earthquake Research Institute, The University of Tokyo, Japan, <sup>4</sup>Isotope Imaging Laboratory, CRI, Hokkaido University, Japan, <sup>5</sup>Department of Natural History Sciences, Hokkaido University, Japan

<sup>1</sup>Institute for Study of the Earth's Interior, Okayama University, Japan, <sup>2</sup>Bayerisches Geoinstitut, University of Bayreuth, Germany, <sup>3</sup>Earthquake Research Institute, The University of Tokyo, Japan, <sup>4</sup>Isotope Imaging Laboratory, CRI, Hokkaido University, Japan, <sup>5</sup>Department of Natural History Sciences, Hokkaido University, Japan

The majority of the dynamical processes in the upper mantle are controlled by creep of minerals. Dislocation creep causes non-Newtonian viscosity and seismic anisotropy whereas diffusion creep causes Newtonian viscosity and no seismic anisotropy. Determination of deformation mechanism in the upper interior is thus essential to understand mantle dynamics. Previous deformation studies on olivine suggested that the shallow regions of the upper mantle should be dominated by dislocation creep and the deeper regions dominated by diffusion creep [Karato, 1992; Karato and Wu, 1993; Hirth and Kohlstedt, 2003]. However, recent study [Fei et al., 2013] demonstrated that those deformation experiments largely misunderstood the creep rate due to the experimental difficulties. Since the creep of olivine is controlled by silicon diffusion, we measured silicon grain-boundary diffusion coefficient in Mg-olivine aggregates as a function of pressure, temperature, and water content. The activation energy, activation volume, and water content exponent are found to be 240–260 kJ/mol,  $1.8 \pm 0.2 \text{ cm}^3/\text{mol}$ , and  $0.22 \pm 0.05$ , respectively. Together with the silicon lattice diffusion data [Fei et al., 2012; 2013], our results predict the diffusion to dislocation creep transition in the upper mantle, which is in contrast with the previously considered model. In the asthenosphere, dislocation creep should dominate because of the high temperature. In the lithosphere, diffusion creep dominates in shallow regions and dislocation creep dominates in deeper parts. The seismic anisotropy jumps at mid-lithosphere discontinuity beneath continents and at Gutenberg discontinuity beneath oceans are caused by the transition from diffusion to dislocation creep. The weak anisotropy in cold lithospheres could be attributed to the fossil anisotropy formed at the spreading ridges. Dominance of diffusion creep in upper lithosphere accounts for the Newtonian rheology suggested by postglacial rebound.

Fei et al., *EPSL* **345**, 95–103 (2012).

Fei et al., *Nature* **498**, 213–215 (2013).

Hirth and Kohlstedt, *Geophys. Monogr.* **138**, 83–105 (2003).

Karato, *JGR* **19**, 2255–2258 (1992).

Karato and Wu, *Science* **260**, 771–778 (1993).

キーワード: diffusion creep, dislocation creep, upper mantle, silicon, grain boundary diffusion, deformation mechanism

Keywords: diffusion creep, dislocation creep, upper mantle, silicon, grain boundary diffusion, deformation mechanism

SIT03-05

会場:418

時間:4月29日 10:00-10:15

## Aluminum incorporation into phase A - a new hydrous silicate in the deep upper mantle Aluminum incorporation into phase A - a new hydrous silicate in the deep upper mantle

蔡闊<sup>1\*</sup>; 井上徹<sup>1</sup>; 藤野清志<sup>1</sup>; 大藤弘明<sup>1</sup>; 栗林貴弘<sup>2</sup>; 坂本尚義<sup>3</sup>

CAI, Nao<sup>1\*</sup>; INOUE, Toru<sup>1</sup>; FUJINO, Kiyoshi<sup>1</sup>; OHFUJI, Hiroaki<sup>1</sup>; KURIBAYASHI, Takahiro<sup>2</sup>; YURIMOTO, Hisayoshi<sup>3</sup>

<sup>1</sup> 愛媛大学地球深部ダイナミクス研究センター, <sup>2</sup> 東北大学大学院理学研究科地学専攻, <sup>3</sup> 北海道大学理学研究院

<sup>1</sup>Geodynamics Research Center, Ehime University, <sup>2</sup>Institute of Mineralogy, Petrology and Economic Geology, Graduate School of Science, Tohoku Universi, <sup>3</sup>Division of Earth and Planetary Sciences, Hokaido University

A new aluminum bearing hydrous silicate was found in the experiments under the deep upper mantle conditions, using phase A ( $Mg_7Si_2O_8(OH)_6$ ) and  $Al(OH)_3$  as the starting materials. Using electron probe micro analysis (EPMA) and secondary ion mass spectrometry (SIMS), the composition was determined to be very close to  $Mg_{5.5}AlSi_2O_8(OH)_6$ , which contained about 12 wt % of water. Almost pure phase was obtained in the subsequent experiments. The powder x-ray diffraction pattern and transmission electron diffraction patterns showed a hexagonal structure, with an abnormal large c axis. This new phase has similar stability region with phase A. At lower pressure and higher temperature, it breaks down into Chondrodite + Garnet + Brucite + Fluid, while at higher pressure and higher temperature, it breaks down into Al-superhydrous phase B + Garnet + Brucite + Fluid.

Besides, present study shows that phase A coexists with this aluminum bearing hydrous phase, with a small amount of aluminum (<1 wt%) incorporated into phase A structure, which predicts that phase A can preserve only trace of aluminum.

According to Inoue's unpublished data, aluminum can easily incorporate into some dense hydrous magnesium silicates and form aluminum bearing hydrous phases such as phase B, superhydrous phase B, and even perovskite. However, rather than aluminum bearing phase A, the present study shows a small amount of aluminum incorporation into phase A structure, and an appearance of a new aluminum bearing hydrous phase, with the composition very similar to phase A but structure very different from it. Further investigations are needed to clarify these two phases.

キーワード: Phase A, Aluminum incorporation, hydrous phase, upper mantle

Keywords: Phase A, Aluminum incorporation, hydrous phase, upper mantle

SIT03-06

会場:418

時間:4月29日 10:15-10:30

## 長白山火山とその付近の深発地震との関係 Changbai intraplate volcanism and deep earthquakes in Northeast Asia

趙 大鵬<sup>1\*</sup>; Tian You<sup>1</sup>  
ZHAO, Dapeng<sup>1\*</sup>; TIAN, You<sup>1</sup>

<sup>1</sup> 東北大学・理  
<sup>1</sup>Tohoku University, Department of Geophysics

The origin of the intraplate volcanoes in Northeast Asia is considered to be associated with upwelling of hot and wet asthenospheric materials in the big mantle wedge above the stagnant Pacific slab in the mantle transition zone. Among these intraplate volcanoes, Changbai is the largest and most active one, and very deep earthquakes (500-600 km depths) in the Pacific slab under East Asia occur ~300 km to the east of the Changbai volcano. Integrating the findings of geophysical, geochemical and petrologic studies so far, we suggest a link between the Changbai volcanism and the deep earthquakes in the Pacific slab. Many large shallow earthquakes occurred in the Pacific plate in the outer-rise areas close to the oceanic trench, and seawater may enter down to the deep portion of the oceanic lithosphere through the active normal faults which generated the large outer-rise earthquakes. The seawater or fluids may be preserved in the active faults even after the Pacific plate subducts into the mantle. Many large deep earthquakes are observed that took place in the subducting Pacific slab under the Japan Sea and the East Asian margin. At least some of the large deep earthquakes are caused by the reactivation of the faults preserved in the subducting slab, and the fluids preserved in the faults within the slab may cause the observed non-double-couple components in the deep earthquake faulting. The fluids preserved in the slab may be released to the overlying mantle wedge through the large deep earthquakes. Because large deep earthquakes occur frequently in the vicinity of the Changbai volcano, much more fluids could be supplied to this volcano than other areas in Northeast Asia, making Changbai the largest and most active intraplate volcano in the region.

### Reference

Zhao, D., Y. Tian (2013) Changbai intraplate volcanism and deep earthquakes in East Asia: A possible link? *Geophys. J. Int.* 195, 706-724.

キーワード: 火山, 深発地震, アジア, スラブ  
Keywords: volcanoes, deep earthquakes, Asia, slab

SIT03-07

会場:418

時間:4月29日 10:30-10:45

## Semiconductor diamond heater (SCD): An innovation for ultrahigh temperature experiments in the Kawai cell

## Semiconductor diamond heater (SCD): An innovation for ultrahigh temperature experiments in the Kawai cell

XIE, Longjian<sup>1\*</sup> ; ITO, Eiji<sup>1</sup> ; YONENDA, Akira<sup>1</sup>

XIE, Longjian<sup>1\*</sup> ; ITO, Eiji<sup>1</sup> ; YONENDA, Akira<sup>1</sup>

<sup>1</sup>Institute for Study of the Earth Interior , Okayama University

<sup>1</sup>Institute for Study of the Earth Interior , Okayama University

We developed the semi-conductor diamond heater in the Kawai high pressure cell. The starting material of the semi-conductor diamond heater is boron(B)-doped burned-graphite. We succeeded to improve the machinability of the B-doped burned-graphite by decreasing porosity. Following is the motivation and the background of the semi-conductor diamond heater project.

It is important to generate extremely high temperature (~3000 °C) in a large sample volume (~0.1mm<sup>3</sup>) in the Kawai apparatus. X-ray transparency is also desirable for in-situ synchrotron analysis. However, any traditional heater used in the Kawai apparatus so far does not satisfy the both requirements simultaneously.

Semiconductor diamond is a candidate material to generate temperatures higher than 3000 °C with low x-ray absorption. Anton Shatskiy (2009) have generated a temperature of 3500 °C by using the semiconductor diamond heater in a large-volume Kawai-type high-pressure apparatus, although their temperature measurement is questionable from a viewpoint of the power-temperature relation. Furthermore, their semi-conducted diamond heater, made of boron and graphite powders, was not machinable and difficult to control the temperatures. It often became unstable at around 1000~1300 °C and impossible to generate higher temperature.

Systematic experiments have done to improve the performance of the semiconductor heater. We used a machinable block of graphite contain 3 wt.% boron as the starting material for the semi-conductor diamond heater. The graphite-diamond transformation started at ~1000-1200 °C at 15 GPa in the Kawai apparatus. After the transformation, we stably generated temperature to 2000 °C. Activation energy of B-doped diamond is about 0.1 eV, which is much lower than that of pure diamond (5.45eV).

### References:

Anton Shatskiy, Daisuke Yamazaki, Guillaume Morard, Titus Cooray, Takuya Matsuzaki et al. , Review of Scientific Instruments 80, 023907 (2009).

キーワード: Semiconductor Diamond Heater, Ultrahigh Temperature, Kawai Cell

Keywords: Semiconductor Diamond Heater, Ultrahigh Temperature, Kawai Cell

SIT03-08

会場:418

時間:4月29日 11:00-11:15

## 最近のグローバルトモグラフィーモデル：我々はどこに向かっているのか？ Recent Global Tomography Models: Where are We Heading for?

竹内 希<sup>1\*</sup>

TAKEUCHI, Nozomu<sup>1\*</sup>

<sup>1</sup> 東京大学地震研究所

<sup>1</sup>Earthquake Research Institute, University of Tokyo

Many high-resolution global tomography models have been obtained and we now have consensus about overall features of 3-D heterogeneous structure in the Earth. Majority of models have been obtained by using conventional ray theory which assumes that scale length of lateral heterogeneities is sufficiently large compared with wavelength of seismic waves analyzed.

Primary efforts in recent studies appear to introduce better theories to overcome resolution limits caused by the above-mentioned assumption. The efforts include delay time tomography with finite frequency kernels (e.g., Montelli et al. 2004, Science; Obayashi et al. 2013, GRL) and waveform tomography with 2-D (e.g., Li and Romanowicz 1996, GJI, Panning and Romanowicz 2004, Science) or 3-D (e.g., Takeuchi 2007, GJI; Takeuchi 2012, EPSL) finite frequency kernels. Waveform tomography with further better theories is also becoming feasible (e.g., Lekic and Romanowicz 2011, GJI; French et al. 2013, Science).

In this presentation, I want to propose another direction to improve resolution: use of a new type of dataset. I will propose to use incoherent part of seismic signals (i.e., scattering waves or coda waves). Scattering waves are sensitive to heterogeneities whose scale length is comparable with wavelength of seismic waves analyzed. Use of such waves therefore should provide new information beyond resolution limit of ray theory. At the time of the presentation, I plan to show feasibility and examples of such analyses to reveal distribution of smaller scale heterogeneities in the subduction zone around Japan.

キーワード: tomography, scattering wave, seismology

Keywords: tomography, scattering wave, seismology

SIT03-09

会場:418

時間:4月29日 11:15-11:30

## Lattice preferred orientation of stishovite in deformation experiment

### Lattice preferred orientation of stishovite in deformation experiment

XU, Fang<sup>1\*</sup> ; YAMAZAKI, Daisuke<sup>1</sup> ; TSUJINO, Noriyoshi<sup>1</sup>  
XU, Fang<sup>1\*</sup> ; YAMAZAKI, Daisuke<sup>1</sup> ; TSUJINO, Noriyoshi<sup>1</sup>

<sup>1</sup>ISEI, Okayama University

<sup>1</sup>ISEI, Okayama University

Seismic observations reveal strong negative anisotropy ( $V_{SV} > V_{SH}$ ) at around 550 km depth in the lower part of mantle transition zone (Visser et al., 2008). The mantle tomography indicates the obvious association of this negative anisotropy with the subduction zones (Panning and Romanowicz., 2006). The observed anisotropy can be caused by lattice preferred orientation (LPO) of constituting material when the material is elastically anisotropic. Majorite and ringwoodite, which are the dominant minerals in this region, are nearly isotropic (Chai et al., 1997; Weidner et al., 1984). On the other hand, stishovite, which may occur in significant amounts in this region derived from the delaminated subducting basaltic layer (Karato et al., 1997) and continental crust (Kawai et al., 2012), shows strong elastic anisotropy ( $V_{SV}/V_{SH}$  is as large as 150%) indicated by the acoustic velocities study (Yoneda et al., 2012) on single crystal of stishovite. Therefore, the LPO of stishovite has a high potential to interpret the seismic anisotropy in the lower part of the transition zone and indicate the geometry of mantle flow.

To investigate the LPO of stishovite, deformation experiments on stishovite were conducted in both simple shear and uni-axial geometry. We prepared starting material of polycrystalline stishovite with grain size of  $\sim 10 \mu\text{m}$  at 12 GPa and 1450 °C in a Kawai-type high-pressure apparatus. Then deformation experiments were carried out at 12 GPa and 1600 °C by Kawai-type apparatus for tri-axial deformation (KATD installed at Tokyo Institute of Technology) and deformation-DIA apparatus (SPEED-Mk. II installed at SPring-8). Sintered diamond piston was used in the uni-axial deformation experiment. Shear strain was  $\sim 1.0$  estimated from the rotation of platinum strain maker after deformation. From the change of sample length, uni-axial tension and compression strain were 0.4 and 0.1 respectively. The microstructure and crystallographic orientation of the deformed samples were investigated by SEM with EBSD.

The EBSD results reveal that the [001] direction of stishovite tends to be parallel to the shear direction. (100) plane, though not so obvious, tends to parallel to the shear plane. The slip system is consistent with rutile TiO<sub>2</sub> (Blanchin and Faisant., 1979), which has the same structure with stishovite. The calculated seismic anisotropy indicates a fast shear wave along shear direction. Polarization anisotropy reported by Visser et al. (2008) can be attributed by a vertical flow and LPO of stishovite in the transition zone. The negative anisotropy along subduction zones in Panning and Romanowicz. (2006) indicates the type A slabs (slabs which penetrate directly into the lower mantle without much deflection in the transition zone) (Karato et al., 2001).

キーワード: stishovite, deformation, LPO

Keywords: stishovite, deformation, LPO

SIT03-10

会場:418

時間:4月29日 11:30-11:45

## 1600~2200 °Cにおける 660 km 地震波不連続面付近のパイロライトの相転移と鉱物化学 Phase transitions and mineral chemistry in pyrolite at 1600-2200C across 660-km seismic discontinuity

石井 貴之<sup>1\*</sup>; 糸谷 浩<sup>1</sup>; 赤荻 正樹<sup>1</sup>  
ISHII, Takayuki<sup>1\*</sup>; KOJITANI, Hiroshi<sup>1</sup>; AKAOGI, Masaki<sup>1</sup>

<sup>1</sup> 学習院大学理学部

<sup>1</sup> Depertment of chemistry, Gakushuin University

It is widely accepted that pyrolite is a model rock which represents the chemical composition of the Earth's upper mantle. Because the post-spinel transition in pyrolite occurs at about 23 GPa along mantle geotherm (e.g. Litasov et al. 2005), it has been accepted that the transition is responsible for the seismic 660-km discontinuity. Slow velocity anomalies by global seismic tomography which may indicate mantle upwelling have been found in the transition zone and the lower mantle, and these regions are higher in temperature than average mantle. To elucidate the origin and dynamics of the mantle plume, informations on phase relations in pyrolite are essential. However, few investigations on phase relations in pyrolite have been made at hot-plume temperatures (1800-2200C) (Hirose, 2002; Nishiyama and Yagi, 2003). In this study, we demonstrated detailed phase equilibrium experiments in pyrolite composition at hot plume conditions.

The starting material was prepared as the oxide mixture in pyrolite composition after McDonough and Sun (1995) excluding minor components (MnO, K<sub>2</sub>O and P<sub>2</sub>O<sub>5</sub>). Quench experiments were made at about 20-28 GPa and 1600-2200C for 2-10 hours using a Kawai-type 6-8 multianvil high-pressure apparatus at Gakushuin University. The starting material was packed with pressure calibrants (MgSiO<sub>3</sub> and pyrope) in a Re multi-sample capsule. A LaCrO<sub>3</sub> heater and a W5%Re-W26%Re thermocouple were inserted in a Cr<sub>2</sub>O<sub>3</sub>-doped MgO pressure medium. Phases of recovered samples were identified with microfocus-Xray diffractometer and SEM-EDS.

The mineral assemblages of MgSiO<sub>3</sub>-rich perovskite (Mpv) + magnesiowustite (Mw) + garnet (Gt) + CaSiO<sub>3</sub>-perovskite (Cpv) and Mpv + Mw + Cpv at 1600-2200C are stable at pressure range of 22-24 GPa and above 24 GPa, respectively. The mineral assemblage of ringwoodite (Rw) + Gt + Cpv at 1600C changes to that of Rw + Mw + Gt + Cpv at 1800-2000C, and Rw disappears perfectly above 2200C. From mass balance calculation of analyzed compositions of the phases, we found that Gt content increases with increasing temperature before and after formation of Mpv. We also calculated the densities in pyrolite at each temperature. The density of average pyrolite mantle (1600C) is higher than pyrolite plume (1800-2200C) across 660-km discontinuity due to increase in Gt content with increasing temperature. Therefore, we conclude that hot-plume ascending nearby 660-km discontinuity has positive buoyancy by the phase transitions.

キーワード: ポストスピネル転移, 660 km 地震波不連続面, マントルプラーム, パイロライト, ポストガーネット転移  
Keywords: post-spinel transition, 660-km seismic discontinuity, mantle plume, pyrolite, post-garnet transition

SIT03-11

会場:418

時間:4月29日 11:45-12:00

## 高圧力条件下での $\delta$ -AlOOH の弾性的性質: マントル遷移層における高 Vs 異常への影響

### Elastic properties of delta-AlOOH under high-pressure: Implications for high Vs anomaly in the mantle transition zone

増野 いづみ<sup>1\*</sup>; 村上 元彦<sup>1</sup>; 大谷 栄治<sup>1</sup>  
MASHINO, Izumi<sup>1\*</sup>; MURAKAMI, Motohiko<sup>1</sup>; OHTANI, Eiji<sup>1</sup>

<sup>1</sup> 東北大学大学院理学研究科地学専攻

<sup>1</sup>Faculty of Science, Tohoku University

delta-AlOOH is a high-pressure polymorph of diaspore (alpha-AlOOH) and boehmite (gamma-AlOOH) (Suzuki *et al.*, 2000). Since delta-AlOOH is identified to be stable from 20 to 120 GPa, and temperatures up to 2300 K, this phase is considered to be a possible carrier and reservoir of water in subducting cold slab into the deep mantle (Ohtani *et al.*, 2001; Sano *et al.*, 2004; 2008). In order to investigate the effect of composition on seismic velocities in subducting slab, it is important to measure the elastic properties of delta-AlOOH at high pressure.

We have conducted high-pressure acoustic-wave velocity measurements of delta-AlOOH using Brillouin spectroscopy and also explored the chemical bonding of delta-AlOOH by Raman spectroscopy at high pressure in a diamond anvil cell. We obtained sharp peaks from transverse acoustic mode ( $V_S$ ) of delta-AlOOH over the entire pressure range explored up to a pressure of 89 GPa. The peaks from longitudinal acoustic mode ( $V_P$ ) of delta-AlOOH were masked by the diamond shear acoustic modes from 35 GPa. The pressure dependence of the aggregate velocities for the delta-AlOOH at 300 K suggests that the hydrogen-bonding symmetrization with the space group changes from  $P2_1nm$  to  $Pnnm$  occurs during compression above 7 GPa. The shear and adiabatic bulk moduli and their pressure derivatives at zero pressure were determined to be  $K_0 = 192.2(8)$  (GPa),  $G_0 = 158.8(3)$  (GPa),  $(dK/dP)_0 = 3.63(6)$ , and  $(dG/dP)_0 = 1.35(6)$  for the pressures above 15 GPa. Raman spectroscopic measurements have shown that the  $B_1$  mode frequencies of  $P2_1nm$  disappeared around 6 GPa and  $A_g$  mode frequencies of  $Pnnm$  appeared above 5.6 GPa, which also indicates the hydrogen-bonding symmetrization around 6 GPa. These results indicate that delta-AlOOH becomes harder by the hydrogen-bonding symmetrization and probably exists as a phase ( $Pnnm$ ) with the symmetric hydrogen bonding in the mantle transition zone and lower mantle.

Shear wave velocities for delta-AlOOH are larger than those of hydrous wadsleyite (by 30 %), hydrous ringwoodite (by 29 %), and majorite (by 29 %). Those of delta-AlOOH are approximately 7 % below those of stishovite. The delta-AlOOH phase thus found to be one of the hardest phases compared to the minerals of mantle transition zone. The existence of delta-AlOOH may contribute to the cause of high  $V_S$  and  $V_P$  anomalies. Shear velocities for sediment containing delta-AlOOH phase are larger than those of pyrolite (by 10 %) and MORB (by 5 %). The subducting slabs often stagnate at the transition zone before reaching the lower mantle. Particularly beneath Korean peninsula, there is a high  $V_S$  anomaly (~2 %) in the lower part of the transition zone (Zhang *et al.*, 2012). The seismic data under the eastern part of northeast China (NEC) also indicates a slight positive anomaly of  $V_S$  (~1 %), but the  $V_S$  value observed around 600 km depth under NEC is ~1 % lower than that beneath Korea. We explain the difference in the  $V_S$  anomalies beneath the NEC and Korea by the amount of sediment containing the delta-AlOOH phase and the stagnating duration. If sediments stagnate at the transition zone before reaching the lower mantle in this region, we can estimate that the higher  $V_S$  anomaly (~1 %) than NEC would correspond to sediments with 13.4 vol% in stagnant slab. The average oceanic crust subduction rate is estimated to be about 8 cm/yr around Japan. Assuming this estimated rate of subduction, the slab stagnation has lasted for at least 30 million years.

Keywords: delta-AlOOH, Brillouin scattering, Raman spectroscopy, subducting slab, high pressure

SIT03-12

会場:418

時間:4月29日 12:00-12:15

## 660-km不連続面を貫通するスラブ内の極深発地震 ( $h>660\text{km}$ ) のメカニズム Mechanisms of ultra-deep earthquakes ( $h>660\text{km}$ ) in a slab penetrating the 660-km discontinuity

深尾 良夫<sup>1\*</sup>; 大林 政行<sup>1</sup>; 吉光 順子<sup>1</sup>

FUKAO, Yoshio<sup>1\*</sup>; OBAYASHI, Masayuki<sup>1</sup>; YOSHIMITSU, Junko<sup>1</sup>

<sup>1</sup> 海洋研究開発機構

<sup>1</sup> IFREE/JAMSTEC


Recent mantle tomography has begun to reveal the characteristic differences between the deep hypocentral distributions associated with stagnant slabs and those associated with penetrating slabs (e.g., Fukao and Obayashi, 2014). We here show that there are differences in focal mechanism as well. Mechanisms of deep shocks within tomographically imaged stagnant slabs (typically in Bonin and Tonga) are characterized by horizontal compression (e.g., Bonnardot et al., 2009). Those within tomographically imaged penetrating slabs (typically in Java and Tonga) are characterized by very steeply dipping compressional axes (e.g., Alpert et al., 2010).

The deepest seismicity is especially active in Tonga, where many shocks occur at depths greater than 660km. Such ultra-deep shocks show in general very unusual mechanisms, typified by nearly vertical tensional axes with a large amount of CLVD component, as demonstrated in Figure 1 (Mechanisms viewed from the side). This figure also shows a remarkable contrast of mechanisms of deepest shocks just above and below the 660km depth. The source region of these ultra-deep shocks ( $h>660\text{km}$ ) is underlain by the greatly deepened post-spinel phase boundary (Niu and Kawakatsu, 1995) so that the source region is at the pre-spinel state while the underlying portion is at the post-spinel state. This situation along with contortion of the slab associated with its interaction with the post-spinel phase boundary (e.g., Cizkova and Bina, 2013) may explain the mechanism change across the 660km depth as observed in Figure 1. We explore the finer velocity structure and hypocentral distribution in the source region by a technique of differential travel time tomography.

キーワード: マントルダイナミクス, トモグラフィー, 深発地震

Keywords: mantle dynamics, tomography, deep earthquakes

Tonga section B



SIT03-13

会場:418

時間:4月29日 12:15-12:30

## A pyrolytic lower mantle with $(\text{Mg}, \text{Fe}^{3+})(\text{Si}, \text{Al}^{3+})\text{O}_3$ perovskite

## A pyrolytic lower mantle with $(\text{Mg}, \text{Fe}^{3+})(\text{Si}, \text{Al}^{3+})\text{O}_3$ perovskite

WANG, Xianlong<sup>1\*</sup> ; TSUCHIYA, Taku<sup>1</sup>  
WANG, Xianlong<sup>1\*</sup> ; TSUCHIYA, Taku<sup>1</sup>

<sup>1</sup>GRC, Ehime University and ELSI, Tokyo Institute of Technology

<sup>1</sup>GRC, Ehime University and ELSI, Tokyo Institute of Technology

To better understand the Earth's lower mantle (LM), thermodynamic properties (TDPs) of LM minerals with Fe and Al dopant should be illustrated more clearly. We have so far reported the TDPs of Fe-bearing MgO,  $\text{MgSiO}_3$  perovskite (Pv) and post perovskite. [1-4] We furthermore study the TDPs of Fe- and Al-bearing Pv, where the internally consistent LSDA+U method and the lattice dynamics method are applied. Two spin states, high (HS) and low spin state, two substitution sites, Mg and Si site, and several possible distribution configurations are considered. In the LM pressure range, HS  $\text{Fe}^{3+}$  substituted at the Mg site with  $\text{Al}^{3+}$  at the adjacent Si site (Fe-Al pair) is the most stable configuration and tends to distribute homogeneously in LM. Furthermore, negative frequency cannot be observed in the Fe-Al pair-bearing Pv, and Al contributes to middle frequency while Fe mainly to low part due to its heavy mass. This indicates that the Fe-Al pair is vibrationally stable. Incorporation of the pair for geophysically relevant concentrations can increase volume of Pv a little and has marginal effects on the TDPs of Pv except for thermal expansivity and Gruneisen parameter. Simulated densities, adiabatic bulk moduli ,and bulk sound velocities show that a composition close to pyrolite is accountable for the reference Earth model.

### References:

- [1] A. Metsue, and T. Tsuchiya, J. Geophys. Res. 116, B08207 (2011).
- [2] A. Metsue, and T. Tsuchiya, Geophys. J. Int. 190, 310 (2012).
- [3] H. Fukui, T. Tsuchiya, and A. Q. R. Baron, J. Geophys. Res. 117, B12202 (2012).
- [4] T. Tsuchiya, and X. Wang, J. Geophys. 118, 83 (2013).

キーワード: First-principles method, Internally consistent LSDA+U, Perovskite, Thermodynamic properties, Pyrolite

Keywords: First-principles method, Internally consistent LSDA+U, Perovskite, Thermodynamic properties, Pyrolite

SIT03-14

会場:418

時間:4月29日 12:30-12:45

## 強い圧縮性を持つ流体中の熱対流に関する線形解析 Linear analysis on the onset of thermal convection of highly compressible fluids

亀山 真典<sup>1\*</sup>  
KAMEYAMA, Masanori<sup>1\*</sup>

<sup>1</sup> 愛媛大学・地球深部研  
<sup>1</sup>GRC, Ehime University

A series of linear stability analysis was performed on the onset of thermal convection in a highly compressible fluid, in order to study the fundamental nature of mantle convection of massive super-Earths in the presence of strong adiabatic compression. We consider the temporal evolution (growth or decay) of an infinitesimal perturbation superimposed to a highly compressible fluid which is in a hydrostatic (motionless) and conductive state in a basally-heated horizontal layer. As a model of pressure-dependence in material properties, we employed an exponential decrease in thermal expansivity and exponential increase in (reference) density with depth. The linearized equations for conservation of mass, momentum and internal (thermal) energy are numerically solved for the critical Rayleigh number as well as the vertical profiles of eigenfunctions for infinitesimal perturbations. The above calculations are repeatedly carried out by systematically varying (i) the dissipation number which measures the effect of adiabatic compression, (ii) the temperature at the top surface and (iii) the magnitude of pressure-dependence in thermal expansivity and reference density.

Our analysis demonstrated that the onset of thermal convection is strongly affected by the adiabatic compression, through modulating the static stability of thermal stratification in the fluid layer. For sufficiently strong adiabatic compression where a sufficiently thick "stratosphere" of stable stratification develops in the layer, for example, the critical Rayleigh number explosively increases with the dissipation number. The explosive changes in the critical Rayleigh number are associated with drastic decreases in the length scales of perturbations both in vertical and horizontal directions. In particular, for very strong adiabatic compression, the vertical motion of fluid is significantly suppressed in a thick "stratosphere", which narrows the incipient convection in a thin sublayer of unstable thermal stratification. In addition, when the effect of adiabatic compression is extremely strong so that the thermal stratification becomes stable in the entire layer, no perturbation is allowed to grow with time regardless of the Rayleigh number and/or the horizontal wavelength. We also found that the effect of adiabatic compression becomes prominent for higher temperature at the top surface of the fluid layer. These findings may imply the crucial importance of adiabatic compression in understanding the dynamics and evolution of the mantles of massive super-Earths, particularly for those orbiting their parent stars very closely.

キーワード: スーパー地球, マントル対流, 断熱圧縮, 熱膨張率  
Keywords: super-Earths, mantle convection, adiabatic compression, thermal expansivity

SIT03-15

会場:418

時間:4月29日 14:15-14:30

部分融解したケイ酸塩から溶融鉄が分離する過程の高温高圧下におけるX線透視を用いたその場観察

**In Situ observation of the Segregation Process of Molten Iron from Partially Molten Silicate using X-ray Radiography**

八木 健彦<sup>1\*</sup>;後藤 弘匡<sup>2</sup>;飯塚 理子<sup>1</sup>;鈴木 昭夫<sup>3</sup>  
YAGI, Takehiko<sup>1\*</sup>; GOTOU, Hirotada<sup>2</sup>; IIZUKA, Riko<sup>1</sup>; SUZUKI, Akio<sup>3</sup>

<sup>1</sup> 愛媛大学地球深部ダイナミクス研究センター, <sup>2</sup> 東京大学物性研究所, <sup>3</sup> 東北大学大学院理学研究科地学専攻

<sup>1</sup>Geodynamics Research Center, Ehime University, <sup>2</sup>Institute for Solid State Physics, University of Tokyo, <sup>3</sup>Department of Earth Science, Tohoku University

溶融鉄が部分融解したケイ酸塩から分離する過程を、5GPa, 1800Kまでの高温高圧下でX線透過法によりその場観察を行った。地球のコアは同様の過程によって生成初期に形成されたと考えられており、その過程の詳細については溶融ケイ酸塩中を液滴が落下するモデルや、固液が混じったケイ酸塩中を浸透流となって溶けた鉄が落下するモデルなどが提案されている。しかし今までの実験では種々異なる結果が示されて、まだ統一的な理解には至っていない。本研究ではMg(OH)<sub>2</sub>、SiO<sub>2</sub>、Feの粉末の一様な混合体を出発物質として、室温で5GPaまで加圧してから、1800Kまで昇温する過程でX線透視によるその場観察を行った。その結果、鉄が凝集して球状になった液滴が落下し、最終的にひとつの大きな玉になる過程が観察された。この観察方法がこのような研究に有力な手段となることが明らかにされた。

キーワード: コア形成過程, 溶融鉄, X線, 高温高圧

Keywords: core formation process, molten iron, x-ray, high pressure and temperature

SIT03-16

会場:418

時間:4月29日 14:30-14:45

## メジャーライト相がマントル対流に与える影響 Influence of majorite on mantle convection

市川 浩樹<sup>1\*</sup>; 亀山 真典<sup>2</sup>; 千秋 博紀<sup>3</sup>; 河合 研志<sup>4</sup>; 丸山 茂徳<sup>5</sup>  
ICHIKAWA, Hiroki<sup>1\*</sup>; KAMEYAMA, Masanori<sup>2</sup>; SENSEHU, Hiroki<sup>3</sup>; KAWAI, Kenji<sup>4</sup>; MARUYAMA, Shigenori<sup>5</sup>

<sup>1</sup> 愛媛大学地球深部ダイナミクス研究センター及び東京工業大学地球生命研究所, <sup>2</sup> 愛媛大学 地球深部ダイナミクス研究センター, <sup>3</sup> 千葉工業大学 惑星探査研究センター, <sup>4</sup> 東京工業大学 地球惑星科学専攻, <sup>5</sup> 東京工業大学 地球生命研究所  
<sup>1</sup>GRC, Ehime University and ELSI, Tokyo Institute of Technology, <sup>2</sup>Geodynamics Research Center, Ehime University, <sup>3</sup>Planetary Exploration Research Center, Chiba Institute of Technology, <sup>4</sup>Department of Earth and Planetary Sciences, Tokyo Institute of Technology, <sup>5</sup>Earth-Life Science Institute, Tokyo Institute of Technology

Influence of  $\text{MgSiO}_3$  majorite on the mantle convection has been investigated by using numerical simulations. According to a first principles study (Yu et al., 2011), wadsleyite decomposes to an assemblage of majorite plus periclase with a large negative Clapeyron slope. Since stability field of majorite is limited at high temperature, downwellings are considered to be unaffected by this phase boundary. On the contrary, the upwelling plumes may be significantly modified by this phase boundary. The asymmetry on upwelling and downwelling caused by the phase transitions may induce strong effects on the thermal evolution and the thermal structure of the mantle.

In this study, we performed 2-D numerical simulations on thermal convection of the mantle incorporating majorite stability field. According to our numerical results, very hot upwelling plumes are strongly influenced by the phase transitions related to majorite. The dynamics of these upwellings are controlled by the release and the absorption of latent heat induced by the transitions as well as interruption of currents due to the large negative Clapayron slope of the transition between wadsleyite and majorite plus periclase.

キーワード: マントル対流, メジャーライト, 相転移  
Keywords: Mantle convection, Majorite, Phase transition

SIT03-17

会場:418

時間:4月29日 14:45-15:00

## Rapid lateral variation of P-wave velocity at the base of the mantle beneath the Western Pacific

### Rapid lateral variation of P-wave velocity at the base of the mantle beneath the Western Pacific

田中 聰<sup>1\*</sup>; 川勝 均<sup>2</sup>; 大林 政行<sup>1</sup>; NECESSAArray プロジェクトチーム<sup>2</sup>

TANAKA, Satoru<sup>1\*</sup>; KAWAKATSU, Hotoshi<sup>2</sup>; OBAYASHI, Masayuki<sup>1</sup>; NECESSAARRAY, Project team<sup>2</sup>

<sup>1</sup>JAMSTEC, <sup>2</sup>ERI, Univ. Tokyo

<sup>1</sup>JAMSTEC, <sup>2</sup>ERI, Univ. Tokyo

We examine P-wave velocity structure at the base of the mantle beneath the Western Pacific, where is the western edge of the Pacific Large-Low Velocity Province (LLVP), by using high-quality seismograms that are provided by the NECESSAArray project. Forward modeling with the reflectivity method is conducted to explain the variation of P-wave travel times as function of epicentral distance near the core shadow zone after station and ellipticity corrections are applied. Additionally PcP-P travel times are also examined to enlarge the survey area. As a result, a rapid variation of P-wave velocity structure at the base of the mantle is detected. Thin (10 to 60 km thickness) and very low velocity (-2 to -6 %) layers at the base of the mantle are intersected with a 100 km thickness and high velocity (+3%) layer, and a slightly fast layer exists at the north of the region with the thin and low velocity layers. Their spatial separations are typically several hundred kilometers and it would be difficult to explain by only a thermal effect. These observations suggest that very complicated thermo-chemical reactions occur near the edge of Pacific LLVP.

キーワード: P-wave velocity, the base of the mantle, the Western Pacific

Keywords: P-wave velocity, the base of the mantle, the Western Pacific

SIT03-18

会場:418

時間:4月29日 15:00-15:15

## 局所的3次元詳細構造のための波形インバージョン手法の開発およびその応用 Methods for inversion of body-wave waveforms for localized three-dimensional seismic structure and an application to D''

河合 研志<sup>1\*</sup>; 小西 健介<sup>2</sup>; ゲラー ロバート<sup>3</sup>; 富士 延章<sup>4</sup>  
KAWAI, Kenji<sup>1\*</sup>; KONISHI, Kensuke<sup>2</sup>; GELLER, Robert J.<sup>3</sup>; FUJI, Nobuaki<sup>4</sup>

<sup>1</sup> 東京工業大学大学院理工学研究科地球惑星科学専攻, <sup>2</sup> ソウル大学, <sup>3</sup> 東京大学大学院理学系研究科地球惑星科学専攻, <sup>4</sup> パリ地球物理研究所

<sup>1</sup>Department of Earth and Planetary Sciences, Tokyo Institute of Technology, <sup>2</sup>School of Earth and Environmental Sciences, Seoul National University, <sup>3</sup>Department of Earth and Planetary Science, Graduate School of Science, University of Tokyo,

<sup>4</sup>Institut de Physique du Globe de Paris

We formulate the inverse problem of waveform inversion for localized 3-D seismic structure, computing partial derivatives of waveforms with respect to the elastic moduli at arbitrary points in space for anisotropic and anelastic media. In this study we minimize computational requirements by using the Born approximation with respect to a laterally homogeneous model, but this is not an inherent limitation of our approach. We solve the inverse problem using the conjugate gradient (CG) method, using Akaike's Information Criterion (AIC) to truncate the CG expansion. We apply our method to invert for three-dimensional shear wave structure in the lowermost mantle beneath Central America using a total of 2154 waveforms at periods from 12.5 to 200 s recorded at stations near the Pacific coast of North America for 29 deep and intermediate-depth events beneath South America. The resulting model shows lateral heterogeneity in the E-W direction which may be associated with a subducted cold slab surrounded by hotter materials with slower velocities. Various tests show that our model is robust.

キーワード: 最下部マントル, 波形インバージョン法, フアラロンプレート

Keywords: Lowermost mantle, Waveform inversion, Farallon plate

SIT03-19

会場:418

時間:4月29日 15:15-15:30

## 磁場でさぐる地球のコアと深部マントルのダイナミクス A magnetic probe into Earth's core and deep-mantle dynamics

櫻庭 中<sup>1\*</sup>

SAKURABA, Ataru<sup>1\*</sup>

<sup>1</sup> 東京大学大学院理学系研究科 地球惑星科学専攻

<sup>1</sup>Department of Earth and Planetary Science, University of Tokyo

It is widely recognized that Earth's core dynamics is an important research subject in understanding the past, present and future states of our planet, firstly because the metallic core is a vast domain accounting for one thirds of Earth's mass and plays a significant role in thermal history, and secondly because it dynamically generates the main geomagnetic field that has historically been observed for several hundred years and geologically recorded in rocks since more than a billion years ago. This review attempts to cover this subject with an attention to general questions: how geomagnetic-field data can be used to advance the deep-Earth science, and what theoretical progresses have been made and could be made. I will deal with some of the following particular topics: (1) various driving sources of convection, such as thermal and compositional buoyancy and inertial forcing (e.g., luni-solar precession); (2) a dynamo without a solid inner core; (3) a dynamo that operates in a part (e.g., an inner part) of the outer core; (4) sensitivity of the geomagnetic field structure (e.g., dipolarity), intensity, and time variations (e.g., reversal frequency) to the above mentioned various parameters.

SIT03-20

会場:418

時間:4月29日 15:30-15:45

## 高圧下における合金融体の音速と密度の関係 Relationship between sound velocity and density of liquid alloy under pressure

寺崎 英紀<sup>1\*</sup>; 西田 圭佑<sup>2</sup>; 浦川 啓<sup>3</sup>; 桑原 庄馬<sup>1</sup>; 田窪 勇作<sup>1</sup>; 下山 裕太<sup>1</sup>; 上杉 健太朗<sup>4</sup>; 竹内 晃久<sup>4</sup>; 鈴木 芳生<sup>4</sup>; 河野 義生<sup>5</sup>; 肥後 祐司<sup>4</sup>; 近藤 忠<sup>1</sup>

TERASAKI, Hidenori<sup>1\*</sup>; NISHIDA, Keisuke<sup>2</sup>; URAKAWA, Satoru<sup>3</sup>; KUWABARA, Souma<sup>1</sup>; TAKUBO, Yusaku<sup>1</sup>; SHIMOYAMA, Yuta<sup>1</sup>; UESUGI, Kentaro<sup>4</sup>; TAKEUCHI, Akihisa<sup>4</sup>; SUZUKI, Yoshio<sup>4</sup>; KONO, Yoshio<sup>5</sup>; HIGO, Yuji<sup>4</sup>; KONDO, Tadashi<sup>1</sup>

<sup>1</sup>Osaka Univ., <sup>2</sup>Univ. Tokyo, <sup>3</sup>Okayama Univ., <sup>4</sup>JASRI, <sup>5</sup>HPCAT

<sup>1</sup>Osaka Univ., <sup>2</sup>Univ. Tokyo, <sup>3</sup>Okayama Univ., <sup>4</sup>JASRI, <sup>5</sup>HPCAT

It is important to understand the relationship between sound velocity and density of liquid Fe-alloys under high pressure for obtaining a constraint of the composition of the molten outer core from observed seismic data. We have studied a relationship between sound velocity and density of liquid alloy based on simultaneous measurement of these properties under high pressure and high temperature. Sound velocity was measured using ultrasonic pulse-echo overlapping method and density was measured employing X-ray absorption method combined with X-ray tomography technique. The measured P-wave velocity and density of liquid Ni?S both increase with pressure. From these data, adiabatic bulk modulus ( $K_{S0}$ ) of the liquid sample can be well constrained to 29 GPa. It is noted that the measured P-wave velocity is found to increase linearly with increasing density. This result provides an important issue in terms of Birch's law for liquid material.

キーワード: 音速, 密度, 液体, 高圧

Keywords: Sound velocity, density, liquid, high pressure

SIT03-21

会場:418

時間:4月29日 15:45-16:00

## 対流駆動型ダイナモにおける回転磁気流体波 Magnetic-Coriolis waves in convection-driven dynamos: Implications for geomagnetic westward drift

堀 久美子<sup>1\*</sup> ; Jones Chris<sup>2</sup> ; Wicht Johannes<sup>3</sup> ; 清水 久芳<sup>1</sup>  
HORI, Kumiko<sup>1\*</sup> ; JONES, Chris<sup>2</sup> ; WICHT, Johannes<sup>3</sup> ; SHIMIZU, Hisayoshi<sup>1</sup>

<sup>1</sup> 東京大学地震研究所, <sup>2</sup> リーズ大学, <sup>3</sup> マックスプランク太陽系科学研究所

<sup>1</sup>Earthquake Research Institute, University of Tokyo, <sup>2</sup>University of Leeds, UK, <sup>3</sup>Max-Planck Institute for Solar System Research, Germany

A prominent feature of the geomagnetic secular variation is the westward motion of the non-dipole part of the field, which is significant in the Atlantic hemisphere with timescales of a few hundred years. Potential mechanisms to account for longitudinal geomagnetic drifts are advection due to large-scale zonal flows in the Earth's core as well as propagation of rotating magnetohydrodynamic (MHD) waves, particularly of slow Magnetic-Coriolis (MC) Rossby waves. More commonly the westward motion is thought to reflect zonal flow advection, an assumption that is used when inverting the secular variation signal for the flow at the top of the core. However, recent geodynamo simulations have successfully reproduced longitudinal magnetic drifts and some authors reported that the drift is at least partly a wave propagation.

To assess to what extent waves could play a role in geomagnetic drift, we explore nonlinear simulations of convection-driven MHD dynamos in rotating spherical shells. By performing a tempo-spatial spectral analysis of simulation data, we identify a slow MC-Rossby mode, that follows the dispersion curve predicted by a quasi-geostrophic linear theory. The result indicates that such waves can be excited in the planetary fluid core and that wave propagation may indeed play a role in the magnetic drifts. This gives a framework for further exploration of different wave types, which can provide valuable information about the physical properties in the deep interior fluid core.

SIT03-22

会場:418

時間:4月29日 16:15-16:30

## 地磁気移動性磁場の生成過程 A generating process of the geomagnetic drifting field

行武 賀<sup>1\*</sup>; 清水 久芳<sup>2</sup>

YUKUTAKE, Takeshi<sup>1\*</sup>; SHIMIZU, Hisayoshi<sup>2</sup>

<sup>1</sup>なし, <sup>2</sup>東京大学地震研究所

<sup>1</sup>None, <sup>2</sup>Earthquake Research Institute, University of Tokyo

地球磁場には数百年にわたりほぼ  $0.3^\circ$  /年の速度で西方移動する移動性磁場と同一場所に停滞して変化する停滞性磁場とが存在する。ここでは移動性磁場の生成過程についてひとつのモデルを提案する。

移動性磁場には二つの大きな特徴が存在する。地球磁場を球関数で展開すると、第一に移動性磁場は主として階数と次数が等しい sectorial 項から構成されている。第二に移動速度が階数次数によらずほぼ同じである。つまり分散性がない。

移動速度に分散性がない事から移動性磁場はあたかも核内の流体運動に凍結されたような状態で西向きの核内流体の流れの中で生成されていると考える。さらに核表層部の現象であるとすると、表層部で圧倒的に卓越する双極子磁場と流体運動の相互作用によって sectorial 項が卓越した移動性磁場が生成される。

単純化して核を3層のモデルで近似する。第1層は核マントル境界に接する薄層でマントルに固着している。第2層は第1層の内側で西向きに回転する層で内部で流体運動により問題の移動性磁場を発生する。第3層は静止流体とする。流体は核内全体で粘性ゼロ、第1層と第3層で電気伝導度有限、第2層で無限大として双極子磁場のもと回転系での計算を行った。さらにマントルの電気伝導度はゼロとする。

肝心なのは第2層内の流体運動でトロイダル流、ポロイダル流に分けて検討した。マントルが絶縁体であるから核マントル境界で電流の鉛直成分がゼロでなければならない。そのためにはトロイダル流は sectorial な流れ、ポロイダル流は meridional な流れに限られる。双極子磁場に作用して sectorial なトロイダル流は sectorial なポロイダル磁場を、meridional なポロイダル流は meridional なポロイダル磁場を誘導する。これらの磁場は第2層全体の西向きの回転に重畠して西向きに回転する西方移動磁場となる。さらにこれらのポロイダル磁場は第1層で減衰を受けた後マントルを通して地表で観測されることになるが、meridional な磁場の回転は回転していると認識されないため移動性磁場として観測されるのは sectorial な磁場だけとなる。

キーワード: 地磁気永年変化, 西方移動, 移動性磁場, 核表面流

Keywords: geomagnetic secular variation, westward drift, drifting field, core surface flow

SIT03-23

会場:418

時間:4月29日 16:30-16:45

## 地球中心核条件における純鉄の電気抵抗率 Electrical resistivity of hcp-Fe under Earth's core conditions

太田 健二<sup>1\*</sup>; 桑山 靖弘<sup>2</sup>; 廣瀬 敬<sup>3</sup>; 大石 泰生<sup>4</sup>  
OHTA, Kenji<sup>1\*</sup>; KUWAYAMA, Yasuhiro<sup>2</sup>; HIROSE, Kei<sup>3</sup>; OHISHI, Yasuo<sup>4</sup>

<sup>1</sup> 東京工業大学 大学院理工学研究科 地球惑星科学専攻, <sup>2</sup> 愛媛大学 地球深部ダイナミクス研究センター, <sup>3</sup> 東京工業大学 地球生命研究所, <sup>4</sup> 高輝度光科学研究センター

<sup>1</sup>Department of Earth and Planetary Sciences, Tokyo Institute of Technology, <sup>2</sup>Geodynamics Research Center, Ehime University, <sup>3</sup>Earth-Life Research Institute, Tokyo Institute of Technology, <sup>4</sup>Japan Synchrotron Radiation Institute

Iron is the primary component of the Earth's core. Convection of the conductive liquid outer core generates the geomagnetic field, and secular cooling of the core induces growth of the solid inner core and dynamics in the Earth's inside. Synchrotron x-ray diffraction study suggests that iron crystallizes in the hexagonal close-packed structure at the inner core conditions (Tateno et al., 2010). Thus, the electrical resistivity of hexagonal close-packed iron (hcp-Fe) is a key piece of information for estimating the transport properties of the core. We report high temperature electrical resistivity for hcp-Fe to 185 GPa measured in a laser-heated diamond anvil cell. We observed resistivity saturation in hcp-Fe under high pressure and high temperature conditions as predicted in a recent laboratory-based model for the conductivity of the Earth's core (Gomi et al., 2013). The saturation effect is significant in estimating electrical and thermal conductivity of the core, which strongly affect the dynamics and thermal evolution of the Earth.

### References

Gomi, H. et al. The high conductivity of iron and thermal evolution of the Earth's core. *Phys. Earth Planet. Inter.* 224, 88-103 (2013).

Tateno, S., Hirose, K., Ohishi, Y., & Tatsumi, Y. The structure of iron in Earth's inner core. *Science* 330, 359-361 (2010).

Keywords: Electrical resistivity, Earth's core, hcp iron

SIT03-24

会場:418

時間:4月29日 16:45-17:00

非等方熱拡散の空間依存性とその地球コアダイナミクスへの影響  
Spatial dependence of anisotropic thermal diffusivity and its influence on dynamics in the Earth's core

松島 政貴<sup>1\*</sup>

MATSUSHIMA, Masaki<sup>1\*</sup>

<sup>1</sup> 東京工業大学

<sup>1</sup>Tokyo Institute of Technology

Small-scale fluid motions in the Earth's core are likely to be highly anisotropic because of rapid rotation of the Earth and a strong magnetic field in the core. We have carried out direct numerical simulations of rotating magnetoconvection to investigate the effect of anisotropic diffusivity on dynamics in the Earth's core, as one of pilot studies. When a computational region is expressed in terms of a rectangular box with periodic boundaries in the three-directions, the prescribed anisotropic thermal diffusivities were found not to influence the character of rotating magnetoconvection, such as kinetic and magnetic energies averaged over the computational region. When a computational region is expressed in terms of a rectangular box with rigid boundary surfaces perpendicular to the gravitational direction, the prescribed anisotropic thermal diffusivities have a significant effect on the character of rotating magnetoconvection; that is, kinetic and magnetic energies can be increased even by a small anisotropy. The degree of increase depends on the direction of anisotropy and the direction of gravity corresponding to location of the computational region. These results suggest that anisotropic thermal diffusivity insignificantly influences dynamics in the bulk of the core, but that it should be effective near rigid boundary surfaces. Therefore, it is likely that anisotropic diffusivity has a more significant effect on MHD dynamos in rotating thin spherical shells. Such an implication can be examined through global numerical simulations of MHD dynamo models with anisotropic diffusivity being variable in the core.

Instead of such a study, we perform further direct numerical simulations of rotating magnetoconvection by prescribing anisotropic thermal diffusivities with spatial dependence; for example, in one case, anisotropic thermal diffusivities are presumed to be effective only near rigid boundary surfaces; in another case, anisotropic thermal diffusivities are presumed to be effective only far from rigid boundary surfaces. Hence this is another pilot study. Kinetic and magnetic energies in the former case seem to be larger than those in the latter case. The result is consistent with that obtained in our previous studies.

キーワード: 非等方性熱拡散, コアダイナミクス

Keywords: anisotropic thermal diffusivity, core dynamics

SIT03-25

会場:418

時間:4月29日 17:00-17:15

## Can a stably stratified layer interrupt the top-down dynamics of Earth's core? Can a stably stratified layer interrupt the top-down dynamics of Earth's core?

中川 貴司<sup>1\*</sup>

NAKAGAWA, Takashi<sup>1\*</sup>

<sup>1</sup>IFREE, JAMSTEC

<sup>1</sup>IFREE, JAMSTEC

Takashi Nakagawa

In some of previous studies of numerical dynamo simulation with a stably stratified region below the outer boundary, the long-wavelength feature of radial magnetic field can be only found on the outer boundary because a stratified layer can filter small-scale features of radial magnetic field generated in the convective region below the stratified boundary [Christensen, 2006; Nakagawa, 2011]. The existence of stably stratified region below the core-mantle boundary (CMB) is recently exposed from high pressure mineral physics [e.g. Pozzo et al., 2012] and seismological data analysis [e.g. Helffrich and Kaneshima, 2013].

Regarding the modeling on geomagnetic secular variation from numerical dynamo simulations, the heterogeneous thermal/chemical anomalies at the core-mantle boundary is important for understanding the time-scale of secular variation such as polarity reversals and excursions suggested from paleomagnetic observations [e.g. Olson et al., 2011; Olson et al., 2013] and current observational magnetic field [Aubert et al., 2013]. However, their investigation was not included in the effects of stably stratified region below the CMB in their dynamo simulations.

Here we introduce several examples of numerical dynamo simulations with both heterogeneous outer boundary prescribed by the CMB heat flux calculated from numerical mantle convection simulations and a stably stratified layer. Preliminary results are found that the large-scale and ?amplitude of thermal/chemical anomalies induced by the heterogeneous boundary condition, that is, thermal wind type flow, may be trapped at the imposed stratified boundary. This may imply that the geomagnetic secular variations related to the core-mantle coupling may be suggested that the core surface flow would be a key physics.

キーワード: Earth's core, heterogeneity, core-mantle boundary, stably stratified layer, thermal wind

Keywords: Earth's core, heterogeneity, core-mantle boundary, stably stratified layer, thermal wind

SIT03-26

会場:418

時間:4月29日 17:15-17:30

## 高温高圧下における 液体鉄ニッケル硫黄合金の弾性波速度測定 Sound velocity measurements of liquid Fe-Ni-S alloy at high pressure and temperature via inelastic X-ray scattering

今田 沙織<sup>1\*</sup>; 中島 陽一<sup>2</sup>; 駒林 鉄也<sup>3</sup>; 廣瀬 敏<sup>4</sup>; 筒井 智嗣<sup>5</sup>; 内山 裕士<sup>5</sup>; 石川 大介<sup>5</sup>; Alfred Baron<sup>2</sup>  
IMADA, Saori<sup>1\*</sup>; NAKAJIMA, Yoichi<sup>2</sup>; KOMABAYASHI, Tetsuya<sup>3</sup>; HIROSE, Kei<sup>4</sup>; TSUTSUI, Satoshi<sup>5</sup>; UCHIYAMA, Hiroshi<sup>5</sup>; ISHIKAWA, Daisuke<sup>5</sup>; ALFRED, Baron<sup>2</sup>

<sup>1</sup> 東京工業大学 地球惑星科学専攻, <sup>2</sup> 理化学研究所 理化学研究所, <sup>3</sup> エジンバラ大学, <sup>4</sup> 地球生命研究所, <sup>5</sup> 高輝度光科学研究センター

<sup>1</sup> Department of Earth and Planetary Sciences, Tokyo Institute of Technology, <sup>2</sup> Material Dynamics Laboratory, RIKEN SPring-8 Center, RIKEN, <sup>3</sup> School of Geosciences, The University of Edinburgh, <sup>4</sup> Earth-Life Science Institute, Tokyo Institute of Technology, <sup>5</sup> Japan Synchrotron Radiation Research Institute

液体 Fe-Ni 合金で構成されている外核には、水素 (H)、炭素 (C)、酸素 (O)、ケイ素 (Si)、硫黄 (S) 等の軽元素が含まれると考えられてきたが、含有される軽元素の種類、含有量は未だ明らかにされていない (e.g. Poirier 1994, Li and Fei 2003, McDonough 2003)。また、近年の観測により、火星や水星にも液体核が存在する可能性があることが報告された (Dehant, Science, Margot et al. 2007)

地震波速度は地球深部の観測量として最も信頼性が高い。液体 Fe 合金の弾性波速度のデータは、観測により得られた地震波速度と直接比較することが出来、外核組成決定に強い制約を与える。しかし実験的困難さから、5 GPa 以上の高圧下において液体 Fe 合金の弾性波速度を測定した例は無い (Nishida et al., 2012)。本研究では、地球、火星、水星外核に含まれる軽元素の候補の一つである硫黄に着目し、高圧発生装置ダイヤモンドアンビルセルと SPring-8 BL35XU、BL43XU 高分解能 X 線非弾性散乱装置を組み合わせ、高圧下における液体 Fe-Ni-S 合金の弾性波速度測定を、10-30 万気圧までの圧力範囲にて行った。出発試料として、 $(\text{Fe}_{0.83}\text{Ni}_{0.17})_3\text{S}$ 、 $(\text{Fe}_{0.64}\text{Ni}_{0.36})_3\text{S}$  の組成をもつ粉末混合物、もしくはマルチアンビル型高圧発生装置を用い、合成した試料を用いた。

今学会では、得られた弾性波速度データを用い、火星核、水星核について議論を行う。

キーワード: 弾性波速度, 非弾性散乱法, 外核, 液体鉄合金, 高温高圧実験

Keywords: sound velocity, inelastic X-ray scattering, planetary outer core, liquid iron alloy, High-PT experiment

SIT03-27

会場:418

時間:4月29日 17:30-17:45

## Experimental approach to the core-mantle boundary region of Mercury

### Experimental approach to the core-mantle boundary region of Mercury

岡本 美宝<sup>1</sup>; 浦川 啓<sup>1\*</sup>  
OKAMOTO, Miho<sup>1</sup>; URAKAWA, Satoru<sup>1\*</sup>

<sup>1</sup> 岡山大学 自然科学研究所

<sup>1</sup>Dept Earth Sci, Okayama Univ

MESSENGER mission revealed precise moment of inertia parameters of Mercury and its surface chemistry [1, 2]. These data allow to model the internal structure of Mercury, which has a large liquid core with ~2000 km radius and a solid outer shell with ~400 km thickness [3, 4]. As density of solid outer portion is apparently higher than that of the expected mantle silicate, the solid outer layer must include dense materials. Recent models [3, 4] showed that the Mercury's core contains sulfur and silicon as light elements due to high S fugacity and low oxygen fugacity of its interior. Those models presented a solid FeS layer at bottom of solid outer shell of Mercury as a dense layer, which separated from liquid outer core as a FeS-rich liquid due to liquid immiscibility of the Fe-S-Si ternary system. To investigate the FeS-rich layer at the top of Mercury's core, we performed the high-pressure experiments on the Fe-S-Si system using a KAWAI-type multi-anvil apparatus.

Pressure is fixed at 5 GPa corresponding to the CMB of Mercury and temperature is 1800 K, which is 200 K above the liquidus of Fe-S-Si system reported by Sanloup and Fei [5]. Fe-S-Si sample was kept for 30 min at this condition, and then it was quenched into room temperature. Oxygen fugacity of run charges was maintained around 3 log unit below IW buffer. Texture and chemistry of recovered samples were examined by electron microprobe.

We found two immiscible liquids in one run charge, which consist of Fe,Si-rich metallic liquid and FeS-rich sulfide liquid. Sulfur content of metallic liquid ranges 6 to 9 at%, which is higher by ~5 at% than those reported by Morard and Katsura [6]. Differences in texture of recovered samples and run duration between this study and Morard and Katsura [6] suggest that the latter experiments were in disequilibrium state. Our data shows the liquid immiscible region has a narrower extent than the previous estimation and the Mercury immiscible Fe-S-Si core must contain at least 6-9 at% sulfur. The quenched FeS-rich liquid phase consists mainly of crystalline FeS (~90 vol%) and Fe-Si alloy. In the case that FeS-rich liquid contacted with MgO sample container,  $(\text{Mg}_{0.8}\text{Fe}_{0.2})\text{S}$  crystalline phase coexisted with FeS-rich liquid. Mg-sulfide phase could be made by Fe-Mg exchange reaction between MgO and FeS-rich liquid. In the Mercury core, when FeS-rich liquid ascends to add the bottom of the CMB due to its buoyancy, it makes a stable low density layer. Mg-sulfide phase is produced under low oxygen fugacity and high sulfur fugacity at CMB, and then it incorporates into mantle. This is consistent with the results of X-ray fluorescence spectrometry on the Mercury's surface, which indicates the presence of Mg and Ca sulfides [2].

#### References

- [1] Margot JL et al., J. Geophys. Res., 117, E00L09, 2012.
- [2] Nittler LR et al., Science, 333, 1847, 2011.
- [3] Smith DE et al., Science, 336, 214, 2012.
- [4] Hauck II SA et al., J. Geophys. Res.: Planets, 118, 1204, 2013.
- [5] Sanloup C and Fei Y, Phys. Earth Planet. Inter., 147, 57, 2004.
- [6] Morard G and Katsura T, Geochim. Cosmochim. Acta, 74, 3659, 2010.

キーワード: 核, 核マントル境界, 水星

Keywords: core, CMB, Mercury