

Estimate of permafrost organic carbon balance in Alaskan boreal and tundra ecosystems using natural radiocarbon tracer

KONDO, Miyuki^{1*} ; UCHIDA, Masao¹ ; UTSUMI, Motoo² ; IWAHANA, Go³ ; YOSHIKAWA, Kenji³ ; IWATA, Hiroki⁴ ; HARAZONO, Yoshinobu³ ; NAKAI, Taro⁵ ; TANABE, Kiyoshi¹ ; SHIBATA, Yasuyuki¹

¹National Institute for Environmental Studies, ²University of Tsukuba, ³University of Alaska, Fairbanks, ⁴Kyoto University, ⁵Nagoya University

The high-latitude regions, where a serious warming is expected, currently store large amounts of soil organic carbon in active-layer soils and permafrost, accounting for nearly half of the global belowground organic carbon pool. Despite the importance of these regions in the present carbon cycle, the soil C fluxes and budget are still only poorly known. Here, we use radiocarbon as the tool for quantifying the C balance of the inputs and decomposition in tundra and boreal soil. We evaluated the C inputs (I) and decomposition rates (k , inverse of turnover time) and net C accumulation (CA), using ¹⁴C approaches.

Tundra and boreal soils show different patterns of depth distribution and C storage. Cumulative organic carbon stocks in boreal forest are 5.3 and 19.2 kgCm⁻², in surface organic layer (0-25 cm), and deep organic and mineral layers (25-70 cm), respectively. Large annual C input (0.25 kgCm⁻² yr⁻¹) and relatively slow decomposition (27 years) lead to rapid CA (0.05 kgCm⁻² yr⁻¹) in surface organic layer in boreal forest. Deep organic and mineral layers including near-surface permafrost show slower rate of input (0.03 kgCm⁻² yr⁻¹) and turnover (617 years) and CA about 20 times slower (0.003 kgCm⁻² yr⁻¹) than surface organic layer. Decomposition organic matter (Rh), which in accord with C losses from both surface and subsurface layers, was 0.23 kgCm⁻²yr⁻¹. This value agreed well with Rh (0.23 kgCm⁻² yr⁻¹) simulated by process-based models that simulate the biogeochemical and hydrologic cycle, where Rh averaged 45% of ecosystem respiration and 59% of soil respiration.

In contrast, large amount of SOC (36.4 kg m⁻²) have accumulated over millennia (turnover time: 4540 yrs) below the thin organic layer in tundra. The CA of mineral layer and permafrost is close to zero (0.003 kgCm⁻² yr⁻¹), and Rh is 0.008 kgCm⁻² yr⁻¹. Our radiocarbon data show that the most SOC in tundra soil was mode of stabilizing OC by permafrost and steady-state SOC stocks under current C balance.