

## Multiscale Interactions In The Genesis Of Tropical Cyclone Observed In PALAU2013

YOSHIOKA, Hiroaki<sup>1\*</sup> ; FUDEYASU, Hironori<sup>1</sup> ; KATSUMATA, Masaki<sup>2</sup> ; YOKOI, Satoru<sup>2</sup> ; TSUJINO, Satoki<sup>3</sup> ; MASUDA, Ayumi<sup>1</sup>

<sup>1</sup>Yokohama National University Graduate School, <sup>2</sup>Japan Agency for Marine-Earth Science and Technology, <sup>3</sup>Hydrospheric Atmospheric Research Center, Nagoya University

To understand the formation of a tropical cyclone (TC), has long been a captivating subject at the frontier of science and remains challenging because of the complex multi-scale interactions involved. During the genesis stage, sustained convective activities, which may stem from a variety of processes in a favorable environment, develop into a surface mesoscale or synoptic vortex. The mesoscale processes in the genesis stage have been the least understood aspect of the lifecycle of a TC. Although the climatological large-scale conditions favorable for TC genesis have been well known since Gray (1968, 1979), the interactions between the large-scale conditions and mesoscale processes have been poorly understood. The main purpose of this study is to conduct a detailed analysis on the multiscale interactions involved in the cyclogenesis based on observational data and numerical simulations.

The Japan Agency for Marine-Earth Science and Technology (JAMSTEC) conducted a field project named the Pacific Area Long-Term Atmospheric Observation for Understanding of Climate Change (PALAU2013) over the northwest Pacific Ocean. In this project, a sounding and radar network was deployed over the ocean during the early summer of 2013. During PALAU2013, the four initial disturbances growing tropical depression (TD) or tropical storm (TS) were observed. This study focused on the disturbances, growing TS (T1304) in association with the temporal changes in large-scale environment. We analyzed re-analysis data (JMA-GSM data), observational data during PALAU2013, and simulation results using WRF-ARW. The radiosondes were launched every 3h on the R/V Mirai and every 6h at Koror and Yap. The Doppler radar was installed on this ship, collected volume-scan every 10 and 7.5 min.

Initial disturbances which occurred at (3N, 175W) in 03UTC June 10, 2013, passed through the observation point R/V MIRAI MR13-03 at (12N, 135E), grew T1304 in 00UTC June 18. The disturbance was developing along the convergence region between the trade easterlies and monsoonal westerlies. Results of radiosonde show that potential temperature was higher in the middle and upper troposphere and CAPE increased as disturbances approached. Moreover, the zonal wind component of the lower troposphere changed to the strong easterly, corresponding to the meridional wind component of the troposphere also changed to the south from the north. The temporal variation of the radar-echo area during the convections showed the organization of convective clouds to form the intense cyclones.

Keywords: Tropical storm, Tropical depression, Tropics, Doppler radar, WRF