

大和川感潮域における栄養塩動態の季節変化 Seasonal variation in nutrient dynamics in the tidal zone of Yamato river

小野寺 真一^{1*}; 齋藤 光代²; 清水 裕太³; 丸山 豊¹; 宮岡 邦任⁴; 金 広哲¹; 有富 大樹¹
ONODERA, Shin-ichi^{1*}; SAITO, Mitsuyo²; SHIMIZU, Yuta³; MARUYAMA, Yutaka¹; MIYAOKA, Kunihide⁴; JIN,
Guangzhe¹; ARITOMI, Daiki¹

¹ 広島大学, ² 岡山大学, ³ 近畿中国四国農研センター, ⁴ 三重大学
¹Hiroshima University, ²Okayama University, ³NORO, ⁴Mie University

In coastal megacities, severe groundwater depression and water pollution occurred. These impacts affected to river environment change. Especially, the river mouth area has been deposited the polluted matters. These areas have characteristics of water level fluctuation which causes river water-groundwater interaction and the associated change in dynamics of nutrients. However, these effects on the nutrient transport in tidal reaches and nutrient load to the sea have not been fully evaluated in previous studies. Therefore, we aimed to clarify the nutrient dynamics with the river water-groundwater interaction in the tidal river of Osaka metropolitan city. We conducted the field survey from the river mouth to the 7km upstream area of Yamato River, which has a length of 68km and a watershed area of 1070 km². In addition, model simulations were also conducted. Spatial variations in radon (²²²Rn) concentrations and the difference of hydraulic potential between river waters and the pore waters suggest that the groundwater discharges to the river channel in the upstream area. In contrast, river water seeped into the groundwater in the river mouth area. It may be caused by the lowering of groundwater level associated with the excess abstraction of groundwater in the urban area. The spatial and temporal variations in nutrient concentrations indicate that nitrate-nitrogen (NO₃-N) concentrations changed temporally and it negative correlated with dissolved organic nitrogen (DON) concentrations. Inorganic phosphorous (PO₄-P) concentrations showed the increasing trend with the increase of the river water level. Based on the mass balance, nutrient reproduction from the river bed was suggested in tidal reach during a summer, especially phosphorus was large.

キーワード: 季節変化, 栄養塩動態, 干潮河川, 汚濁, リン

Keywords: seasonal variation, nutrient dynamics, tidal river, pollution, phosphorus