

## 海洋酸性化が温帯性サンゴの成長に与える影響について Ocean acidification influences on coral growth of temperate species

Kim So<sup>1\*</sup>; 鈴木 淳<sup>2</sup>; 林 正裕<sup>3</sup>; 山本 雄三<sup>3</sup>; 堀田 公明<sup>3</sup>; 磯野 良介<sup>3</sup>; 渡邊 裕介<sup>3</sup>; 山野 博哉<sup>4</sup>; 野村 恵一<sup>5</sup>; 西田 梢<sup>2</sup>; 井上 麻夕里<sup>6</sup>; 張 勁<sup>1</sup>; 野尻 幸宏<sup>4</sup>  
KIM, So<sup>1\*</sup>; SUZUKI, Atsushi<sup>2</sup>; HAYASHI, Masahiro<sup>3</sup>; YAMAMOTO, Yuzo<sup>3</sup>; HOTTA, Kimiaki<sup>3</sup>; ISONO, Ryosuke<sup>3</sup>; WATANABE, Yusuke<sup>3</sup>; YAMANO, Hiroya<sup>4</sup>; NOMURA, Keiichi<sup>5</sup>; NISHIDA, Kozue<sup>2</sup>; INOUE, Mayuri<sup>6</sup>; ZHANG, Jing<sup>1</sup>; NOJIRI, Yukihiro<sup>4</sup>

<sup>1</sup> 富山大学, <sup>2</sup> 産業技術総合研究所, <sup>3</sup> 公益財団法人海洋生物環境研究所, <sup>4</sup> 国立環境研究所, <sup>5</sup> 串本海中公園センター, <sup>6</sup> 東京大学大気海洋研究所

<sup>1</sup>Toyama University, <sup>2</sup>Geological Survey of Japan, AIST, <sup>3</sup>Marine Ecology Research Institute (MERI), <sup>4</sup>National Institute for Environmental Studies, <sup>5</sup>Kushioto Marine Park, <sup>6</sup>Atmosphere and Ocean Research Institute, The University of Tokyo

Carbon dioxide concentration in the atmosphere has steadily increased since the industrial revolution due to burning of fossil fuel and will cause the global warming and ocean acidification. It will raise the ocean temperature around Japan and reduce the seawater pH and then it may bring serious threat to corals dwelling around Honsyu Island, Japan. Last year, our research group did temperature-controlled culture experiments of temperate coral species from the Pacific side of Honsyu Island of Japan under the present level of the partial pressure of CO<sub>2</sub> (pCO<sub>2</sub>). But, synergetic effect of the global warming and ocean acidification on these corals has not been tested yet in detail. In this study, we focus on the how the different pCO<sub>2</sub> levels (past, present, and future) can influence skeletal growth of temperate *Acropora* coral species under the different temperature setting using a precise control system. This system was used to generate six different pCO<sub>2</sub> levels: (i) pre-industrial, ~300 μatm, (ii) present-day pCO<sub>2</sub>, ~400 μatm, and at four near-future conditions, (iii) ~550 μatm, (iv) ~750 μatm, (v) ~1000 μatm and (vi) ~1200 μatm at three temperature conditions (17, 25, and 27 deg C). Our early results suggested a negative influence of higher pCO<sub>2</sub> levels on skeletal growth of temperate *Acropora* corals, but not so sensitive compared to tropical and subtropical *Acropora* corals.

キーワード: 海洋酸性化, 温帯サンゴ, 石灰化, 地球温暖化

Keywords: Ocean acidification, temperate coral, calcification, global warming