

二枚貝の炭素・窒素安定同位体比による養殖域からの懸濁物拡散の評価 Propagation of suspended matter from aquacultures as traced by stable C and N isotope ratios of bivalves

森本直子^{1*}; 梅澤有²; 田中義幸³; レジーノ ジュネビエーブ⁴; 渡邊敦⁵; マックグローン マリア・ローデス⁴; 宮島利宏¹

MORIMOTO, Naoko^{1*}; UMEZAWA, Yu²; TANAKA, Yoshiyuki³; REGINO, Genevieve L.⁴; WATANABE, Atsushi⁵; MCGLONE, Maria lourdes s.d.⁴; MIYAJIMA, Toshihiro¹

¹ 東京大学 大気海洋研究所, ² 長崎大学 水産学部, ³ 海洋研究開発機構 むつ研究所, ⁴ フィリピン大学 海洋科学研究所, ⁵ 東京工業大学大学院 情報理工学研究科

¹ Atmosphere and Ocean Research Institute, The University of Tokyo, ² Faculty of Fisheries, Nagasaki University, ³ Mutsu Institute for Oceanography, Japan Agency for Marine-Earth Science and Technology, ⁴ Marine Science Institute, University of the Philippines, ⁵ Graduate School of Information Science and Engineering, Tokyo Institute of Technology

Recently there is growing concern about the impact of densely-deployed aquacultures on coastal marine ecosystems in the Philippines. As suspension-feeding bivalves are expected to reflect local food sources, their effectiveness as an environmental indicator were examined by analyzing stable carbon and nitrogen isotope ratios of bivalves living in aquaculture and neighboring seagrass areas. As a whole, the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ of bivalves collected in the seagrass areas ranged from -13.1 to -11.0 and from +4.0 to +6.6, respectively, but in seagrass area where water mass from aquaculture area passed through typically lower values (-18.9 ~ -16.1 and +2.7 ~ +5.2, respectively) were observed, and they were the lowest in the aquaculture area (-24.4 ~ -19.8 and +3.4 ~ +4.3, respectively). It suggests that bivalves mainly fed on sinking particles, and presumably also seagrass-derived particles in seagrass areas. Higher C/N ratio was observed at sites where impact of aquaculture was larger. Although the interspecies differences and food selectivity etc. may affect the variability of the bivalve $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ to some extent, these results demonstrated that stable isotope ratios of bivalves could be used as an effective indicator to evaluate propagation areas and actual effects of suspended matter resulting from anthropogenic source on ecosystems.

キーワード: 懸濁物食二枚貝, 海草, 養殖, 炭素・窒素安定同位体比

Keywords: suspension-feeding bivalve, seagrass, aquaculture, stable carbon and nitrogen isotope ratios