

## Fluorometric analysis of photosymbiosis: Toward quantitative validation of ecological proxy of planktic foraminifers

TAKAGI, Haruka<sup>1\*</sup> ; KIMOTO, Katsunori<sup>2</sup> ; FUJIKI, Tetsuichi<sup>2</sup> ; KURASAWA, Atsushi<sup>2</sup> ; HIRANO, Hiromichi<sup>3</sup>

<sup>1</sup>CSE Grad. School, Waseda University, <sup>2</sup>Japan Agency for Marine-Earth Science and Technology, <sup>3</sup>Dep. Earth Sci., Sch. Edu., Waseda University

Endosymbiosis of planktic foraminifers with photosynthetic algae (photosymbiosis) is established especially among species which dominate in warm, low-nutrient surface water. Here, photosymbiosis probably plays an important role for host foraminifers, and can be considered as an adaptive ecology to live in such oligotrophic oceans. Therefore, back in geologic time, photosymbiosis could have been involved with species adaptive radiation as well. In such viewpoint, stable isotopic change of foraminiferal test through ontogeny, attributed to change of symbiont photosynthetic effect, has been used as an indicator to detect fossil photosymbiosis. However, how host-symbiont association change through ontogeny, if any, is practically unknown and has never been quantified. Here, we offer new insights for photosymbiosis based on photosynthetic characteristics of symbionts, obtained by in vivo fluorometric analysis (Fast Repetition Rate Fluorometry, FRRF).

We cultured two symbiont-bearing species, *Globigerinoides sacculifer* and *Globigerinella siphonifera*, and conducted FRRF measurement on individual host-algal consortium during the culture period. FRRF can identify photosymbiosis of individual foraminifer instantly in a non-destructive manner, and gives us various photosynthetic characteristics of symbionts, i.e., maximum fluorescence yield ( $F_m$ , index of chlorophyll content), photochemical efficiency ( $F_v/F_m$ , index of potential photosynthetic activity), and effective absorption cross-section of photosystem II ( $\sigma_{PSII}$ , capability of the absorbed energy to promote a photochemical reaction).

Sequential FRRF analyses on single individuals revealed that  $F_m$  increases with growth, and then decrease drastically at the end of their life, which means that the algal biomass per individual foraminifer increases through ontogeny, but the symbionts are rapidly digested at the end.  $F_v/F_m$  and  $\sigma_{PSII}$  values were constant through ontogeny, though  $F_v/F_m$  drops in correspondence with the decrease of  $F_m$ . Compared between the two species, average values of both  $F_v/F_m$  and  $\sigma_{PSII}$  showed statistically significant differences.  $F_v/F_m$  was significantly higher in *Gs. sacculifer*, which means that symbionts are more actively photosynthesizing in *Gs. sacculifer*. Because  $F_v/F_m$  is mainly depends on nutrient availability, it is a direct evidence of nutrient (metabolite) flow from host to symbionts. On the other hand,  $\sigma_{PSII}$  was higher in *Gn. siphonifera*, indicating that this species can utilize low light energy more efficiently, i.e., more " low-light-adapted " than *Gs. sacculifer*. Actually, it is consistent with inferred habitat preference of *Gn. siphonifera*, which is relatively deeper than *Gs. sacculifer*.

These FRRF results provide us information of foraminiferal photosymbiosis both quantitatively and qualitatively. When the information is combined with test geochemistry mentioned above, it will presumably enable us to quantify the photosynthetic activity from foraminiferal tests. Then, it can be applied to fossil specimens as a validated ecological proxy of photosymbiosis.

Keywords: planktic foraminifers, photosymbiosis, Fast Repetition Rate Fluorometry