

Snowball Earth and GCM simulation Snowball Earth and GCM simulation

丸山 茂徳^{1*}; 鈴木 遼平²; 大淵 濟³

MARUYAMA, Shigenori^{1*}; SUZUKI, Ryohei²; OHFUCHI, Wataru³

¹ 東京工業大学地球生命研究所, ² 東京工業大学, ³ 独立行政法人海洋研究開発機構

¹ELSI Tokyo Institute of Technology, ²Tokyo Institute of Technology, ³JAMSTEC

Numerical simulation of snowball Earth, using out-of-date supercomputer program has been performed recently in USA, France and Germany. It seems to be difficult to reconstruct Snowball state by their simulation, while freezing more than 55% of ocean. If continents are gathered along the equatorial region such as Rodinia in the case of Sturtian and Marinoan Snowball Earth in Neoproterozoic, total surface irradiance (TSI) seems plausible to be 95% of present day and CO₂ level as same as today. However, if the atmospheric CO₂ is 2-6 times more than today, Snowball state cannot appear (Voigt et al., 2011). More realistic CO₂ concentration of Neoproterozoic Earth was 20-50 times more than today. In addition, the temperature fluctuation of Snowball Earth period, from Sturtian to Marinoan, was -40 °C to +40 °C and vice versa within a short period <10 m.y. which seem to be impossible because input and output of CO₂ by plate tectonics usually takes time more than several hundreds of millions years.

GCM simulation exaggerates positive feedback of CO₂ too much. It is time to remodel GCM, considering the amount of clouds and its effect.