

南アフリカ、Onverwacht 層群における初期太古代の硫黄非質量依存同位体分別 S-MIF geochemistry of the Early Archean in the Onverwacht Suite, South Africa

三島 郁^{1*}; 上野 雄一郎¹; de Wit Maarten²; Furnes Harald³; 齋藤 拓也¹
MISHIMA, Kaoru^{1*}; UENO, Yuichiro¹; DE WIT, Maarten²; FURNES, Harald³; SAITOU, Takuya¹

¹ 東京工業大学大学院地球惑星科学専攻, ² ネルソンマンデラメトロポリタン大学, ³ ベルゲン大学

¹Department of Earth and Planetary Sciences, Tokyo Institute of Technology, ²EON-Africa Earth Observatory Network, and Faculty of Science, Nelson Mandela Metropolitan University, ³Department of Earth Science and Centre for Geobiology, University of Bergen

The recent study of sulfur mass independent fractionation (S-MIF) in the Archean sedimentary rocks represented that multiple sulfur isotope ratios ($^{32}\text{S}/^{33}\text{S}/^{34}\text{S}/^{36}\text{S}$) could be useful new tracer for Archean sulfur cycles. Farquhar et al. (2000) first discovered that Archean sedimentary rocks before 2.4 Ga have $\Delta^{33}\text{S}$ anomaly, whereas no such anomaly was found in younger samples. This contrast implies the rise of atmospheric oxygen content that fundamentally changed atmospheric sulfur cycle. The hypothesis are based on the studies from Western Australia and South Africa (Kaufman et al., 2007; Ono et al., 2009; Zerckle et al., 2013). High-resolution stratigraphic studies provide a detailed view into the late Archean marine sulfur cycle, which can help our understanding of both atmospheric and biological processes. In the early Archean, S-MIF data are almost from hydrothermal sulfate and sulfide. For comparing early and late Archean data precisely, it is necessary to investigate stratigraphical and petrological distributions and variations of the multiple sulfur isotopes. We have studied Early Archean sedimentary sulfides which are well preserved in the Barberton Greenstone Belt, South Africa. Sulfur isotope analysis of extracted sulfide of sedimentary rocks from Barberton Greenstone Belt, show a clear MIF ($>1\text{‰}$) and $\delta^{34}\text{S}-\Delta^{33}\text{S}$, $\Delta^{33}\text{S}-\Delta^{36}\text{S}$ correlation. The Noisy Complex which consists of fluvial sediments and diamictite show negative $\delta^{34}\text{S}-\Delta^{33}\text{S}$ correlation, and $\Delta^{36}\text{S}/\Delta^{33}\text{S}$ slope of -0.72. On the other hand, the Kromberg Formation which consists of deep marine sediments show positive $\delta^{34}\text{S}-\Delta^{33}\text{S}$, and scattered $\Delta^{36}\text{S}/\Delta^{33}\text{S}$ slope. $\delta^{34}\text{S}-\Delta^{33}\text{S}$, $\Delta^{33}\text{S}-\Delta^{36}\text{S}$ relation from each stratigraphic level shows somewhat different trend, possibly reflecting local environment and/or bacterial sulfate reduction activity.

キーワード: 南アフリカ, 硫黄, 非質量依存同位体分別
Keywords: South Africa, Sulfur, MIF