

重元素同位体比から探る鉍化熱水の起源 Sr-Nd-Pb isotopic compositions of hot spring water in the Toyoha Mine, Hokkaido Japan: Implications for the origin of hy

中村 仁美^{1*}; 岩森 光²; 荒木 修平³; 稗田 裕樹³; 藤永 公一郎³; 加藤 泰浩³; 中井 俊一⁴; 木村 純一²; 常 青²
NAKAMURA, Hitomi^{1*}; IWAMORI, Hikaru²; ARAKI, Shuuhei³; HIEDA, Yuki³; FUJINAGA, Koichiro³; KATO, Yasuhiro³
; NAKAI, Shun'ichi⁴; KIMURA, Jun-ichi²; CHANG, Qing²

¹ 東京工業大学, ² 海洋研究開発機構, ³ 東京大学, ⁴ 東京大学地震研究所

¹Titech, ²JAMSTEC, ³Univ. of Tokyo, ⁴ERI, Univ. of Tokyo

Chemistry and dynamics of slab-derived fluids in subduction zones have been rigorously studied by high pressure experiments, geochemical and hydrological modellings, and geophysical observations [1-5]. Surface manifestation of deep slab-derived fluids are now suggested by geochemistry, such as slab fluid-like chemical affinities found in volcanic rocks [6,7] and in hot spring waters [8]. In this study, we aim to examine the presence (or absence) of slab derived fluid signatures in hot spring water related with the Toyoha Mine ore deposits in Hokkaido, one of the largest hydrothermal vein-type deposits in Japan. We applied Sr-Nd-Pb isotope analyses of the hot spring water and compared the results to those from the volcanic rocks and the ore minerals from the Toyoha Mine.

For this purpose, we have examined a ferric co-precipitation pre-concentration method for the hot spring water from the Toyoha Mine. This was necessary because the abundances of Nd and Pb were very low, less than several ppb for Nd, in particular. The method has previously been applied to brines with high chlorine concentration at Arima hot spring [9], and the method worked well with the Toyoha hot spring water. The concentrated sample has been analyzed by Q-ICP-MS and MS-ICP-MS for both element abundances and Sr-Nd-Pb isotopic compositions. We examined origin of the hot spring water by using Sr-Nd-Pb isotope systematics in comparison with the data from the ore deposit, volcanic rocks related with the ore deposition, and the basement rocks of the Toyoha Mine. A recent study has shown that Sr-Nd-Pb isotopic ratios of sulfide ores in the Toyoha Mine exhibit a high contribution of slab-derived fluid from the Pacific Plate slab [10]. Our preliminary results on the hot spring water suggest that the water may also preserve the slab-fluid signatures and/or may also be affected by the chemical components in the basement rocks.

[1] Schmidt and Poli, 1998, EPSL [2] Hacker et al., 2003, JGR [3] Iwamori, 1998, EPSL [4] Arcay et al., 2005, PEPI [5] Cagnioncle et al., 2007, JGR [6] Pearce et al., 2005, G3 [7] Nakamura et al., 2008, NGeo [8] Kusuda et al., in revision [9] Nakamura et al., submitted [10] Hieda, 2013, Master Thesis, Univ. of Tokyo

キーワード: 温泉水, 同位体, 豊羽, 鉍床, 熱水
Keywords: hot spring, isotope, Toyoha, mine, ore