

## Left-right reversal in unicellular eukaryotes, planktonic foraminifera

UJIE, Yurika<sup>1\*</sup> ; ASAMI, Takahiro<sup>1</sup>

<sup>1</sup>Department of Biology, Shinshu University

Aquatic unicellular organisms are little motile and passively disperse in general. Holoplankton, which spend their entire life-cycle floating in the water column, are likely carried by water flow and exposed to diverse conditions of environment. Their morphology may vary over wide distribution ranges by phenotypic plasticity or allelic variation. Among these organisms, planktonic foraminifera are an excellent system to examine diversity and evolution in cellular responses to the environment because of two reasons: (1) occurrence in every ocean and (2) visible asymmetry in coiled shell. Both left- and right-coiled forms are often found within single morphospecies. Their coiling direction has traditionally been thought to change phenotypically depending on environmental factors, especially water temperature, based on coil-morph distributions but without statistical evidence. Molecular phylogenetic studies have revealed that morphospecies often contain multiple cryptic species. The arguments on the role of temperature for coil reversal most probably confused cryptic species into single taxa. In the present study, we examined the dependence of morph frequency on temperature by focusing on populations that are dimorphic for coiling direction and occur across wide ranges of temperature. *Globorotalia truncatulinoides* includes five genetically isolated species, and each of them is dimorphic for coiling direction. The statistically meaningful regression analysis was possible in three species that are distributed in global ranges. The results showed that morph frequency does not depend on water temperature in warm or cold seasons or on the annual mean temperature. Moreover, the geographic patterns of frequency variation among water masses in these species suggest that gene flow affects morph frequency. The majority exhibits the same coiling direction among populations that inhabit water masses connected by ocean circulation system. In contrast, morph frequency greatly varies between unconnected water masses regardless of climatic conditions. The present results, therefore, reject temperature-dependence of coiling direction and suggest the presence of genetic basis for coiling direction in planktonic foraminifera. Our study provides a base to explore the evolution on left-right asymmetry in unicellular eukaryotes.

Keywords: left-right asymmetry, coiling direction, cryptic species, unicellular eukaryote