

## The 24 September 2013 tsunami in the Makran region, northwestern Indian Ocean

HEIDARZADEH, Mohammad<sup>1</sup> ; SATAKE, Kenji<sup>1\*</sup>

<sup>1</sup>Earthquake Research Institute (ERI), The University of Tokyo

Tsunami waves were observed in the northwestern Indian Ocean following the Mw 7.7 Pakistan inland earthquake on 24 September 2013. We analyze eleven tide gauge records as well as one DART record of this tsunami and perform numerical modeling of tsunami. The tsunami registered a maximum wave height of 109 cm in Qurayat tide gauge station (Oman). Spectral analysis showed that the most governing period of the tsunami waves was around 12 min though wavelet analysis showed that parts of the tsunami energy were partitioned into other period bands of 7 and 16 min. Distribution of aftershocks in the region showed that all of them were located inland indicating that the tsunami was generated by submarine geological phenomena triggered by the earthquake. Tsunami backward ray tracing showed that the tsunami source was possibly located at offshore Jiwani (Pakistan) and the tsunami was most likely generated immediately after the main shock. Tsunami modeling assuming a pile-up structure at the location of the new island was not successful in reproducing the observed sea level records. A landslide source with a length of about 15-20 km, a thickness of 100-150 m located at 61.72°E and 24.60°N seems capable of fairly reproducing the observed sea level records. This event was the second tsunami recorded in the Makran region since 1945, and may be evidence for hazards from landslide-generated waves following seismic activities.

Keywords: Northwestern Indian Ocean, Tsunami, Makran subduction zone, Landslide, Spectral analysis, Numerical modeling