

真核生物の細胞膜に存在するステロールおよびその代替物質に関する謎 Enigmas concerning sterols and their surrogates in eukaryotic cell membranes

瀧下 清貴^{1*}; 矢吹 彬憲¹; 力石 嘉人¹; 高木 善弘¹; 吉田 尊雄¹; 大河内 直彦¹

TAKISHITA, Kiyotaka^{1*}; YABUKI, Akinori¹; CHIKARAISHI, Yoshito¹; TAKAKI, Yoshihiro¹; YOSHIDA, Takao¹; OHKOUCHI, Naohiko¹

¹ 海洋研究開発機構

¹Japan Agency for Marine-Earth Science and Technology

A large fraction of eukaryotes and bacteria respectively possess sterols and hopanoids, which function as potent stabilizers of cell membranes. Sterols are also associated with fluidity and permeability of eukaryotic cell membranes, and are key to fundamental eukaryotic-specific cellular processes such as phagocytosis. Several steps of *de novo* sterol biosynthesis require molecular oxygen. For example, the epoxidation of squalene is the first oxygen-dependent step in the sterol pathway; the epoxidized squalene is then cyclized to either lanosterol or cycloartenol by the enzyme oxidosqualene cyclase. In contrast, prokaryotic hopanoid biosynthesis does not require molecular oxygen as a substrate, and the squalene is directly cyclized by the enzyme squalene-hopene cyclase.

Until now, it was unclear how bacterivorous unicellular eukaryotes that are abundant in anoxic or low oxygen environments could carry out phagocytosis. These eukaryotes cannot obtain sterols from food bacteria as the latter generally lack them and sterols cannot be synthesized *de novo* in the absence of molecular oxygen. We have previously provided evidence that the molecule tetrahymanol is synthesized by some anaerobic/microaerophilic eukaryotes and possibly functions as an analogue of sterols in these organisms. Nevertheless, neither sterol, nor tetrahymanol, nor their related molecule has been found in the other anaerobic/microaerophilic eukaryotes, and so it is still enigmatic how these organisms maintain their fluid and permeable membrane system specific to eukaryotes.

One more area of confusion is regarding sterols in bivalves with chemosynthetic bacteria inhabiting areas of deep-sea hydrothermal vents and methane seeps, such as *Calyptogena* spp. and *Bathymodiolus* spp. In general, bivalves cannot synthesize sterols *de novo* and it is necessary for them to obtain these molecules from small eukaryotic prey. On the other hand, *Calyptogena* spp. and *Bathymodiolus* spp. mainly or exclusively acquire nutrients produced by their bacterial symbionts, rather than from eukaryotes rich in sterols. Nevertheless, these "chemosynthetic bivalves" contain sterols. More curiously, *Calyptogena* spp. have intermediate metabolites of phytosterols (24-methylenecycloartanol, cycloeucalenol, and obutusifoliol), while *Bathymodiolus* spp. have high amounts of cholesterol typical of animals. Little attention has been given to how chemosynthetic bivalves produce or acquire these kinds of sterols.

In my talk, I will discuss potentially controversial topics regarding sterols and their surrogates in eukaryotic cell membranes, which do not appear in biochemical and geochemical textbooks.

キーワード: 真核生物, ステロール, テトラヒマノール, 細胞膜

Keywords: eukaryotes, sterols, tetrahymanol, cell membrane