

## 地球内起源ニュートリノによる新しい地球観測 Geo-neutrinos for advanced earth studies

田中 宏幸<sup>1\*</sup>  
TANAKA, Hiroyuki<sup>1\*</sup>

<sup>1</sup> 東京大学地震研究所

<sup>1</sup>Earthquake Research Institute, The University of Tokyo

Neutrinos generated in Earth (geo-neutrinos) gives us information about the distribution of Uranium (U), thorium (Th), and potassium (K) inside Earth. Beta-decays of radionuclides U/Th/K inside Earth produce low energy anti-electron neutrinos (U and Th produces  $7.41 \times 10^7$  neutrinos  $\text{kg}^{-1}\text{s}^{-1}$  and  $1.62 \times 10^7$  neutrinos  $\text{kg}^{-1}\text{s}^{-1}$  respectively (without considering neutrino oscillation)) that traverse through Earth without being disturbed due to their extremely small interaction cross section with matter. Recent geo-neutrino observations have produced results that have a potential to support and clarify the current concerns of earth science: estimating the amount of contribution to the surface heat flux; constraining existing Earth's compositional estimates; and clarifying the origin of low shear velocity regions found at the core mantle boundary (CMB). Today, there are two detectors capable of measuring geoneutrinos: KamLAND, in Japan, and Borexino, in Italy. The KamLAND research team has found  $116 \pm 28$ , 27 geoneutrino candidate events (generated through the decay processes of  $^{238}\text{U}$  and  $^{232}\text{Th}$ ) during 2,991 days of geoneutrino observation (Gando et al. 2013). The contribution from geonuclear reactions to the heat flow, estimated from examination of the geoneutrino flux, reached  $11.2 \pm 7.9$ , 5.1 TW. Although the volume of the Borexino detector (280 t) is much smaller than that of KamLAND (1,000 t), the background from reactor neutrinos is much lower than that for KamLAND because there are no nuclear power plants in Italy. Borexino detected  $14.3 \pm 4.4$  geoneutrino candidates over 1353 days of observation. Both measurement results are consistent each other, and also reject the fully radiogenic model, which assumes that the total Earth's surface heat flux is completely originated from radiogenic heat from U, Th and K. The upper limit on the fully radiogenic heat flux hypothesis (Herndon 1996) was set to be 4.5 TW at 95% confidence level (Bellini et al. 2013).

Gando A, Gando Y, Hanakago H, Ikeda H, Inoue K, et al. 2013. Reactor on-off antineutrino measurement with KamLAND. Phys. Rev. D 88:033001.

Herndon JM. 1996. Substructure of the inner core of the Earth. Proc. Natl. Acad. Sci. USA 93:646-48.

Bellini, G., Ianni, A., Ludhova, L., Mantovani, F., McDonough, W.F. 2013. Geo-neutrinos, Prog.Part.Nucl.Phys. 73:1-34

キーワード: ニュートリノ, ウラニウム, トリウム, マントル  
Keywords: Neutrino, Uranium, Thorium, Mantle