

PEM06-P10

会場:3階ポスター会場

時間:4月30日 18:15-19:30

Na ライダー/EISCAT レーダー連携観測によって検出したオーロラ起因の Na 層変動 Aurora-induced sodium layer variation detected by coordinated observation with sodium lidar and EISCAT radar

津田 卓雄^{1*}; 野澤 悟徳²; 川原 琢也³; 川端 哲也²; 斎藤 徳人⁴; 和田 智之⁴; 小川 泰信¹; 大山 伸一郎²; Hall Chris⁵; 堤 雅基¹; 江尻 省¹; 鈴木 臣²; 高橋 透²; 中村 卓司¹

TSUDA, Takuo^{1*}; NOZAWA, Satonori²; KAWAHARA, Takuya³; KAWABATA, Tetsuya²; SAITO, Norihito⁴; WADA, Satoshi⁴; OGAWA, Yasunobu¹; OYAMA, Shin-ichiro²; HALL, Chris⁵; TSUTSUMI, Masaki¹; EJIRI, Mitsumu K.¹; SUZUKI, Shin²; TAKAHASHI, Toru²; NAKAMURA, Takuji¹

¹ 国立極地研究所, ² 名古屋大学太陽地球環境研究所, ³ 信州大学工学部, ⁴ 理化学研究所, ⁵ トロムソ大学

¹National Institute of Polar Research, ²Solar-Terrestrial Environment Laboratory, Nagoya University, ³Faculty of Engineering, Shinshu University, ⁴RIKEN, ⁵University of Tromsøe

Sodium atom layer is generally distributed at 80-100 km. One of mysterious subjects on high-latitude sodium layers is relationship between auroral particle precipitation and sodium atom layer variation. A previous study suggested a sodium column density decrease during a geomagnetic active period due to that the particle precipitation accompanied by electron density enhancement could induce ionization of sodium atom through their ion-molecule chemistry. Another study pointed a possibility of sodium density increase. For this reason, it is suggested that auroral precipitating particle bombardment on meteoric smoke particles can sputter sodium atoms from the smoke particles. On the other hand, ionospheric electric field, which may become more significant near auroral precipitating regions, could induce ion motions (i.e. can generate sodium ion convergence and/or divergence), and then also could affect generation and/or loss processes of sodium atoms through their ion-molecule chemistry. Thus, for the examination of the causality, it is vitally important to distinguish the effects of auroral particle precipitation and ionospheric electric field. Using a sodium lidar (which was installed in early 2010) and European incoherent scatter (EISCAT) radar at Tromsøe, Norway (69.6N, 19.2E), we have investigated, for the first time, that the actual effect of the particle precipitation to the sodium density variations without electric filed injection. In the nighttime observation on 24-25 January 2012, we detected a significant decrease of sodium atom density coincided with electron density enhancements (implying strong particle precipitations) and low ion temperatures (implying no electric field injections). These results strongly suggested that auroral particle precipitations induced sodium atom density decrease. Furthermore we discuss observed time response in the sodium density decrease.

キーワード: Na ライダー, EISCAT レーダー, Na 層, オーロラ降下粒子, 電離圏電場

Keywords: Na lidar, EISCAT radar, Na layer, Auroral particle precipitation, Ionospheric electric field