

イオン誘起微粒子核生成 II: 水クラスターイオンの自由エネルギー Ion-induce nucleation experiment II: free energy of the water-cluster ion

日高 宏^{1*}; 中井 陽一²; 小島 隆夫³; 渡部 直樹¹
HIDAKA, Hiroshi^{1*}; NAKAI, Yoichi²; KOJIMA M., Takao³; WATANABE, Naoki¹

¹ 北海道大学低温科学研究所, ² 理化学研究所仁科センター, ³ 理化学研究所原子物理研究室
¹Inst. Low Temp. Sci., Hokkaido Univ., ²RIKEN Nishina Center, ³RIKEN Atomic Physics Laboratory

Ion-induce nucleation in gas phase is an important mechanism for grain formation in various circumstances. However, the number of works regarding this formation mechanism is very limited. To investigate the elementary processes of ion-nucleation mechanism, we recently developed a new apparatus (See, the presentation by N. Watanabe in this session). Using this apparatus, the cluster ion formation with an ion core mass-selected, which is the first stage of nucleation, can be observed quantitatively. In this presentation, we show the results of experiment on water-cluster ion formation in which free energies with the size of cluster have been determined. The experiment was performed at temperatures in range of 230-400 K with the supersaturation ratio of about 10^{-3} - 10^{-2} .

キーワード: 星間塵, クラスターイオン, 核生成
Keywords: interstellar grain, cluster ion, nucleation