

## 高圧下における FeO-SiO<sub>2</sub> 系の融解関係と沈み込んだ縞状鉄鉱層の行方 Melting relation on FeO-SiO<sub>2</sub> system at high pressure and the fate of the subducted banded iron formations

加藤 千恵<sup>1\*</sup>; 野村 龍一<sup>1</sup>; 廣瀬 敬<sup>2</sup>; 三宅 亮<sup>3</sup>; 大石 泰生<sup>4</sup>

KATO, Chie<sup>1\*</sup>; NOMURA, Ryuichi<sup>1</sup>; HIROSE, Kei<sup>2</sup>; MIYAKE, Akira<sup>3</sup>; OHISHI, Yasuo<sup>4</sup>

<sup>1</sup> 東京工業大学大学院地球惑星科学専攻, <sup>2</sup> 東京工業大学地球生命研究所, <sup>3</sup> 京都大学大学院地球惑星科学専攻, <sup>4</sup> 高輝度光科学研究センター

<sup>1</sup>Department of Earth and Planetary Sciences, Tokyo Institute of Technology, <sup>2</sup>Earth-Life Science Institute, Tokyo Institute of Technology, <sup>3</sup>Division of Earth and Planetary Sciences, Kyoto University, <sup>4</sup>Japan Synchrotron Radiation Research Institute

Subduction of banded iron formations (BIFs) may have played a significant role on the evolution of the core-mantle boundary (CMB) region and the chemical stratification at the topmost core. Almost all of the BIFs that had been deposited on the seafloor must have been subducted into the mantle and only a small portion was left at the surface today. Because of their high density, BIFs may have fallen down toward the CMB region. The amount of subducted BIFs is estimated to be  $2 \times 10^8 \text{ km}^3$ , which roughly matches with the total volume of the ultra-low velocity zones (Dobson and Brodholt, 2005, *Nature*). BIFs would be composed mainly of FeO and SiO<sub>2</sub> in the lower mantle because its oxidation state is close to iron-wustite buffer. We have performed melting experiments on FeO-SiO<sub>2</sub> system by laser-heated diamond-anvil cell technique at 25 - 140 GPa. FE-EPMA and FIB-SEM chemical analysis and observation of the texture of the recovered samples revealed that the liquidus phase was SiO<sub>2</sub> when starting from Fe<sub>2</sub>SiO<sub>4</sub> fayalite, and compositions of the quenched melt suggested that the composition of the eutectic point was extremely FeO-rich (<0.6 wt% SiO<sub>2</sub>). The solidus curve was constrained by observing the existence or no-existence of tiny quenched melt pools in each recovered samples. The solidus temperature at CMB pressure was 3,100 to 3,300 K that were lower than the solidus of pyrolite and the geotherm (Nomura et al., 2014, *Science*). These results imply that when the BIFs reach the CMB, they generate FeO-rich liquid that would be mixed with the basal magma ocean (Labrosse et al., 2007, *Nature*). This liquid would form a thin layer spread along CMB and react with the topmost core. Silicon content in liquid iron varies inversely to the oxygen fugacity when equilibrium with silicate melt (Ricolleau et al., 2011 *EPSL*). Assuming silicon-rich bulk core, exchange of silicon and oxygen would occur between the topmost core and FeO-rich silicate melt. This mechanism may account for the seismic wave speed anomaly observed at the topmost core which is thought to be honor to chemical stratification (Helffrich and Kaneshima, 2010 *Nature*; Buffett and Seagle, 2010 *JGR*).