

SGL42-09

会場:419

時間:4月29日 11:40-11:55

高精度 U-Pb ジルコン年代測定における結晶形態観察および微量元素存在度分析の意義：石鎚コールドロンに産する火成岩類を例に Significance of external morphology and zircon chemistry for precise U-Pb zircon dating

竹原 真美^{1*} ; 堀江 憲路² ; 谷 健一郎³ ; 吉田 武義⁴ ; 外田 智千² ; 清川 昌一⁵

TAKEHARA, Mami^{1*} ; HORIE, Kenji² ; TANI, Kenichiro³ ; YOSHIDA, Takeyoshi⁴ ; HOKADA, Tomokazu² ; KIYOKAWA, Shoichi⁵

¹九州大学 大学院理学府 地球惑星科学専攻, ²国立極地研究所, ³海洋研究開発機構地球内部ダイナミクス領域, ⁴東北大 学 大学院理学科 地学専攻, ⁵九州大学 大学院理学研究院 地球惑星科学部門

¹Department of Earth and Planetary Sciences, Graduate School of Sciences, Kyushu University, ²National Institute of Polar Research, ³Institute for Research on Earth Evolution, Japan Agency for Marine-Earth Science and Technology, ⁴Institute of Earth Sciences, Graduate School of Science, Tohoku University, ⁵Department of Earth and Planetary Sciences, Faculty of Sciences, Kyushu University

Improvement of U-Pb zircon dating by microbeam analysis has been provided opportunity of discussion about more detailed geological events. Recent analytical precision of less than 2% at Paleogene zircon allows us to investigate shorter period events such as crystallization differentiation in magma chamber. However, the highly precise U-Pb age data yield an importance of confirming their accuracy and assaying disturbance of U-Pb system and incorporation of exotic components. In this study, we introduce data processing method of the highly precise ages based on zircon morphology, trace element abundances as well as statistics.

The precise U-Pb zircon dating by using a sensitive high-resolution ion microprobe (SHRIMP II) at National Institute of Polar Research, Japan, was applied to igneous rocks of the Tertiary Ishizuchi Cauldron in the Setouchi volcanic belt of Miocene age in northwestern Shikoku. A primary ion beam of about 10 nA was used to sputter an analytical spot of about 40 μm diameter. A retardation lens system was utilized as a means to increase signal-to-noise ratio, and a secondary ion optics including slits of source and collector was adjusted to maximum transmission of the secondary ion under suitable mass resolution avoiding isobaric interferences on Pb isotopes. The surfaces of grain mounts were carefully washed with diluted HCl and ultra pure water to remove Pb contamination. A correction for common Pb was made on the basis of the measured ^{204}Pb and the model for common Pb composition.

Weighted mean ages were calculated from $^{206}\text{Pb}/^{238}\text{U}$ ratios corrected by ^{207}Pb . In order to ensure the accuracy of U-Pb age, age known zircon, OD-3, was analyzed together with unknown sample. Concentrations of Hf and rare earth element (REE) in zircons were also measured at the same analytical spot of U-Pb dating by SHRIMP.

Zircon grains from the Bansyodani-biotite-rhyolite were divided to two types based on the external morphology: sharply euhedral type and relatively rounded edge of prism and pyramid type. $^{206}\text{Pb}/^{238}\text{U}$ data of whole zircon grains were widely scattered beyond analytical uncertainty and show a weighted mean of 14.78 ± 0.18 Ma (mean square weighted deviation, MSWD: 3.4). On the other hand, the euhedral zircons yielded the weighted mean of 14.21 ± 0.19 Ma (MSWD: 1.0), whereas the relatively rounded zircons were older than the euhedral zircons, which suggestss the incorporation of exotic components.

Zircon chemistry supported the classification by the morphology and the U-Pb dating. An average of Hf contents of the euhedral zircons were 9523 ppm ranging from 8883 to 10496 ppm and those of the relatively rounded zircons were 8475 ppm ranging from 7616 to 8803 ppm. Hf contents of the euhedral zircons were higher than those of relatively rounded zircons. C1 chondrite-normalized REE patterns of the euhedral zircons were characterized by a large fractionation between light REE and heavy REE, large positive Ce anomalies and large negative Eu anomalies. In contrast, those of the relatively rounded zircons were enrichment of light REE, weaker anomalies of Ce and Eu. Difference of the zircon chemistry between the euhedral zircons and the relatively rounded zircons reflects source melt composition. Therefore, the external morphology, Hf content, and REE abundance are useful criteria for the data processing of the highly precise U-Pb age data.