

ゲスト分子の配向秩序化による Filled Ice Ih 構造メタンハイドレートの相変化 Phase changes of filled ice Ih methane hydrate induced by the orientational ordering of the guest molecules

田中 岳彦^{1*}; 平井 寿子¹; 松岡 岳洋²; 大石 泰生³; 八木 健彦¹; 大竹 道香⁴; 山本 佳孝⁴; 中野 智志⁵
TANAKA, Takehiko^{1*}; HIRAI, Hisako¹; MATSUOKA, Takahiro²; OHISHI, Yasuo³; YAGI, Takehiko¹; OHTAKE, Michika⁴
; YAMAMOTO, Yoshitaka⁴; NAKANO, Satoshi⁵

¹ 愛媛大学地球深部ダイナミクス研究センター, ² 岐阜大学, ³ 高輝度光科学研究センター, ⁴ 産業技術総合研究所, ⁵ 物質・材料研究機構

¹Geodynamics Research Center, Ehime University, ²Gifu University, ³Japan Synchrotron Radiation Research Institute, ⁴The National Institute of Advanced Industrial Science and Technology, ⁵National Institute for Materials Science

Oriental ordering of guest methane molecules in a filled ice Ih structure of methane hydrate (MH) was observed above 15 to 20 GPa at room temperature in a previous Raman study. However, the change in the fundamental structure was not observed at the pressure region by X-ray diffractometry. In this study, low-temperature and high-pressure experiments were performed with filled ice Ih structure of methane hydrate under pressure and temperature conditions of 2.0 to 77.0 GPa and 30 to 300 K, respectively, using diamond anvil cells and a helium-refrigeration cryostat. Distinct changes in the axial ratios of the host framework were revealed by In-situ X-ray diffractometry. Splitting in the CH vibration modes of the guest methane molecules, which was previously explained by the orientational ordering of the guest molecules, was observed by Raman spectroscopy. The pressure and temperature conditions at the split of the vibration modes agreed well with those of the axial ratio changes. The results indicated that orientational ordering of the guest methane molecules from orientational disordered-state occurred at high pressures and low temperatures, and that this guest ordering led to the axial ratio changes in the host framework. Existing regions of the guest disordered-phase and the guest ordered-phase were roughly estimated by the X-ray data. In addition, above the pressure of the guest-ordered phase, another high pressure phase was developed at a low-temperature region.

キーワード: メタンハイドレート, X 線回折, 高圧, ラマン分光

Keywords: Methane Hydrate, X ray diffraction, high pressure, Raman spectroscopy