

ACG06-01

会場:101A

時間:5月26日 11:00-11:20

インドネシア海大陸の速い水循環：「AM ラジオ」仮説 Quick water cycle over the Indonesian maritime continent: An "AM radio" hypothesis

山中 大学^{1*}

YAMANAKA, Manabu D.^{1*}

¹ 海洋研究開発機構大気海洋相互作用研究分野／神戸大学大学院理学研究科

¹DCOP, JAMSTEC / DP-GSS, Kobe U

平成 17～21 年度「地球観測システム構築推進プラン (JEPP)」による海大陸レーダー網 (HARIMAU) 構築および 21～25 年度「地球規模課題対応国際科学技術協力 (SATREPS)」による海大陸最先端研究拠点 (MCCOE) 構築により、海風水蒸気輸送、雲・降雨、河川水輸送の全てが日周期変化することが示された。

従ってインドネシア海大陸域の水循環は極めて速く、大規模現象や経年・季節内変動に伴って（日周期海陸風循環強度を通じて）雨量等の増幅はするが、日周期は基本的に維持される。この様相は「AM ラジオ」に似ている。その意味では、中緯度のような総観規模現象の連続的な観測・予測よりも、地域ごとの観測や毎日新たに（何度も）予測する方が適している。

キーワード: インドネシア海大陸, 雲対流, 水循環, 日周期, HARIMAU レーダー網, SATREPS-MCCOE

Keywords: Indonesian maritime continent, cloud convection, water cycle, diurnal cycle, HARIMAU, MCCOE

MJO role on Intraseasonal variation of stable isotope of Precipitation in Indonesia Maritime Continent

MJO role on Intraseasonal variation of stable isotope of Precipitation in Indonesia Maritime Continent

BELGAMAN, Halda aditya^{1*}; ICHIYANAGI, Kimpei¹; TANOUE, Masahiro²; SUWARMAN, Rusmawan³; YOSHIMURA, Kei⁴; MORI, Shuichi⁵; KURITA, Naoyuki⁶
BELGAMAN, Halda aditya^{1*}; ICHIYANAGI, Kimpei¹; TANOUE, Masahiro²; SUWARMAN, Rusmawan³; YOSHIMURA, Kei⁴; MORI, Shuichi⁵; KURITA, Naoyuki⁶

¹GSST Kumamoto Univ., ²School of Engineering, The University of Tokyo, ³Bandung Institute of Technology, ⁴AORI, The Univ. of Tokyo, ⁵JAMSTEC, Japan, ⁶Nagoya Univ.

¹GSST Kumamoto Univ., ²School of Engineering, The University of Tokyo, ³Bandung Institute of Technology, ⁴AORI, The Univ. of Tokyo, ⁵JAMSTEC, Japan, ⁶Nagoya Univ.

MJO (Madden-Julian Oscillation) is one of the disturbances for Asian Monsoon in the Maritime Continent Area. Intraseasonal variability of precipitation in the Indonesia Maritime Continent (IMC) is mainly due to The Madden-Julian Oscillation (MJO), cold surge, or other synoptic scale disturbance. This study examined the relationship between MJO and stable isotope in precipitation over the IMC. Observation and simulation model data from isotope circulation model was used. From 2001 - 2009, 10 main MJO events were detected and 6 of the event occurred at Boreal Winter season (Asian Monsoon). Temporal and spatial analysis from simulated stable isotopes model reveals that isotopic variation is correlated to MJO event in different phase for different stations. Generally, $\delta^{18}\text{O}$ in precipitation became lighter in most observation station at phase 3, 4 and 5 at western part of IMC, and phase 4, 5, and 6 at north east part of IMC. Spatial distribution of $\delta^{18}\text{O}$ in precipitation for each MJO phase show that mainly MJO governed $\delta^{18}\text{O}$ variability in intraseasonal timescale for IMC area. Further investigation showed clear signal of MJO in $\delta^{18}\text{O}$ was observed at Bukit Tinggi (GAW) station only; it is confirmed with result from Isotope Circulation Model (ICM). When MJO is in active phase (enhanced) precipitation in western part of IMC, precipitable water was came from Indian Ocean and South IMC Ocean.

キーワード: asian monsoon disturbance, stable isotope, Madden Julian Oscillation, moisture transport analysis
Keywords: asian monsoon disturbance, stable isotope, Madden Julian Oscillation, moisture transport analysis

南シナ海夏季モンスーンオンセットの年々変動および10年規模変動に対する海洋の役割

Relative role of the ocean for interannual and decadal variations in summer monsoon onset over the South China Sea

今川 新^{2*}; 樋口 篤志¹

IMAKAWA, Shin^{2*}; HIGUCHI, Atsushi¹

¹ 千葉大学 環境リモートセンシング研究センター, ² 千葉大学 大学院理学研究科

¹Center for Environmental Remote Sensing (CEReS), Chiba University, ²Graduate School of Science, Chiba University

In this study, we reveal a difference of mechanisms with interannual variation a 15-years variation in the South China Sea Summer Monsoon (SCSSM) onset. The SCSSM onset occurs in active convections over the South China Sea (SCS), when convections are active over the Philippine Sea (PS). Variations in the SCSSM onset are affected by the variations in the convective activities over SCS and PS. The increase in the sea surface temperature (SST) over the Western Pacific in recent years causes the more active convection over the PS, therefore the SCSSM onset is advanced in 15-years variation as already pointed out by Kajikawa and Wang (2012).

Moreover, we elucidate the relation between the SCSSM onset date anomaly and aggregated SST over the SCS from January to March. The correlation is good between the SCSSM onset date anomaly and the aggregated SST over SCS, while a poor relation is found between the 15-years variation oriented signal in onset date and the SST in SCS. When SCSSM onset date is delayed, the aggregated SST in SCS has a tendency to high temperature. On the other hand, the correlation is good between the 15-years variation in the SCSSM onset date and the aggregated SST over the PS. The SST anomalies over the SCS are influenced by the frequencies and strength of cold surges during boreal winter. Strong and longer cold surges bring the lower SST over the SCS. The cold surges are brought about by the anticyclone over the Eastern Eurasian continent. An arctic sea-ice decline instigates the high-pressure deviation over the Eurasian Continent. Then relatively lower land surface temperature in middle latitude, and bring the cold surge over the SCS.

キーワード: 夏季モンスーン, オンセット, 南シナ海

Keywords: summer monsoon, onset, South China Sea

フィリピンにおける降雨と大気循環の気候学的季節変化 Climatological seasonal changes of rainfall and circulation in the Philippines

松本 淳^{1*}; Nguyen-Le Dzung¹; Villafuerte Marcelino II²
MATSUMOTO, Jun^{1*}; NGUYEN-LE, Dzung¹; VILLAFUERTE, Marcelino II²

¹ 首都大学東京 地理学教室, ² フィリピン宇宙気象局

¹Department of Geogreaphy, Tokyo Metropolitan University, ²PAGASA, Philippines

Climatological seasonal changes of rainfall and lower tropospheric circulation in the Philippines were analyzed by utilizing 5-day mean TRMM 3B-42 and station rainfall data provided by PAGASA, and ERA-Interim wind data for the period 1998-2013. In particular, climatological onset and withdrawal processes of the southwest monsoon were investigated.

It was found that the onset of southwest monsoon occurred abruptly in mid-May. It started from the north in the Philippines both in rainfall and wind, which showed a peculiar feature in this region. After the onset, anti-cyclonic flow from the Pacific high was predominant, and it changed into cyclonic flow in mid-June. Easterlies still remained in the south until early July, afterwards SW monsoon covered the whole country and enhanced from late July.

Southwest monsoon began to retreat from the north in mid- September, and fully retreated from the southern tip of the Philippines in late October.

Acknowledgment: Part of this study was supported by The JSPS KAKENHI, The GRENE program of the MEXT, and the Asian Human Resource Fund of the Tokyo Metropolitan Government.

キーワード: モンスーン, 季節変化, モンスーン開始, モンスーン撤退, 雨季

Keywords: monsoon, seasonal changes, monsoon onset, monsoon withdrawal, rainy season

Long-term Regional Precipitation Disparity in Northwestern China and Its Driving Forces

Long-term Regional Precipitation Disparity in Northwestern China and Its Driving Forces

LEE, Harry F.^{1*}

LEE, Harry F.^{1*}

¹Department of Geography, The University of Hong Kong

¹Department of Geography, The University of Hong Kong

Precipitation in Northwestern China (NW China) is characterized by salient regional differences. Yet, the long-term regional precipitation disparity in NW China still remains insufficiently-explored. In the present study, we base on historical documentation to derive the fine-grained precipitation indices of two macro regions in NW China between AD580 and 1979 to (1) determine the multi-decadal to centennial regional precipitation disparity in NW China; and (2) find the major driving forces behind it. Wavelet analysis is applied. Our results show that there is significant regional discrepancy of precipitation change in NW China over extended period. Besides, the association between the regional precipitation disparity in NW China and various modes of atmospheric circulation (Asian Summer Monsoon, Arctic Oscillation, Pacific Decadal Oscillation, and North Atlantic Oscillation) is significant and characterized by a regime shift during the transition from the warm episode to the Little Ice Age in the 14th century. Most importantly, the ~180 to 240 year cycle of the El Niño-Southern Oscillation is found to be the most prominent pacemaker of regional precipitation disparity in NW China at the long-term temporal scales. Our findings help to demonstrate which atmospheric circulation is primarily responsible for the long-term regional precipitation disparity in NW China, which may have important implications for water resource management there.

キーワード: precipitation, moisture, atmospheric circulations, Asian Summer Monsoon, El Nino-Southern Oscillation, North-western China

Keywords: precipitation, moisture, atmospheric circulations, Asian Summer Monsoon, El Nino-Southern Oscillation, North-western China

Future Projection of Himalayan Monsoon Season Precipitation by CMIP5 Models under Warming Climates

Future Projection of Himalayan Monsoon Season Precipitation by CMIP5 Models under Warming Climates

Kadel Indira^{1*}; 山崎 剛¹

KADEL, Indira^{1*}; YAMAZAKI, Takeshi¹

¹Department of Geophysics, Tohoku University

¹Department of Geophysics, Tohoku University

I. Introduction: Livelihood of the Himalayan people depends largely upon precipitation. Any variability on its regular pattern exacts profound social and economical costs.

In response to increasing anthropogenic forcing, the global monsoon precipitation is expected to amplify over most of the global region [1]. However, the response varies from region to region owing to various factors such as topography, forcing etc. [2]. Recent studies found that monsoon precipitation over India is likely to increase under the global warming scenario with enhanced variability [3, 4]. Similarly, it has also been found that precipitation over the Himalaya particularly over Nepal is poorly comparable with all India rainfall [5]. Since, study on precipitation over the world's unique geographical region, Himalaya under warming scenario is very limited, this study is aimed to investigate the future precipitation change over the Himalaya.

II. Models, Data and Methodology: We employed 26 general circulation models (GCMs) participating in Coupled Model Intercomparison Project Phase 5 (CMIP5) to evaluate their skill in simulating present monsoon season (June-September) precipitation (MSP) climatology (1979-2005) over Himalaya (20-35 °N latitude and 75-94 °E longitude). The skill of the models was judged based on their ability to represent annual cycle, spatial pattern and variability using Taylor diagram. Global Precipitation Climatology Project Version 2.2 monthly data was used for the model evaluation. The best model was selected whenever normalized standard deviation (STD) lies in 0.8-1.2, correlation coefficient (CC) >0.5 and centered root mean square (RMS) difference <1. To assess the future change, simulations from Representative Concentration Pathway 2.6 (RCP2.6), 4.5 (RCP4.5) and 8.5 (RCP8.5) of selected best models of 2070-2099 were compared with the historical runs of 1970-1999. Monthly precipitation of each model from one ensemble (r1i1p1) run was used.

III. Results and Discussions: All the models reproduced annual cycle of mean precipitation over the Himalaya more or less accurately with CC is greater than 0.7. Among them, only 13 models met the criteria mentioned in section II for reproducing annual cycle. We found that simulating the spatial pattern over the complex terrain became a major challenge for the models. Out of the 26 GCMs, only nine of them were comparatively better than others to reproduce spatial pattern with high CC, less variability and error. Likewise, simulation of the STD of MSP was also found to be a problematic one. Taylor diagram of MSP indicated that only nine models performed relatively well. Our analysis revealed that neither of the models performed equally to simulate all three matrices used in this study. Thus, only five models; HadGEM2-ES, MPI-ESM-LR, MPI-ESM-MR, NorESM1-M and NorESM1-ME were selected based on their skill to simulate all three matrices. Ensemble projections of the best models under RCP8.5 scenario indicated that MSP might be increased by 11 % compared to present climatology with amplified variability over heavy rainfall zone. Multi-model mean annual cycle of monthly precipitation showed the enhancement of monthly precipitation during monsoon season.

IV. Conclusion: Precipitation over the Himalaya simulated by 26 GCMs was evaluated. No model was found to simulate all aspects of MSP equally. Relatively, high resolution earth system models performed better than the low resolution models. Ensemble mean of five reliable models projected that the MSP will increase over the Himalaya in future warmer climate with enhanced variability over the heavy precipitation zones.

References:

- [1] Lee, J.-Y. and B. Wang, 2014, *Clim. Dyn.*, **42**, 101-119
- [2] Turner, A. G. and H. Annamalai, 2012, *Nat. Clim. Chang.*, **2**, 587-595
- [3] Menon et al., 2013, *Earth Syst. Dyn.*, **4**, 287-300
- [4] Sharmila et al., 2015, *Glob. Planet. Chang.*, **124**, 62-78
- [5] Shrestha et al., 2000, *Int. J. Climatol.*, **20**, 317-327

ACG06-P01

会場:コンベンションホール

時間:5月26日 18:15-19:30

Keywords: Monsoon precipitation, Future projection, CMIP5, Himalaya