

Quantifying warming effect due to urbanization based on comparative measurements of the surface energy budget

ANDO, Tomoyoa^{1*} ; UEYAMA, Masahito¹ ; KOSUGI, Yoshiko²

¹Graduate School of Life and Environmental Sciences, Osaka Prefecture University, ²Graduate School of Agriculture, Kyoto University

Land use changes from natural ecosystems to urban built-up have strong influence on local climates. The urban heat island has received much attention as an important environmental problem. However, detailed mechanisms have not been clear enough. In this study, we conducted comparative measurements of the surface energy balance at an urban built-up and a forest, and evaluated factors contributing to surface warming due to the land use change based on a methodology called temperature decomposition.

We measured the energy balance at the urban built-up in Sakai, Osaka and at a cypress plantation (Kiryu Experimental Watershed) in Shiga. Sensible and latent heat fluxes were measured using the eddy covariance method. The ground heat flux for the urban area was estimated using the objective hysteresis model with relevant plot-scale experiments. In this study, we evaluated the urbanization effect from the forest plantation to the urban in terms of changes in surface temperatures associated with surface albedo, aerodynamic resistance, Bowen ratio, and ground heat flux.

In daytime, the urbanization increased net surface temperatures by 5.5 K in the summer (June to August 2014) and 3.8 K in the winter (January, February, and December 2014). The daytime warming was mostly contributed by increased surface resistance due to the urbanization in the summer (4.4 K) and the winter (3.7 K). In the urban built-up, the decreased surface roughness due to a high building density restricted the heat transfer redistribution into the atmosphere. Consequently, we found that the decreased surface resistance was the most effective factor to warm the urban surface temperatures in daytime. In nighttime, the urban surface temperature was higher than those in the forest in the summer (1.9 K) and the winter (2.5 K). Change in ground heat flux most strongly contributed to the nighttime warming in the summer (4.2 K) and the winter (5.2 K). In the urban built-up, a lot of incoming energy was stored in buildings and impervious roads in daytime. This energy was radiated and warmed up the urban surface in nighttime. The estimated net increases in the surface temperature were lower than those estimated using the bulk model in daytime and nighttime. The discrepancy could be caused by uncertainties associated with estimated radiative fluxes, energy imbalance of turbulent fluxes, and no consideration of anthropogenic heat in the urban built-up.

A decline of a Japanese oak by sulfuric acid of an air pollutant, and reproduction by charcoal

OMORI, Teiko^{1*}

¹FOMER TOHO UNIVERSITY

A fossil fuel generates sulfuric acid by combustion. Sulfuric acid becomes an air pollutant, moves by wind and adheres to trees. Sulfuric acid which adhered to trees is dropped on a root by rain, and acidifies soil. If soil acidifies, a metal ingredient will dissolve. The dissolved metallic ion is absorbed into the trees. The phosphoric acid in trees combines with metallic ion. The combined phosphoric acid becomes inactivate and trees wither. The combined phosphoric acid becomes inactivate and trees wither. If the tannin contained in trees combines with a metal ion, it loses the insect control effect. Rain water becomes alkaline solution by charcoal. An alkaline solution changes metal ion into hydroxide. Trees cannot absorb metal hydroxide. As a result, trees are saved. If charcoal is scattered on the tree of the declined Japanese oak, it dropped a seed after 3 years and budded after 4 years.

Keywords: air pollutant, acidifies soil, acidifies soil, tannin, metallic phosphate

Long-term warming effect on soil respiration in warm-temperate evergreen broad-leaved forest in Kyusyu

TERAMOTO, Munemasa^{1*} ; LIANG, Naishen¹ ; TAKAGI, Masahiro² ; ZENG, Jiye¹

¹Center for Global Environmental Research, National Institute for Environmental Studies, ²Faculty of Agriculture, University of Miyazaki

Soil respiration is composed of heterotrophic respiration and plant root respiration, and is the second largest carbon flux in the terrestrial ecosystem. As the heterotrophic respiration increases exponentially with soil temperature, its positive feedback to the global warming has become a concern.

In the middle of December 2008, a multi-channel automated chamber measurement system was installed at the Tano forest science station of Miyazaki University, which is a warm-temperate broad-leaved forest site in Kyusyu. We prepared 10 trenched plots with 5 of them artificially warmed by +2.5°C by infrared heaters 1.6 m above the surface for long-term measurement of warming effect on soil respiration.

The average value of soil respiration in control and warmed plots were, respectively, 3.45 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ and 3.53 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ in 2009, 3.81 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ and 4.07 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ in 2010, 3.31 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ and 3.81 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ in 2011, 3.02 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ and 3.79 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ in 2012, 3.08 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ and 3.33 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ in 2013. The annual warming effect increased soil respiration by 5.4% in 2009, 9.9% in 2010, 18.4% in 2011, 29.5% in 2012, and 11.5% in 2013. The warming effect showed an increasing trend until 2012, but decreased in 2013 due to fewer amount of rainfall in summertime, which caused a much lower moisture in soil.

Keywords: soil respiration, global warming, chamber, forest soil




Fig 1. Seasonal variation of soil temperature, soil moisture and CO₂ efflux of each treatment.

Control of Soil Carbon Dynamic of Southeast Asian Ecosystems

LIANG, Naishen^{2*} ; TERAMOTO, Munemasa² ; TANG, Zhenghong³ ; FLETCHER, Christine D.⁴

¹NIES, ²NIES, ³XTBG, ⁴FRIM

The tropical ecosystems have been estimated to be a large carbon source (1.3 Pg C yr^{-1}) due to deforestation and forest degradation, even the global terrestrial carbon sink has been large (1.1 Pg C yr^{-1}) in recent decades. Reducing Emissions from Deforestation and Forest Degradation (REDD) is an effort to create a financial value for the carbon stored in forests, offering incentives for tropical region to reduce emissions from forested lands and invest in low-carbon paths to sustainable development. The ultimate goal of this study is to update REDD mechanism through improved forest management by evaluation of effects of logging and land-use change on soil carbon emission of tropical forests. This study was conducted in a lowland primary forest at Pasoh Forest Reserve ($2^{\circ}58'N$, $102^{\circ}18'E$; $75\sim 150\text{m a.s.l.}$) and a mountainous tropical forests at Temenggor concession area ($5^{\circ}33'N$, $101^{\circ}36'E$; $800\sim 900\text{m a.s.l.}$) in Peninsular Malaysia. About 50~65% biomass was harvested and soil temperature increased about $3^{\circ}C$ with SMS, resulting value of the carbon stock lost about $2,577 \text{ US\$ ha}^{-1}$ following the first year of logging. On the other hand, under low-impact harvest condition, only about $1,773 \text{ US\$ ha}^{-1}$ was lost following the first year of logging. Result suggests that this low-impact harvest system would achieve about $804 \text{ US\$ ha}^{-1}$ of REDD credit partially contributed from mitigating soil degradation of about $169 \text{ US\$ ha}^{-1}$.

Keywords: Automated chamber, LULUC, moisture, soil carbon, tropical ecosystem

Spatial variations in larch and soil nitrogen isotope ratio along forest-grassland gradient in northern Mongolia

FUJIYOSHI, Rei^{1*} ; TSUJIMURA, Maki² ; LOPEZ, Larry c.m.³ ; SUGIMOTO, Atsuko⁴

¹Graduate School of Environmental Science, Hokkaido University, ²Faculty of Life and Environmental Sciences, University of Tsukuba, ³Faculty of Agriculture, Yamagata University, ⁴Faculty of Environmental Earth Science, Hokkaido University

The underlying processes that affect the Nitrogen isotope ratio of plant and soil, which is known as a good indicator of N dynamics and shows dependency on climate, is not yet clearly understood.

In this study, we compare the nitrogen isotope ratio of single plant species (*Larix sibirica* Ledeb.) and soils along the forest-grassland gradient of a forest-steppe ecotone in Mongolia. Larch needles and soils were sampled during the growing season (May- August) from 2004-2012 at several sites in seven study areas.

The results showed the clear spatial pattern in needle, soil nitrogen isotope ratios, and its difference ($\Delta\delta^{15}\text{N}$) along forest-grassland gradient and the pattern corresponded to humus type (mor/mull). $\Delta\delta^{15}\text{N}$ also had significant correlations with needle $\delta^{13}\text{C}$ and C/N ratio of bulk soil, suggesting the $\Delta\delta^{15}\text{N}$ change relates to water, light conditions and soil N availability. Isotope mass-balance model was applied to investigate the processes affecting the spatial pattern of $\Delta\delta^{15}\text{N}$.

Keywords: ecotone, Mongolia, nitrogen isotope ratio, plant and soil

Seasonal variation in ratio of soil respiration to ecosystem respiration at Takayama estimated from $\delta^{18}\text{O}$ measurement

MURAYAMA, Shohei^{1*}; SAWADA-TAKAMURA, Chikako²; SAIGUSA, Nobuko³; ISHIDOYA, Shigeyuki¹; WATANABE, Tsutomu⁴; ITO, Akihiko³; MORIMOTO, Shinji⁵; KONDO, Hiroaki¹; MAEDA, Takahisa¹; MURAOKA, Hiroyuki⁶; USAMI, Tetsuyuki¹; AOKI, Shuji⁵; NAKAZAWA, Takakiyo⁵

¹AIST, ²AORI, The University of Tokyo, ³NIES, ⁴ILTS, Hokkaido University, ⁵Graduate School of Science, Tohoku University, ⁶River Basin Research Center, Gifu University

Forest ecosystems are one of the important reservoirs in the global carbon cycles. However, environmental factors governing variation in carbon dioxide (CO_2) flux between the atmosphere and the forest ecosystems have not been fully understood, which leads to very large uncertainty in future predictions of response of forest ecosystems to climate change. For more precise prediction of the future global carbon budgets, better understandings of each process in the carbon cycle in the ecosystem are necessary. A stable oxygen isotopic ratio ($\delta^{18}\text{O}$) in CO_2 is a unique tracer reflecting not only the carbon cycle but also the hydrological cycle. Using difference of isotopic fractionation in ^{18}O between photosynthetic and respiratory processes and between soil and leaf respiratory processes, gross CO_2 fluxes in each of the processes have been estimated in some previous studies at relatively short time scales from diurnal to a few months. In this study, isotopic measurements of not only CO_2 in the atmosphere and soil air but also soil water and atmospheric water vapor were made at a cool-temperate deciduous forest, Takayama (TKY; 36°08'N, 137°25'E, 1420 m a.s.l.) in the growing seasons during 2006-2009. From the obtained data, $\delta^{18}\text{O}$ values in CO_2 emitted due to soil (R_s) and leaf respirations and total ecosystem respiration (R_{ec}) during the nighttime were calculated. Then, seasonal variation in relative contribution of R_s to R_{ec} was estimated from mass balance equations. The obtained result was compared with variation in the ratio of R_s to R_{ec} estimated from soil chamber and eddy covariance flux measurements. The result from the $\delta^{18}\text{O}$ measurements showed that the ratio of R_s to R_{ec} increased from about 0.5 in late spring to almost 1 in early autumn though the estimated ratios were very scattered. Such a seasonal pattern was similar to that estimated from the flux measurements. In our presentation, factors governing the seasonal variation will also be discussed by comparison with the results simulated for TKY using a process-based ecosystem model (VISIT) and a multi-layer canopy model (MINCER).

Keywords: oxygen isotope, forest ecosystem, ecosystem respiration, soil respiration

Long-term observation of CO₂ flux over a larch forest in northern foot of Mt. Fuji

TAKAHASHI, Yoshiyuki^{1*} ; SAIGUSA, Nobuko¹ ; IDE, Reiko¹ ; HIRATA, Ryuichi¹

¹Center for Global Environmental Research, National Institute for Environmental Studies

Larch forest is representative vegetation type of widespread region in northeast Eurasia where vulnerability to climate change is expected to be significant. Since length of growing season of the vegetation is affected by temperature, evaluation of climate change feedback in the carbon exchange process in the ecosystem has great importance. National Institute for Environmental Studies has been operating long-term observation of CO₂ flux over a larch plantation at the Fuji Hokuroku flux observation site in northern foot of Mt. Fuji since 2006. We have accumulated 9-year records of observational data set. The observation is to be continued until 2025. In this presentation, we will investigate year-to-year variation of CO₂ exchange and its relationship with environmental factors based on the observational data.

Keywords: carbon dioxide, flux, forest ecosystem, climate change, long-term monitoring, micrometeorology

Effects of waterlogging on carbon isotope discrimination of *Larix gmelinii* during photosynthesis

LI, Fang^{1*} ; SUGIMOTO, Atsuko¹

¹Graduate school of environmental science, Hokkaido university

The carbon isotope discrimination during photosynthesis reflects various environmental factors, such as VPD, solar radiation, soil moisture, salinity, air pollution and so on. It has been used to estimate environmental condition, or calculation of carbon isotope mass balance equation to know a flow of carbon. It is well known that soil moisture is one of the major factors controlling carbon isotope discrimination: discrimination decreased under drought condition due to a decrease of stomatal conductance. On the other hand, there is almost no research on carbon isotope discrimination under extremely wet condition. *Larix gmelinii* is one of the major species of larch trees, which constitute eastern Siberian Taiga. In 2007, extreme wet event has been reported near Yakutsk. In this research, pot experiment with *Larix gmelinii* saplings was conducted to obtain carbon isotope discrimination during photosynthesis under waterlogging condition which may happen more frequently in the future. Under waterlogging condition, both photosynthesis rate and stomatal conductance of all three larch saplings used for experiment decreased, while only one sapling among three showed clear decrease in carbon isotope discrimination. Although the decrease in carbon isotope discrimination was not so clear, needles on current year stem, which seemed to be formed with C fixed during the experiment, showed slightly higher $\delta^{13}\text{C}$ in waterlogging treatment than that in control. This result suggests that waterlogging condition makes stomatal conductance decreased.

Keywords: carbon isotope discrimination, photosynthesis, stomatal conductance, waterlogging, larch, experiment

Effects of the expansion of vascular plants in *Sphagnum*-dominated bog on carbon balance

HIRANO, Takashi^{1*} ; YAMADA, Hiroyuki¹ ; TAKADA, Masayuki² ; FUJIMURA, Yoshiyasu³ ; FUJITA, Hiroko⁴ ; TAKAHASHI, Hidenori⁵

¹Research Faculty of Agriculture, Hokkaido University, ²Faculty of Humanity and Environment, Hosei University, ³Nippon Koei, ⁴Field Science Center for Northern Biosphere, Hokkaido University, ⁵Hokkaido Institute of Hydro-Climate

Peatland ecosystems have accumulated a huge amount of soil organic carbon for millennia. However, it is reported that the soil carbon has become vulnerable because of global warming and land-use change. Such disturbances potentially enhance oxidative peat decomposition, and consequently a large amount of CO₂ is emitted into the atmosphere. In Sarobetsu Mire in northern Hokkaido, which has the largest *Sphagnum*-dominated bog in Japan, vascular plants, *Sasa*, have invaded into *Sphagnum* bog because of the change of the hydrological environment, which was due to drainage for agriculture. To understand the effects of the expansion of *Sasa*-dominated area in *Sphagnum* bog on carbon balance, we measured CO₂ flux using the eddy covariance technique at a *Sphagnum*-dominated area (B site) and a *Sasa*-dominated area (T site) and compared CO₂ balance between the two sites.

Flux measurement was conducted during the four snow-free seasons from mid-April through early November in 2007 to 2010. Eddy sensors of CO₂ (LI7500, Licor) and wind speed (CSAT3, Campbell) were installed at the heights of 2.0 and 2.3 m, respectively, at B and T sites. The outputs from the sensors were recorded with a datalogger (CR1000, Campbell) at 10 Hz. Net ecosystem CO₂ exchange (NEE) was calculated on a half-hourly basis as a sum of eddy CO₂ flux and CO₂ storage change calculated from CO₂ concentration measured by LI7500. NEE was partitioned into gross primary production (GPP) and ecosystem respiration (RE) using an empirical conventional method.

Both GPP (gross ecosystem photosynthesis) and RE were larger at T site than B site. In 2008 with dry summer, cumulative NEE for 6.5 months of the snow-free season was -129 and -179 gC m⁻², respectively, at T site and B site. The negative NEE values indicate that the two peatland ecosystems functioned as CO₂ sinks in the season. In addition, sink strength was higher at T site. In 2010 with hot, wet summer, however, cumulative NEE was -238 and -159 gC m⁻², respectively, at T site and B site. Although NEE increased largely at B site, it decreased at T site. *Sphagnum* moss grew more under the hot and moist environment, which increased GPP more than RE. In contrast, although growth and resultant GPP also increased at T site, RE increased more than GPP because of high temperature. This result suggests that net CO₂ uptake will decrease by the invasion of *Sasa* plants under the warming environment in the near future.

Keywords: Peatland, CO₂ flux, Eddy covariance technique, Global warming

Detection of vegetation phenology across Japan by using digital time-lapse cameras

IDE, Reiko^{1*}; OGUMA, Hiroyuki¹; NAKAJI, Tatsuro²; NASAHARA, Kenlo³; ICHII, Kazuhito⁴; NAGAI, Shin⁴; ISHIDA, Akito⁵; NAKAMURA, Kazuhiko⁶; FUJIWARA, Akio⁶; WATANABE, Ryuichi⁷

¹National Institute for Environmental Studies, ²Hokkaido University, ³University of Tsukuba, ⁴Japan Agency for Marine-Earth Science and Technology, ⁵Kyoto Prefectural University, ⁶University of Tokyo, ⁷Shinsyu University

Many recent studies have reported earlier timings of leaf flush and later timings of leaf fall around the world with progress of recent global warming. Seasonal change (phenology) of vegetation is an indicator for the climate change, and also is one of the important parameters for terrestrial ecosystem models to estimate the carbon balance. For detection of the long-term changes in vegetation across wide areas, remote sensing technique is expected as a useful tool. However, due to the limitations of spatial resolution and observing frequency, it is difficult to detect the species specific environmental responses by satellite. Therefore, we collected high temporal and spatial resolution images by using digital time-lapse cameras. The purpose of this study is to detect phenology at a community or a species level for various types of vegetation across Japan, and to analyze the relationships between phenology and temperature.

We used the images of live camera archives such as "Phenological Eyes Network (PEN)", "Internet Nature Information System" of the Ministry of the Environment, "Mt. Hiei live camera" of Kyoto Prefectural University, "Cyberforest" of Tokyo University and so on, which were taken in 2002-2014. Images of approximately 120 site-year at 20 sites across Japan were analyzed. The areas of interest for vegetation a community or a tree within the images were selected for analysis. RGB digital counts in each pixel in the areas were extracted and Green Ratio Index ($GR = G / (R + G + B)$), which varies seasonally reflecting leaf greenness, was calculated at daily step. The time series of GR showed the maximum rate of change on the timings of leaf flush and leaf coloring, consequently, the dates of start/end of green leaves (SOG/EOG) were estimated. Furthermore, we investigated the relationships of the inter-annual variations between SOG/EOG and the temperature at each site.

The estimated SOG/EOG showed large variations across both years and areas, although earlier or later phenological trend during 2002-2014 was not recognized so far. Nationwide earlier SOG in 2002, 2004 and 2009, and later EOG in 2005, 2010 2012 were found. When the green leaved season started earlier than other years, it tended to end relatively later. Vice versa, when the green leaved season started later than other years, it tended to end relatively earlier. Such tendency of earlier or later SOG sometimes differed from the districts even in the same year. SOG or EOG of each community or species had high correlations with the mean temperature during respectively March-April or September at each sites. The impact of the climate change was discussed from the view point of species-specific and site-specific temperature sensitivities of the estimated SOG/EOG.

For assessment of the impact of climate change on vegetation, we need to observe the phenology at more sites for longer term. We plan to install more cameras and to analyze much more data including the uploaded images on the Internet. In addition, it is necessary to keep updating the existing camera systems and to administrate the observation networks sustainably.

Keywords: Time-lapse camera, Phenology, RGB, temperature sensitivity




Fig.1 Study sites of PEN(○), Internet Nature Information System(▲), Cameras of National Institute for Environmental Studies(◇), Mt.Hiei live camera(☆), Cyberforest(*),and Mt Chokai live camera(+).

Development of Above Ground Biomass estimation method using satellite data in Japan

BORJIGIN, Habura^{1*} ; HAYASHI, Masato¹ ; SAIGUSA, Nobuko¹ ; YAMAGATA, Yoshiki¹


¹National Institute for Environmental Studies

As a measurement method of Forest Above Ground Biomass, getting basic information of tree category, height and DBH from fieldwork, then calculate the volume, multiply expansion factor and wood density to get the result is widely used. This method can measure the forest above ground biomass correctly, however it is hard for us to make wall-to-wall map when monitoring large area of land. Satellite Remote Sensing is an effective way to estimate large area of forest distribution. In this study, our purpose is to investigate forest above ground biomass in the whole terrestrial land of Japan using ground truth data (fieldwork data from forest above ground biomass) and satellite data. We mainly used MODIS reflectance data from 2000 to 2009, Land Surface Reflectance data, Vegetation Index, LAI then using Random Forest method to estimate biomass.

We divided ground truth data to two groups, one group (90%) used as building Random Forest learning model. After the learning model was built, another group of data (10%) was used for estimating above ground biomass. Then we did the regression analysis between the estimation result and ground truth data. Accuracy assessment was carried out by calculation of coefficient of determination and RMSE from regression analysis. The results showed R² is 0.6, RMSE is 26.37 (t ha⁻¹). Estimation accuracy from deciduous forest and evergreen forest, R² is 0.4 and 0.53, RMSE is 24.29 (t ha⁻¹) and 27.34 (t ha⁻¹), respectively. Evergreen forest showed an higher accuracy. Also, we found out that low biomass (<100 t ha⁻¹) and high biomass (>200 t ha⁻¹) showed bigger estimation error. We compared the estimation result (acquired from forest above ground biomass of whole Japan) to prefectural data and forest registration, the comparison results showed coefficient of determination is 0.95, slope is 1.68 times, which is much lower than verification result (1.86 times) from Forestry and Forest Products Research Institute.

In this study, we confirmed that combine satellite data and fieldwork data, using Random Forest machine learning model, the large area of forest above ground biomass can be effectively estimated. In future study, compare the Random Forest to other models then get more accurate estimation is one of our goals. Also, carbon emission from deforestation, typhoon and other disturbance caused forest biomass change should be considered as well.

Keywords: Satellite data, MODIS, Above Ground Biomass, RandomForest




Forest resources monitoring using spaceborne LiDAR over Borneo

HAYASHI, Masato^{1*} ; BORJIGIN, Habura¹ ; SAIGUSA, Nobuko¹ ; YAMAGATA, Yoshiki¹ ; HIRANO, Takashi²

¹National Institute for Environmental Studies, ²Research Faculty of Agriculture, Hokkaido University

Forests serve as a primary reservoir of terrestrial carbon. Recently, the technology for forest resources monitoring at large-scale is required for the purpose of understanding the global carbon cycle or supporting REDD+ scheme. Satellite remote sensing can provide the most practical mean for the large-scale observation, and spaceborne LiDAR is particularly expected. This sensor transmits a laser pulse from space to the Earth's surface, and records the time-varying return signal intensity as a waveform. The waveform makes it possible to analyze vertical structure near the surface. ICESat/GLAS was the only spaceborne LiDAR so far, which was operated by NASA from 2003 to 2009. However, there are some future plans of spaceborne LiDAR, such as ICESat-2, GEDI, and MOLI, and they are expected to be used for forest monitoring. This study aims to clarify the potential of spaceborne LiDAR to quantitatively observe forest biomass at large-scale. We targeted Borneo as a study area, where is one of the most important areas for REDD+ because the forest area in the island has been rapidly decreasing in recent years. First, we measured above-ground biomass (AGB) by the Bitterlich method at 37 plots, where corresponded with the GLAS footprints. And, we used the field measured data as the training data to develop an empirical model for estimating AGB from the GLAS waveform parameters. As a result of cross-validation, the accuracy of this model (root-mean-square error) was 32.1 Mg ha⁻¹. Next, we applied the developed model to the 127,862 points GLAS data over Borneo to assess the forest resources in the island. As a result, some features were found as follows: (1) the average AGB in Borneo was 183.1 Mg ha⁻¹, (2) the AGB of evergreen broadleaved forest was nearly two times larger than that of mangrove forest or shrub land, and (3) the total AGB in Borneo was 9.81 Gt. Next, we developed a wall-to-wall map of forest biomass over Borneo using MODIS satellite image (see figure), because GLAS observed at only discrete points. These results indicate that spaceborne LiDAR is a suitable sensor for forest resources monitoring at large-scale.

Keywords: Forest biomass, Spaceborne LiDAR, ICESat/GLAS, Borneo



Development of Sun-Induced Chlorophyll Fluorescence database based on ecosystem tower measurement

KATO, Tomomichi^{1*}; TSUJIMOTO, Katsuto²; NASAHARA, Kenlo³; AKITSU, Tomoko³; NAGAI, Shin⁴; ONO, Keisuke⁵; SAITOH, Taku M.⁶; MURAOKA, Hiroyuki⁶; NODA, Hibiki⁷; SAIGUSA, Nobuko⁷; IDE, Reiko⁷; TAKAHASHI, Yoshiyuki⁷

¹Research Faculty of Agriculture, Hokkaido University, ²School of Agriculture, Hokkaido University, ³Faculty of Life and Environmental Sciences, University of Tsukuba, ⁴Japan Agency for Marine-Earth Science and Technology, ⁵National Institute for Agro-Environmental Sciences, ⁶River Basin Research Center, Gifu University, ⁷National Institute for Environmental Studies

Terrestrial ecosystems, forest, grassland and so on, absorb atmospheric CO₂ as a greenhouse gas by photosynthesis, and are thought to mitigate global warming. Estimation of geographical extent of their photosynthetic activity is very crucial for the understanding of global climate change in future. However, conventional vegetation indices (for ex, NDVI, EVI, etc.) representing the greenness of ecosystem, reduce the accuracy for photosynthesis estimation in the particular situations: for ex., the overestimation in evergreen forest in winter and in drought.

Chlorophyll fluorescence is emitted from chloroplast to release the overflowed energy of incident sunlight (so-called as Sun-Induced Fluorescence; SIF). Recently, several studies proved that SIF could be utilized for photosynthesis estimation at the ecosystem spatial scale (Zarco-Tejada et al., 2013, AFM, etc.) as shown by the strong correlation between SIF and gross primary production (GPP). On the other hand, the availability of SIF is reduced due to small number of ground-based measurement though highly evaluated potential of them.

This study compiles the SIF derived at five different ecosystems at tower-based flux stations in Japan: paddy field in Mase, grassland in Tsukuba university, deciduous broad-leaf and evergreen needleleaf forests in Takayama, deciduous needleleaf forest in Fujihokuroku. The SIF is calculated in the O₂-A band around the wavelength of 760 nm by Fraunhofer line depth (FLD) method. We will compare the SIF to eddy GPP flux during 2005-2013 and show the preliminary analysis on the availability of the SIF for the estimation of ecosystem photosynthesis.

Keywords: Ecosystem Photosynthesis, Remote Sensing, Flux measurement, Satellite measurement

Estimation of water storage and evaporation at seasonal wetlands in Namibia by satellite remote sensing

MIZUOCHI, Hiroki^{1*} ; NASAHARA, Kenlo²

¹University of Tsukuba, ²University of Tsukuba

North-central Namibia, classified as a semi-arid region, forms vast seasonal wetlands during the rainy season. Especially the seasonal ponds near the local farmers' house (*ondombe*) are assumed as important water resources for agriculture. Local farmers mainly have cultivated only pearl millet (*Pennisetum glaucum*), but nowadays they require the introduction of rice cropping in the seasonal ponds to mitigate the risk of poor harvests.

To introduce sustainable rice cropping, evaluation of available water resource and water budget in the seasonal ponds is required. However, there has been few such researches due to the difficulty of the hydrological monitoring because the seasonal ponds are distributed widespread area, and the water storage in the seasonal ponds fluctuate even during a few days. Thus we utilized satellite remote sensing together with field measurement to estimate the chronological water storage and evaporation in the seasonal ponds.

Study site covers north-central Namibia and the part of southern Angola (16:29:43S-19:05:20S, 14:24:59E-17:00:53). Also we set three test sites (each site is 5.4 km by 5.4 km) in the study site for field measurement. First we prepared three different types of satellite data (AMSR-E and AMSR2, MODIS, Landsat ETM+) from 2002 to 2013. We integrated these data by new data fusion technique (database unmixing; Mizuochi *et al.*, 2014, *Remote Sensing*) to detect water covered area with temporally and spatially high resolution. Second, we made a field laser measurement in 11 typical seasonal ponds to generate regression model between water covered area and water storage in the seasonal ponds. Then we translated water covered area detected by satellite remote sensing into water storage based on the regression model. Finally, we estimated evaporation from seasonal ponds derived from satellite data, comparing with precipitation data obtained by field observation to reveal the hydrological features in this region.

As the result of database unmixing, the spatial resolution was improved from original resolution of AMSR-E and AMSR2 (25 km) to Landsat ETM+ resolution (30 m). Temporal resolution was also dramatically improved: original Landsat ETM+ provided only 1.8% available data through the study period, whereas database unmixing provided 88.5% available Landsat-like data through the study period. The laser measurement provided 110 samples which shows the relationship between water covered area and water storage in seasonal ponds. Based on these samples we generated regression model, and eventually translated water covered area detected by database unmixing into chronological water storage data from 2002 to 2013. Both the integral water storage and maximum value of water storage for each year reflected the precipitation situation of each year. For example, in the rainy season between 2012 and 2013, which was reported as the severe drought once in 30 years, the water storage was significantly less than the other year's water storage. While in the rainy season between 2008 and 2009, during which the flooding occurred in the surrounding rivers, both the annual integral water storage and maximum water storage were high. By comparing evaporation estimated by the satellite data with the precipitation data in three test sites, it was shown that the evaporation from the seasonal ponds was several to ten percent of the amount of water input by the precipitation. This result shows that both water infiltration into the ground and evaporation from the soil surface are important process to express this region's water budget.

This study not only revealed this region's water storage and water evaporation but also proposed new approach for hydrological monitoring with high spatiotemporal resolution in seasonal wetlands. Further researches should be done in the future on improvement of the algorithm or validation of these result.

Keywords: semi-arid, seasonal pond, database unmixing, laser measurement

Process-based modeling for mountain pasture dynamics in unusual warm and long snow-free wintertime

KATATA, Genki^{1*} ; GROTE, Ruediger² ; ZEEMAN, Matthias² ; MAUDER, Matthias² ; OTA, Masakazu³ ;
LU, Haiyan² ; KIESE, Ralf²

¹Karlsruhe Institute of Technology (KIT-IMK/IFU)/Japan Atomic Energy Agency, ²Karlsruhe Institute of Technology (KIT-IMK/IFU), ³Japan Atomic Energy Agency

Alpine plant ecosystems are known as a carbon source mainly in colder periods when soil respiration is exceeding gross primary productivity. However, there is still little information about the specific impact of snow cover and thus the response of these ecosystems to climate warming and accompanying snow cover shortening is unclear. Here, we coupled a multi-layer atmosphere-SOIL-VEGetation model (SOLVEG) with a detail snow scheme, a grass growth scheme, and a soil microbiology scheme to investigate managed grassland-snowcover dynamics. The employed grass growth module can simulate key processes under cold environment such as leaf formation, elongation and death, tillering, carbon allocation, and acclimation on temperature on photosynthetic activity and frost damages. We applied the modified SOLVEG to pre-alpine grassland sites in TERrestrial ENVironmental Observatories (TERENO) networks in Germany for a year with an exceptional small amount of snowfall (2013-2014). The modified model reproduced temporal changes in observations of surface energy and CO₂ fluxes, soil temperature and moisture, and aboveground biomass. Our simulations and measurements demonstrate that grasses at lower elevation continuously assimilate atmospheric CO₂ even in the middle of winter season. On the other hand, dead leaf biomass increases due to frosts over cold snow-free days. As a result, snow-free wintertime carbon uptake was not as large as expected (13% of the annual carbon uptake), and it almost balanced with wintertime soil respiration. However, under temperature rise conditions, grass ecosystems act as a strong sink of CO₂ from winter to early spring (25% of the annual carbon uptake) due to a decrease of frost damages of foliage. Future climate developments may enhance the importance of wintertime carbon uptake of typical mountain grass species in the world.

Keywords: managed grassland dynamics, snow-free period, land surface model, photosynthesis, frost damage, European Alps

Model-Aided Analysis of FACE Effects on Rice Canopy Photosynthesis, Transpiration, and Water Use Efficiency

IKAWA, Hiroki^{2*} ; OUE, Hiroki¹ ; YOSHIMOTO, Mayumi² ; KOBAYASHI, Kazuhiko³ ; OKADA, Masumi⁴

¹Faculty of Agriculture, Ehime University, ²National Institute for Agro-Environmental Sciences, ³Graduate School of Agricultural and Life Sciences, The University of Tokyo, ⁴National Agricultural Research Center for Tohoku Region, ⁵Faculty of Agriculture, Iwate University

Sustainable agricultural practice requires promising crop productivity with efficient water use. Given the projected increase in atmospheric CO₂ concentration [CO₂], our understanding on the CO₂ effects on rice productivity (i.e., photosynthesis) and water use (i.e., transpiration) on a leaf scale improved in the last few decades, particularly with Free-Air CO₂ Enrichment (FACE) experiments that enable a simulation of a future agricultural field with high [CO₂]. However, very few information is yet available as to how the investigation on a leaf-level response of photosynthesis and transpiration to [CO₂] is translated to the whole canopy photosynthesis and transpiration (Shimono et al., *Glob. Change Biol.*, 2013; Yoshimoto et al., *Agric. For. Meteorol.*, 2005). This is partly due to the limited size of a FACE ring where it is difficult to apply top-down measurements, such as the eddy covariance technique. In this study, we quantified the effect of [CO₂] on rice canopy photosynthesis, transpiration and water use efficiency, using a multi-layer model with the model parameters obtained from single-leaf photosynthesis and transpiration measurements. With the model parameters carefully determined, we delineated the effects of [CO₂] on the canopy photosynthesis and transpiration through the changes in physiological and micrometeorological conditions for a better understanding on future rice productivity and water use.

Improving the soil sub-model of the process-based terrestrial ecosystem model to apply tropical swamp forests

HIRATA, Ryuichi^{1*} ; ITO, Akihiko¹ ; HIRANO, Takashi² ; SAIGUSA, Nobuko¹

¹National Institute for Environmental Studies, ²Research Faculty of Agriculture, Hokkaido University

We applied conventional model (Vegetation Integrative Simulator for Trace Gases; VISIT) to tropical peat forest data obtained from Palangkaraya (PDF) site in Kalimantan but it cannot simulate carbon balance accurately. Especially, ecosystem respiration (RE) showed opposite seasonal variation with observed RE because of misunderstanding of soil respiration process. Soil respiration process in conventional ecosystem model is controlled by only soil temperature and soil water content. In contrast, CO₂ release from peat is regulated by not only these components but also water table. Soil respiration data obtained from PDF site shows that the response of soil respiration to wetness is also different from that in conventional model. In order to improve the response of soil respiration, we modify soil submodel of the model. Ground water level is simulated by tank model. We modify soil respiration function using the relationship between ground water level and soil respiration.

Keywords: carbon balance, ground water level, Central Kalimantan

Global estimation of soil nitrous oxide emission using a semi-empirical model


HASHIMOTO, Shoji^{1*}

¹Forestry and Forest Products Research Institute

Nitrous oxide (N_2O) flux is one of the major greenhouse gas fluxes from terrestrial ecosystems. N_2O is generally released from soil surface to the atmosphere. In this study, a semi-empirical model was developed through modifying a model for soil CO_2 flux (Raich et al. 2002), and the global distribution of N_2O flux from soil was examined. The model consists of the functions of nitrogen, air temperature, and precipitation and the parameters of the functions were determined using a global dataset of N_2O emission (Stehfest and Bouwman 2006). The model was applied at a spatial resolution of 5-minute and at a monthly time resolution.

The preliminary calculation revealed that the total amount of N_2O emission was 14 Tg N yr^{-1} , and the monthly flux showed a clear seasonality, and was highest in August and lowest in February. The dry natural land is the major source of N_2O emission while the dry cultivated land was the second major source. Latitudinally, the flux was high around 30°N — 40°N and 10°S — 10°N . These results were still based on limited data, especially for wet ecosystems, and will be updated in future.

Keywords: soil, nitrous oxide, model, nitrogen



Variability in aboveground woody biomass in miombo woodland under fire disturbances

SAITO, Makoto^{1*} ; LUYSSAERT, Sebastiaan² ; POULTER, Benjamin³ ; WILLIAMS, Mathew⁴ ; CIAIS, Philippe² ;
BELLASSEN, Valentin⁵ ; RYAN, Casey⁴ ; YUE, Chao² ; CADULE, Patricia² ; PEYLIN, Philippe²

¹NIES, ²LSCE, ³Montana State University, ⁴University of Edinburgh, ⁵INRA Dijon

Miombo woodland is a savanna that spreads extensively over Africa, but it subjects to frequent fires (fire return intervals of 1-3 years) that are probably caused by human activities, such as cultivation, deforestation, and fuel wood collection. These activities result in degradation of miombo woodland. The degradation of large areas of miombo woodland involves the risks of rapid change in the current stable state of ecosystems to an alternative state and of increasing fire frequency and intensity. This study combined a process-based ecosystem model ORCHIDEE-FM with a fire regime model SPITFIRE to quantify the relative importance of fire regime and climate change in regulating aboveground woody biomass in miombo woodland under current and near-future climate conditions. The model was developed based on observations of resprouting and tree topkill rates in individual tree size classes for varying fire intensity. The model demonstrated that fire has a large impact on aboveground woody biomass and vegetation structure in miombo woodland. Aboveground woody biomass in miombo woodland significantly varies with fire regime with fire intensity and fire return interval. A shorter fire return interval and higher fire intensity results in a greater reduction of aboveground woody biomass by reducing the mean tree size. Although fire return interval in miombo woodland depends on human activities, fire intensity depends on the amount of grass fuel and on the season. For the current fire regime in miombo woodland, adaptive fire management is necessary to maintain today's aboveground woody biomass by controlling moderate fire intensity and frequency under current climate conditions. Under near-future climate with elevated CO₂ concentration and warmer climate, miombo woodland is perhaps more tolerant of fire disturbances by increase in the resprouting capacity and growth rate of woody plants.

Keywords: savanna, miombo, fire, aboveground woody biomass

Arctic terrestrial model intercomparison and its site-specific difference - physical process and biogeochemical cycle.

MORI, Junko^{1*} ; SAITO, Kazuyuki² ; MIYAZAKI, Shin¹ ; YAMAZAKI, Takeshi³ ; ISE, Takeshi⁴ ;
ARAKIDA, Hazuki⁵ ; HAJIMA, Tomohiro² ; IJIMA, Yoshihiro² ; MACHIYA, Hirokazu¹ ; SUEYOSHI, Tetsuo¹ ;
YABUKI, Hironori² ; GTMIP, Group⁶

¹National Institute of Polar Research, ²Japan Agency for Marine-Earth Science and Technology, ³Tohoku University, ⁴Kyoto University, ⁵RIKEN, ⁶GTMIP group

GRENE-TEA Model Intercomparison Project (GTMIP) is one of the modeling group's activities in the terrestrial research project of the GRENE Arctic Climate Change Research Project (GRENE-TEA). One of the challenges for Arctic model development is scarcity of site observation data available for validation of the models. GTMIP stage 1, site simulations at GRENE-TEA observation sites using the common driving data which was elaborately combining reanalysis and site observation data, assesses the inter-model variations and the site-specific differences, investigates their attributions to the implemented processes and their complexity of the models, and propose possible improvement of physical and biogeochemical processes for modeling of the Arctic terrestrial (excl. glaciers and ice sheets).

So far, 16 terrestrial process models participate in GTMIP stage 1: a permafrost model (FROST), physical snow models (SMAP and SNOWPACK), land surface models (2LM, HAL, JULES, MATSIRO with several versions, and SPAC-Multilayer), physical and biogeochemical soil dynamics model (PB-SDM), terrestrial biogeochemical models (BEAMS, Biome-BGC, STEM1 and VISIT), dynamic global vegetation models (LPJ and SEIB-DGVM coupled with a land surface model (Noah-LSM) or standalone), and a coupled hydrological and biogeochemical model (CHANGE). Driving and validation data were produced for the four GRENE-TEA observation sites; Fairbanks (USA), Kevo (Finland), Tiksi (Russia) and Yakutsk (Russia), which have different characteristics in snow accumulation, permafrost condition, vegetation and continentality. The 30plus-year forcing data is primarily prepared from reanalysis data to compensate missing values and limitation in the coverage period of the observation data. First, we made 30 minutes-interval forcing dataset (level 0.2) based on ERA-Interim. Air temperature and precipitation are corrected with CRU and GPCP respectively. Then, the site specific data (level 1.0) was created by fitting the level 0.2 data to the observation data at a site to inherit the characteristics of the site with enough temporal coverage for driving models. Target period was set from 1980 to 2013 (34 years). Model outputs are compared and evaluated in terms of the metrics in the five categories: energy budget, snowpack (annual maximum snow depth, snowpack duration etc.), phenology, subsurface thermal and hydrological condition (ground temperature profile, active layer/seasonal freezing thickness etc.) and carbon budget.

Physical models reproduced mean annual latent heat flux better, especially at Fairbanks and Yakutsk, than biogeochemical models that tend to show larger values than the observed, and great inter-model variability at Kevo. While dedicated snow models are good at reproducing maximum snow depth except for Tiksi, those models with invariant snow density show relatively lower value than the observation. At Tiksi, a tundra site on an Arctic coast, all the models overestimate snow depth. Annual gross primary production is reproduced well at Yakutsk. However at Fairbanks, the inter-model variation is large. Annual net ecosystem production (NEP) does not have such large inter-model variation at all sites examined. However, at Fairbanks, model outputs relatively overestimate to the observation value. On the contrary, NEP is underestimated at Yakutsk. Causes and attributions of the variability in reproducibility of the models will be discussed further.

Keywords: Circum-Arctic region, Terrestrial process model

A design for the new Earth System Model with terrestrial carbon/nitrogen cycle processes

HAJIMA, Tomohiro^{1*} ; WATANABE, Shingo¹ ; KAWAMIYA, Michio¹ ; WATANABE, Michio¹ ; NOGUCHI, Maki¹ ; OKAJIMA, Hideki¹ ; TATEBE, Hiroaki¹ ; ITO, Akinori¹ ; YAMAZAKI, Dai¹ ; TACHIIRI, Kaoru¹ ; KATO, Etsushi² ; SUGIYAMA, Masahiro³ ; ITO, Akihiko⁴ ; OHGAITO, Rumi¹

¹Japan Agency for Marine-Earth Science and Technology, ²The Institute of Applied Energy, ³UTokyo Policy Alternatives Research Institute, ⁴National Institute for Environmental Studies

Changes in the natural environment that are the result of human activities are becoming evident, and these changes are interrelated and can not be investigated without interdisciplinary collaboration between scientific fields. In order to understand the interrelated mechanism within the Earth's environment and to evaluate/project the anthropogenic impact on it, Earth system models (ESMs) have been developed by incorporating the biogeochemical processes of terrestrial/marine ecosystems into atmosphere-ocean general circulation models. These models have joined the Coupled Model Intercomparison Project (CMIP) and contributed to the assessment report of IPCC. We finished a process of reviewing our modeling and scientific activities up to the IPCC 5th assessment report, and we have started to design a new generation of ESMs. In this presentation, we briefly summarize the scientific/modeling tasks with ESM, and illustrate the design of the new ESM we are now developing.

Keywords: terrestrial ecosystems, carbon cycle, nitrogen cycle, Earth system models, Climate change projection, Biogeochemistry

Estimation of carbon cycle in a cool-temperate coniferous forest by continuous monitoring of carbon isotopic ratio

TAKANASHI, Satoru^{1*} ; DANNOURA, Masako² ; NAKANO, Takashi³ ; KOMINAMI, Yuji¹ ; MIYAMA, Takafumi¹

¹Forestry and Forest Products Research Institute, ²Kyoto University, ³Mount Fuji Research Institute, Yamanashi

In recent years, carbon assimilation has been estimated at various vegetation and forests are expected as carbon reservoir for the future climate changes. In a forest ecosystem, gas-exchange processes are complicated and difficult to estimate carbon flows because leaf, trunk, root and soil have distinctive gas-exchange characteristics and are interactively affected by environmental factors within a canopy. Fluxes and concentrations of ¹²CO₂ isotope compositions provide information about ecosystem physiological processes and their response to environmental variation. We continuously measured the concentrations of ¹²CO₂ and ¹³CO₂ inside the automated closed dynamic soil chambers and the surrounding atmosphere of a cool-temperate coniferous forest near Mt. Fuji in Japan (AsiaFlux site code:FJY) using a tunable diode laser spectrometer (TDLS: G2101i, Picarro Inc.). The $\delta^{13}\text{C}$ of the atmospheric CO₂ were fluctuated according the changes of CO₂ concentration and fluctuations of the carbon isotope signal from soil respired CO₂ ($\delta^{13}\text{C}_{\text{R}_s}$) was not detected, although Keeling plots to estimate $\delta^{13}\text{C}_{\text{R}_s}$ were very sensitive and the results were very scattered.

To understand carbon flows in a tree, we carried out in situ ¹³C pulse-labelling experiments for a mature pine tree (*Pinus densiflora*, Tree height: 20m) at the FJY site. The experiments were carried out September 2012, December 2012, and July 2013, covering the canopy of the tree by a plastic film chamber and introducing ¹³CO₂ into the labelling chamber. The internal ¹²CO₂ and ¹³CO₂ was monitored by the TDLS to calculate a carbon assimilation by the tree during the labelling experiments. Carbon efflux from a trunk surface were monitored by the TDLS using four closed dynamic trunk chambers installed in different heights (15.5, 11.1, 7.3 and 3.8m height) of the tree. The carbon flow speed were estimated from the arriving time of ¹³CO₂ pulse to the trunk chambers. The pattern of labelled ¹³CO₂ efflux in winter was different from other seasons. The speeds ranged from 0.04 to 0.24 m/hr and relatively slow in winter. The amount of carbon respired from the trunk surface were ranged 14-20% of the assimilated carbon. In winter, the tree respired carbon from relatively upper trunk surface and lower in other seasons.

Keywords: forest, carbon cycle, isotope, laser spectrometry, carbon dioxide, respiration

Methane dynamics in a temperate forest revealed by plot-scale and ecosystem-scale flux measurements

SAKABE, Ayaka^{1*}; KOSUGI, Yoshiko¹; TAKAHASHI, Kenshi²; ITOH, Masayuki³; UHEYAMA, Masahito⁴; IWATA, Hiroki⁵; ATAKA, Mioko¹

¹Graduate School of Agriculture, Kyoto University, ²Research Institute for Sustainable Humanosphere, Kyoto University, ³Center for Southeast Asian Studies, Kyoto University, ⁴Graduate School of Life and Environmental Sciences, Osaka Prefecture University, ⁵Faculty of Science, Shinshu University

Soils play important roles as CH₄ sources and sinks. CH₄ is produced in anoxic environments, including submerged soils, by methanogenic bacteria. On the contrary, CH₄ is oxidized by methanotrophic bacteria in upland soils. In general, forest soils are recognized as the efficient sinks for atmospheric CH₄, because of their CH₄ oxidation capacity in water-unsaturated soil (Le Mer and Roger, 2001). However, we hypothesized that forest ecosystems, especially in wet warm climates such as Asian monsoon climate, are not always CH₄ sink. In this study, we examined the CH₄ dynamics in a temperate Asian monsoon forest (Kiryu Experimental Watershed: 35°N, 136°E), which included wet areas along riparian zones within the watershed. In order to reveal the spatio-temporal variations of CH₄ fluxes, we combined multi-point plot-scale CH₄ flux measurements using chamber methods and ecosystem-scale CH₄ flux measurements using a micrometeorological method, relaxed eddy accumulation (REA) method (Businger and Oncley, 1990; Hamotani et al., 1996, 2001).

Intensive manual chamber measurements of CH₄ fluxes at 60 points in the wet areas and within the water-unsaturated forest floor, respectively, showed that the wet areas had a greater spatial and temporal variability of CH₄ fluxes than the forest floor. This indicates that accurate consideration of CH₄ fluxes from any wet areas is important in order to evaluate the CH₄ budget within the forests. From biweekly continuous manual chamber measurements of CH₄ fluxes at 9 points in the wet areas and the forest floor, respectively, hotspots of CH₄ emissions were observed during summer and fall immediately after intensive precipitation in the wet areas. On the other hand, in the forest floor, seasonal variations of CH₄ fluxes were not simply associated with temperature variations. In contrast, CH₄ absorption increased at some measurement plots in spring before intensive summer rainfall. In addition to the manual chamber measurements, we observed the environmental responses of CH₄ fluxes at a half-hourly time resolution, by using automated chamber measurements at three plots on the water-unsaturated forest floor. We found that the CH₄ absorption flux was greatly weakened by summer intensive rainfall, but recovered and peaked after rainfall as the soil water content decreased. The responses of CH₄ fluxes to rainfall were different for each plot. In a dry soil plot with a thick humus layer, CH₄ fluxes decreased abruptly at the peak of rainfall intensity, and it increased gradually after rainfall. In a wet soil plot and a dry soil plot with a thinner humus layer, such abrupt decreases in CH₄ fluxes were not observed, and CH₄ fluxes gradually switched from a sink to neutral following rainfall. Simultaneous measurements of CO₂ fluxes provided useful information when considering the controlling factors affecting complex CH₄ fluxes in terms of gas diffusivity and microbial activity.

The ecosystem-scale CH₄ flux measurements revealed that the Japanese cypress forest switched seasonally between being a sink and source of CH₄, and the pattern differed year by year. CH₄ fluxes tended to be a source during summer and fall, and switched to a sink during dry period. At hourly to daily timescales, the CH₄ fluxes were sensitive to rainfall; rain events increased CH₄ emission, decreased CH₄ absorption, or shifted CH₄ absorption to CH₄ emission. The results show that the temperate forest containing riparian zone acted as a CH₄ source seasonally, through the increased CH₄ emission in the wet areas and/or the decreased CH₄ absorption on the water-unsaturated forest floor in response to changing soil temperatures and/or the soil water status. The Asian monsoon rainfall was found to strongly influence temporal variations in CH₄ fluxes at both plot-scale and ecosystem-scales.

Response of Siberian larch in Mongolia to environmental variability for the last century

KITAYAMA, Asami^{1*} ; SUGIMOTO, Atsuko² ; YONENOBU, Hitoshi³ ; MIJIDSUREN, Byambasuren⁴ ;
LOPEZ, Larry⁵

¹Graduate school of environmental Science, Hokkaido University, ²Faculty of Earth Environmental Science, Hokkaido University, ³College of Education, Naruto University of Education, ⁴Plant Protection Research Institute, Mongolian state university of Agriculture, ⁵Faculty of Agriculture, Yamagata University

Mongolian forest locates the southern boundary of Siberian Taiga forest. Patchy forests are observed only on the permafrost in moisture area in Mongolia where annual precipitation exceeds about 300mm. Therefore, this means moisture condition is a major factor controlling forest distribution. Recently, increase in temperature and decrease in precipitation have been reported in large area of Mongolia (Batima, 2006), and that these environmental changes may be a potential trigger tree mortality, change in forest distribution and future thaw of permafrost. In this study, we selected five areas showing different moisture condition and growth environment, Terelj (47N, 107E) in Khenty mountain region, Tsagaannuur (51N, 99E) and Hatgal (50N, 100E) near Khuvsgul Lake in the northern forest area, and Tariat (48N, 100E) and Uyanga (51N, 102E) in Hangayn mountain region respectively. Tree-ring cores were collected from Siberian larch and tree-ring width was measured for each areas. Carbon isotope ratio in tree-rings were analyzed to understand tree response to environmental variability for two areas of these.

Tree-ring width at Terelj negatively correlated with temperature and positively correlated with precipitation, and carbon isotope ratio positively correlated with temperature and negatively correlated with precipitation of previous and current summer. Especially, tree-ring width shows more clear correlation with precipitation in previous summer than that in current summer, whereas carbon isotope ratio shows more good correlation with precipitation in current summer than that in previous summer. These results suggest that stem growth depends on the moisture condition in previous year, while tree-ring is formed with carbon assimilated in the current year. Siberian larch showed very narrow ring width in recent years (after 1997) with very high carbon isotope ratio at Terelj, suggesting severe drought stress. On the other hand, tree-ring width and carbon isotope ratio in northern forest area did not show clear correlation with climate parameters. This results imply that trees in Tsagaannuur and Hatgal grow under relatively wet condition compared to trend at Terelj. Narrow ring width and high carbon isotope ratio found for Terelj was not observed trees in northern forest area.

Keywords: Monoglia, Larch, Tree-ring, Carbon isotope ratio, Drought stress

In situ measurement of soil carbon monoxide flux in temperate forest

KAWAI, Amy^{1*} ; TAKAHASHI, Kenshi² ; SAKABE, Ayaka³ ; KOSUGI, Yoshiko³

¹Faculty of Agriculture, Kyoto University, ²Research Institute for Sustainable Humanosphere, Kyoto University, ³Graduate School of Agriculture, Kyoto University

Carbon monoxide plays an important role in the tropospheric chemistry via its reaction with the OH radical. Soil uptake of CO is a sink of secondary importance, except for the chemical reaction with the OH. However, no reports are available on soil CO flux continuously measured in the field of forest. Therefore, it is still unknown about the dynamics of CO in the temperate forest ecosystem, especially the response to the micrometeorological and soil environmental change. In this study, we intended to clarify the effects of the differences of the soil environment by measuring soil CO flux with the closed-chamber method and a laser spectrometer.

Soil CO flux was observed in Kiryu hydrological study site (KEW), in Shiga Prefecture, Japan. KEW has 55-year-old cypress forests. Priority species of the forest floor is *Eurya japonica* growing sparsely. CO fluxes between soil surface and the atmosphere were measured using the closed chambers, in which changes in CO concentration during the chamber closure were monitored in-situ with the laser spectrometer. Three chambers were set up in three locations under different soil environment. Also, the soil moisture content and soil temperature in the three plots were measured.

We found a significant absorption of atmospheric CO by soil surface. Focusing on the differences with the plot, less absorption was found in the plot of less humus layer. More CO absorption was observed when the CO concentration of surrounding area was high.

Application of three sap flow techniques for Japanese cedar: attempting to estimate the characteristics of sap movement

IIDA, Shin'ichi^{1*} ; TAKEUCHI, Shin'ichi² ; SHIMIZU, Takanori¹ ; TAMAI, Koji¹

¹Forest Hydrology Laboratory, Forestry and Forest Products Research Institute, ²Minami Kyushu University

Forests cover about 66% of land area of Japan, and Japanese cedar (*Cryptomeria japonica*) occupies 18% of total forested area. Recently, the water balance of a stand of Japanese cedar has been made clear quantitatively by using eddy-covariance method and sap flow technique in Kyushu Island, Japan (Kumagai et al., 2014; Shimizu et al., 2015). Although Japanese cedar is one of the most representative planted species in Japan, few studies have been carried out except in Kyushu Island. The most common technique of sap flow is thermal dissipation method (TDM, Granier, 1985) in Japan. Heat ratio method (HRM, Burgess et al., 2001) and heat field deformation method (HFD, Nadezhdina et al., 2012) have been widely used among foreign countries, and advantages and disadvantages of these techniques were described (e.g., Vandegehuchte and Steppe, 2013). However, HRM and HFD have not been applied for Japanese cedar in Japan. In this study, we applied TDM, HRM and HFD for a tree of Japanese cedar, and carried out preliminary comparison among three techniques. Based on field measurements, we attempted to estimate the sap flow characteristics within the target tree by three techniques.

We conducted measurements in a mature stand of Japanese cedar, whose age is 62, within Tsukuba Experimental Watershed located in southern part of Mt. Tsukuba, Japan. We picked up a tree of Japanese cedar whose height is 24.9 m and diameter at breast height is 40.4 cm, and applied sensors of TDP, HRM and HFD. We used handmade sensors for TDM (e.g., Iida et al., 2013) and sensors for HRM and HFD manufactured by ICT international Pty Ltd (type SFM1 and HFD8, respectively). Japanese cedar has white zone, in which water movement stops, near the sapwood area. We injected acid fuchsin into a stem, and determined colored area as sapwood. The length of TDM sensor was 20 mm, and sap flux density, which is mean value along the sensor length, was computed by the calibration equation by Granier (1985). The width of sapwood was 44 mm, and additional TDM sensor was inserted into the sapwood at the depths from 20 to 40 mm. On the other hand, the length of HRM sensor was 35 mm, and the sap flow movement was detected at the depths of 12.5 and 27.5 mm. For HFD, the sensor length was 96 mm, and the depths of sap flow detecting were 20, 30, 40, 50, 60, 70, 80 and 90 mm.

The sap flux densities measured at the depth from 20 to 40 mm with TDM was larger than that at the depth from 0 to 20 mm. Similar trend was found in the outputs of HRM and HFD. The results of measurements by three techniques show that active sap flow movement occurred up to the depth of 40 mm of sapwood. The diurnal variations in sap flow movement measured by three techniques were almost same, suggesting the applicability of these techniques to evaluate the characteristics of sap flow for a Japanese cedar. However, additional measurements including thermal diffusivity of wood are necessary to obtain sap flux density by HRM and HFD (e.g., Vandegehuchte and Steppe, 2013): in current stage, we have only qualitative comparisons among three techniques. Moreover, to compare the sap flux densities measured by TDM, HRM and HFD, we have to make clear the azimuthal changes in sap flux density within the target tree: three sensors inserted into different part of a stem in this study. We will present the results including these topics.

Cited paper

- Burgess, 2001. Tree Physiol., 21:589-598.
Granier, 1985. Ann. Sci. For., 42: 193-200. [English translation, in Evaporation, Benchmark Papers in Hydrology 2. Gash JHC, Shuttleworth, WJ (eds). IAHS Press: Oxfordshire; 61-63].
Iida et al., 2013. JARQ, 47: 319-327.
Kumagai et al., 2014. J. Hydrol., 508: 66-76.
Nadezhdina et al., 2012. Trees, 26:1439-1448.
Shimizu et al., 2015. J. Hydrol., 522: 250-264.
Vandegehuchte and Steppe, 2013. Function. Plant Biol., 40: 213-223.

Keywords: Thermal dissipation probe, Heat ratio method, Heat field deformation

Seasonal change in photosynthetic function of cedar and cypress forest estimated by tower-based chlorophyll fluorescence

TSUJIMOTO, Katsuto^{1*}; KATO, Tomomichi²; HIRANO, Takashi²; SAITOH, Taku M.³; NAGAI, Shin⁴; AKITSU, Tomoko⁵; NASAHARA, Kenlo N.⁵

¹Graduate School of Agriculture, Hokkaido University, ²Research Faculty of Agriculture, Hokkaido University, ³River Basin Research Center, Gifu University, ⁴Japan Agency for Marine-Earth Science and Technology, ⁵University of Tsukuba

Estimating the carbon cycle in terrestrial ecosystem is very important for predicting the climate change in future. Remote sensing observation with vegetation indices (VIs, for ex., Normalized Differential Vegetation Index: NDVI) has contributed greatly to estimate terrestrial carbon budget. Because conventional VIs represent generally the greenness of ecosystem, however, there has been the less accuracy especially for winter time in evergreen forest where the VIs tend to overestimate the photosynthesical activity due to its nearly constant greenness.

Chlorophyll fluorescence is emitted from the chlorophyll a and b to release the excess sun-light energy, and has been used for detecting the photosynthesis stress in numerous ecophysiological studies so far. Recently, the chlorophyll fluorescence has been utilized to represent the gross primary production (GPP), ecosystem photosynthesical activity, by the satellite remote sensing studies (e.g. Frankenberg *et al.*, 2011, GRL). However, small number of chlorophyll fluorescence observation on the ecosystem scale at the ground reduces its availability despite its highly expected usefulness.

The aim of this study is to clarify the relationships between chlorophyll fluorescence, and photosynthesis and light use efficiency (LUE) by the ground based measurement in coniferous evergreen forest. The observations were carried out in the plantation forest consisting of mature Japanese cedar and Japanese cypress in Takayama, Japan, from 2007 to 2008. Downward and upward spectral radiances were measured with hemispherical spectroradiometer (MS-700, Eko Instruments, Japan) mounted at 30m-high above the ground surface. We calculated the Sun-Induced fluorescence (F_S) around the O_2 -A band from the spectral data with the Fraunhofer Line Depth method. The GPP was calculated from the carbon fluxes measured with eddy covariance at the top of the tower (Nagai *et al.*, 2012). Both F_S and GPP were averaged for 30 minutes.

F_S showed the strong correlation to GPP linearly in the diurnal course (sunny day (2007/08/10): $r^2 = 0.81$, cloudy day (2007/07/24): $r^2 = 0.87$) and logarithmically in the seasonal change (2008 (half-hour): $r^2 = 0.68$, 2008 (daily): $r^2 = 0.87$). The GPP was fitted against the F_S for each month by the following rectangular hyperbolic curve:

$$GPP = \alpha GPP_{SAT} F_S / (GPP_{SAT} + \alpha F_S),$$

where α is the initial slope of the F_S -GPP curve and GPP_{SAT} is the saturated GPP at high F_S emission. These two parameters showed the clear seasonal change. The root-mean-square error (RMSE) calculated for 2008 was $4.56 \mu\text{molCO}_2 \text{ m}^{-2} \text{ s}^{-1}$. We also investigated the relationship between F_S and LUE in daily averages. The F_S -LUE relationship could be regressed by logarithm curve for each month ($r^2 = 0.46 \sim 0.95$). The seasonal changes in the regression coefficients for F_S -GPP and F_S -LUE curves were thought to be induced by the seasonal variation in the temperature-dependency of photosynthesis and the phenology.

We conclude that F_S can be utilized to estimate GPP and LUE in evergreen forest, and that relationship between F_S and GPP is influenced by environmental factors such as air temperature.

Keywords: Remote sensing, Light use efficiency, Eddy covariance, Carbon cycle

Year-to-year variability in the timing of start and end of growing season in deciduous forests in Japan

NAGAI, Shin^{1*} ; SAITOH, Taku²

¹Japan Agency for Marine-Earth Science and Technology, ²Gifu University

Accurate detection of the spatio-temporal variability in the timing start (SGS) and end of growing season (EGS) is important but challenging task to evaluate the ecosystem functioning and service in deciduous forests under meteorological and climate changes. Here, (1) we detected the spatio-temporal variability in the timing of SGS and EGS in Japan from 2001 to 2013 by analysing Terra/Aqua MODIS satellite-observed daily green-red vegetation index (GRVI) with a 500-m spatial resolution; (2) we examined the spatio-temporal variability in the timing of blooming of cherry, leaf-flushing of ginkgo, leaf-colouring of ginkgo and maple, and leaf-fall of ginkgo and maple based on dataset of *in situ* phenological observation; and (3) we evaluate the relationship between year-to-year variability in the timing of satellite-observed SGS and EGS and those of *in situ*-observed SGS and EGS. We found that (1) the year-to-year variability in the timing of SGS, EGS, blooming, leaf-flushing, -colouring, and -fall showed characteristics along vertical and/or horizontal gradients; and (2) the year-to-year variability in the timing of SGS correlated with those of blooming of cherry and leaf-flushing of ginkgo in Hokkaido and Tohoku.

Keywords: satellite data, biometeorological data, deciduous forest, Japan, phenology, year-to-year variability

Differences in inter-annual variation in flowering and budding dates at two different climatological sites in Japan

INOUE, Tomoharu^{1*} ; NAGAI, Shin¹

¹Japan Agency for Marine-Earth Science and Technology

To investigate the relationship between the inter-annual variability of the flowering and budding phenology and their geographical characteristics in Japan, we analysed the first flowering and budding dates of various plant species during winter and spring (Japanese camellia, persimmon, ginkgo, mulberry, narcissus, cherry, Japanese apricot, rhododendron, dandelion, Japanese wisteria, and Japanese lawn grass) from 1953 to 2011 at two different climatological study sites (Takayama, a northern colder site; Gifu, a southern warmer site). We found that (1) fewer than half of the species showed a trend of earlier phenology at both sites, (2) mean phenological dates at Gifu were distributed at wider range (DOY [day of year] 21.9 to 145.2) than those at Takayama (DOY 95.5 to 165.5), and (3) the species with earlier flowering or budding at Gifu showed higher variability and advanced phenology compared with plants that had later flowering and budding. These findings (a) suggest the possibility that flowering and budding phenology in central Japan has a localized response to increasing air temperature and (b) suggest the importance of the long-term and continuous observation of flowering and budding phenology for a range of plant species at multiple sites (Inoue and Nagai, *under review*).

Keywords: phenological observation, flowering date, budding date