

Tectonic features around the Abukuma ridge and their relationship with tectonic framework of the Northeastern Japan Arc

ARATO, Hiroyuki^{1*} ; CHIYONOBU, Shun¹

¹Faculty of International Resource Sciences, Akita University

The Abukuma ridge is a significant buried uplift zone bordering the east of southernmost part of the forearc basin extending from the central axial Hokkaido through offshore Shimokita and Kitakami to offshore Joban area in the Pacific side of northeast Japan.

Its tectonic characteristics and formation mechanisms might be decisive information to elucidate not only geotectonic framework of the Northeastern Japan Arc but spreading processes of the northern portion of Japan Sea.

In order to do so, following items must be clarified by the seismic dataset and well geological information at the Abukuma ridge and its adjacent area: i) structural geological characteristics of fault groups and folds, ii) their extending trend and direction, and iii) sedimentological and stratigraphic relationship of erosional or starved discontinuous surfaces.

Accordingly the authors are conducting re-interpretation of the following seismic surveys: i) approximately 2,000 square kilometers "Southern Abukuma ridge" three-dimensional seismic data acquired by METI in 2009, and ii) approximately 2,500 kilometers "Offshore Southern Sanriku and Kashima" two-dimensional seismic data acquired by MITI in 1986.

Three MITI wells, moreover, "Kesenuma-oki," "Soma-oki" and "Joban-oki," and several commercial exploratory wells had been drilled in the above mentioned waters. The "Kesenuma-oki" well, spudded in 1984, had drilled thick upper Cretaceous section and reached early Cretaceous granite basement at the total depth of 2,027.00 meter. The "Soma-oki" well, spudded in 1990, had drilled Neogene and Paleogene successions and reached upper Cretaceous at the total depth of 3,500.00 meter. The "Joban-oki" well, spudded in 1991, had drilled thick lower Miocene and upper Cretaceous overlain by relatively thin middle to late Miocene and Quaternary sediments, and confirmed Turonian at the total depth of 3,170.00 meter. Geological information from those wells are utilized as chronostratigraphic and lithostratigraphic controls for the seismic interpretation.

It became gradually clear as tentative results of the interpretation that:

i) the above mentioned forearc basin in the studied area may be subdivided into a few portions by several structural lineations characterized by strike-slip components, ii) the basin therefore may not have simple north-south elongated geometry, iii) such lineations are possibly continuous with some geologic tectonic lines or geotectonic discontinuities specified by subaerial geological descriptions in the northeast Japan, on the basis of their location and direction, and iv) several groups of a number of faults, identified at the southern part of Abukuma ridge in the 3D seismic dataset, may possibly be interpreted as a record for stress field directions of each geologic age.

Keywords: Abukuma ridge, offshore Miyagi Prefecture, offshore Fukushima Prefecture, Northeastern Japan Arc

Tephrochronology in the Cretaceous forearc basin, northern Japan

TAKASHIMA, Reishi^{1*}; KUWABARA, Sato²; NISHI, Hiroshi¹

¹The Center for Academic Resources and Archives, Tohoku University, ²Department of Earth Science, Graduate School of Science, Tohoku University

1. Introduction

Tephrochronology is useful method for basin analysis because tephra provide isochronous horizons across the different facies. Although numerous studies of tephra correlations have long been conducted on the Quaternary strata, there are much fewer attempts to correlate pre-Quaternary tephra because most of the pre-Quaternary tephra are not appropriate for applying conventional tephrochronological methods on account of alteration of glass. On the other hand, several studies based on the heavy mineral chemistry of tephra have been successful in identifying and correlating highly altered tephra (e.g., Sell and Samson, 2011a, b). Detailed tephra correlations by these studies solved various controversies on the Euro-American Ordovician biostratigraphy, biogeography and sequence stratigraphy. Consequently, prevailing tephrochronologic study of the pre-Quaternary strata appears to bring significant progress in the various field of earth sciences because it can provide a much higher precision of chronostratigraphic correlation than that of the radiometric dating in both local and regional scale.

The Yezo Group, exposed in Hokkaido, Japan, is represented by the mid-Cretaceous — Paleocene marine sequences which were deposited in a fore-arc basin along east of the active Asian continental margin. This group contains abundant macro- and microfossils as well as felsic tuffs. However, it has been very difficult to correlate between shallow and deep sea facies in detail because of difference in fossil fauna and sedimentary facies. In order to establish detailed depositional model in such old basin, we establish the Cretaceous tephrochronology in the Yezo Group based on the heavy mineral chemistry of phenocrysts within the tuffs.

2. Method

We collected 30 tuff samples from the whole horizons of the Yezo Group in Tomamae, Oyubari and Urakawa areas. The samples were separated into light and heavy fractions using sodium polytungstate. The separated heavy fraction was collected and handpicked under a binocular microscope to collect the apatite and biotite grains. 20 biotite phenocrysts per sample were analyzed major elements at the Department of Earth Science, Tohoku University using a JEOL 7330. 20 apatite phenocrysts per sample were analyzed major and minor elements at the Institute for Material Research, Tohoku University using a JEOL 8530F. Apatite analyses followed the method described in Gross et al. (2013).

3. Result and discussion

Apatite is one of the most common accessory mineral in the volcanic rocks, and is highly resistant to weathering, diagenetic alteration and diffusion processes. Furthermore, apatite shows a wide variety of trace-element compositions because the structure of apatite is highly tolerant of structural distortion and chemical substitutions. Therefore, chemical fingerprinting of apatite is ideal method to discriminate the altered tephra. Each tuff exhibits unique trend in Cl, Mg, Fe, and Mn concentrations in apatite, which demonstrates that apatite chemistry is useful for discriminating tuffs in the Yezo Group. Although biotite is one of major phenocrysts in the tuffs of the Yezo Group, it is less useful than apatite because of its weakness against diagenesis and weathering. However, Mg number and TiO₂ content of biotite are most effective discriminator of the tuffs in the Yezo Group, and each tuffs are distinguishable on the bivariate plot of the Mg number and TiO₂ contents.

Based on the above result, we correlated tuffs of the Yezo Group among the Tomamae area and the other areas in Hokkaido. As a result, we confirmed that at least 3 tuffs (tuffs at Lower Cenomanian, Turonian/Coniacian boundary and Santonian/Campanian boundary) can be traced widely (more than 100 km distance) throughout Hokkaido and across the different sedimentary facies.

Keywords: Cretaceous, Tephrochronology, Apatite, Yezo Group

Formation of forearc basins and its relationship to subduction zone dynamics

NODA, Atsushi^{1*}

¹Geological Survey of Japan, AIST

Forearc basins along subduction zones are important for studying sediment routing systems, submarine resources, and natural hazards. However, there are still many unresolved problems about them, for example, how do they evolve? or how do they interact with growth of accretionary prisms or intensity of tectonic erosion of the same age? Forearc basins were proposed as basins developed at landward side of accretionary prisms, whose depocenters migrated seaward as growth of the accretionary prisms [1] or landward as uplift of the outer arc highs [2]. However, there are some forearc basins without accretionary prisms along tectonic erosion-dominated subduction zones, which are controlled mainly by normal faults [3]. In addition, it was realized that sediment flux at the trench is important to control volume of outer wedge [4], suggesting it could also influence formation or style of evolution of forearc basins. In fact, subsidence curves of modern and ancient forearc basins are more variable than those of foreland, rift, or strike-slip basins [5].

In this study, I tried to understand formation of forearc basins by means of geological and geomorphological characters of 37 forearc basins with the outer wedges (frontal and middle prisms) in the world. I also measured width (W_{basin}) and sediment thickness (T_{basin}) of forearc basin, width (W_{wedge}) and slope angle (α) of outer wedge, slab dip angle under the wedge (β), orthogonal convergent rate of subducting plate (V_{orth}), thickness of trench fill sediments (T_{trench}).

As a result, forearc basins can be divided into 5 types (compressional/extensional accretionary, compressional/extensional strike-slip, and non-accretionary). A character of the compressional accretionary type is landward tilt of the basin due to uplift of outer arc high associated with backthrust or splay fault. Basin formation of this type may be related to self-similar growth or thickening of the outer wedge. In the extensional accretionary type, seaward migration of outer arc high by gravitational force, generating listric normal faults in the basins, can subside the basin. Internal or basal friction may be too weak to keep the taper angle of the wedge. The non-accretionary basins do not have conspicuous outer arc highs, resulting from tectonic erosion-induced subsidence. Strike-slip type shows intermediate characters between accretionary and non-accretionary types, but whose depocenters occasionally migrate parallel to the trench. Some basins show changes from non-accretionary type to accretionary one, and vice versa.

Width of wedge (W_{wedge}) has a positive correlation with T_{trench}/V_{orth} as a proxy of sediment flux at the trench. T_{basin} has a positive correlation to W_{basin} for the accretionary type, but is basically constant for the non-accretionary type. W_{basin}/T_{basin} of the accretionary type is almost constant regardless of W_{wedge} , but shows a negative correlation to W_{wedge} for the non-accretionary type. For all types of forearc basins, T_{basin} has a positive correlation to W_{wedge} or T_{trench} .

Variations of sediment flux at subduction zones can influence degree of growth/decay of outer wedges, and then, may switch the type of forearc basins. Changes from accretionary to non-accretionary type may cause subsidence of outer arc high because of tectonic erosion, which leads to remove accommodation space and erode the basin strata, and then produces a new basin with large W_{basin}/T_{basin} . On the other hand, changes from non-accretionary to accretionary type may uplift a part of the basin as an outer arc high, and generate a new basin at the landward side.

[1] Karig, D.E., 1974, *Ann Rev Earth Planet Sci* 2, 51–75; [2] Seely, D.R., 1979, *AAPG Memoir*, 29, 245–260; [3] von Huene, R. and Lallemand, S., 1990, *GSA Bull*, 102, 704–720; [4] Clift, P.D. and Vannucchi, P., 2004, *Rev Geophys*, 42, RG2001; [5] Xie, X. and Heller, P.L., 2009, *GSA Bull*, 121, 55–64.

Keywords: forearc basin, subduction zone, sediment flux, accretionary prism, sedimentary basin

Relationship between trends in Miocene basin development and outcrop-based depositional cycles and geological structures

NARISAWA, Sayaka^{1*} ; KURITA, Hiroshi²

¹Graduate School of Science and Technology, Niigata University, ²Faculty of Science, Niigata University

During the Late Oligocene-Middle Miocene, the Japanese archipelago was formed under control of extensional tectonics. At that time, a series of extensional basins developed in the back arc side of Honshu Island. The development of the Niigata basin, which is the largest basin in the coast of the Japan sea, has been argued in various ways. The Tsugawa-Aizu province (Shimadu, 1973) is important for the good understanding of the early stages of development of the Niigata basin. According to Toyoshima (2014), Miocene and later geological structures of Northeast Japan were formed by combination of multiple trends such as NNE-SSW (Niigata trend), NW-SE to WNW-ESE (transverse trend) and NNW-SSE to N-S (Tanagura trend). Relationship of the structural trends and basin development, especially such as field-scale sedimentary and structural processes, has not been considered in detail in the Niigata basin which includes the Tsugawa-Aizu province.

This outcrop-based study discusses basin development in Northeast Japan by sedimentary facies and structural analysis, considering cyclicity and basin geometry of the Mikawa and Tsugawa basins in the Tsugawa-Aizu province. The study area, the Tsugawa area in Aga, Niigata, has various scale basins which include the Mikawa basin in the west and the Tsugawa basin in the east. We estimate vertical changes of depositional environments by field work on Miocene deposits from the Tsugawa Formation to the Nomura Formation in the Yagiyama area, eastern part of the Tsugawa basin.

Comparison between the Mikawa basin and the Tsugawa basin show that the cyclicity of these basins is similar to each other. This study considers basin development also from fault trends by fault analysis. In general, geological structures in the Mikawa area show NNW-SSE to N-S trends, while those in the Yagiyama area have an obvious NW-SE trend.

Keywords: Miocene, back arc, Northeast Japan, sedimentary facies, fault

Crustal structure off western Noto Peninsula to the Yamato Basin observed by seismic reflection survey

NO, Tetsuo^{1*} ; SATO, Takeshi¹ ; KODAIRA, Shuichi¹ ; MIURA, Seiichi¹ ; ISHIYAMA, Tatsuya² ;
SATO, Hiroshi²

¹JAMSTEC, ²ERI

Since recent years, the source fault of the Japan Sea has been reviewed through several projects (e.g., Ministry of Land, Infrastructure, Transport and Tourism, 2014). As a part of these efforts, we conducted research in collaboration with the "Multidisciplinary research project for construction of fault model in the high strain rate zone" consisting of marine seismic surveys using a multichannel seismic system (MCS) and the ocean-bottom seismographs (OBSs) in the eastern part of the Japan Sea. As a result, these studies revealed the relationship between the distribution of crustal structure and seismic activity or shortening structure in the eastern part of the Japan Sea (e.g., No et al., 2014; Sato et al., 2014). However, observations to evaluate the potential damage of earthquakes and tsunamis had not been carried out fully west of Noto Peninsula and off western Hokkaido in the Japan Sea. Therefore, crustal structure data other than those of the eastern part of the Japan Sea are required.

In July and August 2014, we conducted a marine seismic survey to study the crustal structure from the region off western Noto Peninsula to the Yamato Basin. Some studies of crustal structure had been carried out in this survey area (Ludwig et al., 1975; Katao, 1988; Hirata et al., 1989). However, these studies were not able to obtain the spatial variations of the crustal structure and the relationship with active structure. Recently, seismic acquisition technology for deep and detailed seismic imaging has improved. By using this recent technology, we were able to obtain important data to study the relationship between the active structure and crustal structure in the vicinity of the survey area. Based on the results of these studies, although it is important to identify where the source faults have developed in the Japan Sea, structural factors are an important key for determining the development and size of the source faults. In addition, because there is ODP Leg 127 site 797 (Tamaki et al., 1990) directly beneath our seismic survey line, we contributed to the study on the formation of the Yamato Basin by examining the relation between the ODP results and our results.

We present an outline of the data acquisition and preliminary results of the data processing and interpretations from this study. Asymmetrical anticlines with reverse faults are well developed off the northern to western Noto Peninsula. The basement around the continental shelf and continental slope shows larger deformation. In the Yamato Basin, a clear reflector estimated to be the Moho can be identified in the vicinity of about 9 s (two-way time). The characteristics of the sedimentary layer are common within the survey area. For example, a strong coherent reflector that is estimated to be an opal-A/opal-CT BSR (bottom simulating reflector) (Kuramoto et al., 1992) was confirmed in all survey lines in the Yamato Basin. Furthermore, a coherent reflector across the entire crust was confirmed in some of the survey lines in the Yamato Basin.

Keywords: Japan Sea, Seismic survey, Yamato Basin, off Noto Peninsula, Yamato Rise, source fault

Structural analyses of sedimentary basins using gravity anomaly and gravity gradient tensor data

KUSUMOTO, Shigekazu^{1*}

¹Graduate School of Science and Engineering for Research (Science), University of Toyama

Seismic exploration and gravity surveys have often been employed to estimate subsurface structures, and gravity surveys have been used most often as rough surveys because they can cover a wide area in a short period with low cost. In addition, gravity surveys have other advantages regarding data interpretation; it is possible to interpret their map data intuitively and easily. For example, we could interpret low-gravity areas as thick sedimentary layers or low-density layers/materials with much accuracy. We can obtain detailed information on subsurface structures, such as density distribution and shape, by quantitative analyses. However, these analyses require a lot of time and prior information on subsurface structures (e.g., basic subsurface structure and density contrast) for solving problems.

In such a situation, semi-automatic interpretation methods, which are considered as intermediate methods between quantitative analyses and qualitative interpretations, can be used. In this method, structural boundaries are identified from gravity anomalies without constraint conditions on geology and geophysics. Types of high-pass filtering such as horizontal first derivative and vertical first derivative are well-known techniques.

In recent years, gravity-difference surveys have been carried out worldwide, and filtering and semi-automatic interpretation techniques using gravity gradient tensor have been developed and suggested. For example, Shape Index defined by xy components of gravity gradient tensor expressing curvature of gravity field (e.g., Cevallos, 2013), Dimensionality Index defined by combination of eigenvalues of the tensor (e.g., Pedersen and Rasmussen, 1990; Beiki and Pedersen, 2010), and a technique that estimates locations of causative source using eigenvectors of the tensor (e.g., Beiki and Pedersen, 2010; Beiki, 2013).

In general, it is necessary to obtain gravity gradient data with a gradiometer for performing analyses using gravity gradient tensor. However, these data can be derived by combining integration and differential of surface gravity anomalies in the Fourier domain (e.g., Mickus and Hinojosa, 2001).

In this presentation, the author will show derivations of all components of gravity gradient tensor from surface gravity-anomaly data using the Fourier method and will report results obtained by applying techniques using the tensor to structural analyses of structural basins.

[References]

Beiki, M., 2013, *Jour. Appl. Geophys.*, 90, 82-91. Beiki M., and Pedersen, L. B., 2010, *Geophysics*, 75, I37-I49. Cevallos, C., Kovac, P., and Lowe, S. J., 2013, *Geophysics*, 78, G81-G88. Mickus, K. L., and Hinojosa, J. H., 2001, *Jour. Appl. Geophys.*, 46, 159-174. Pedersen L. B., and Rasmussen, T. M., 1990, *Geophysics*, 55, 1558-1566.