

## アカスタ片麻岩体苦鉄質岩の Re-Os 同位体組成：冥王代玄武岩質地殻の解明にむけて Re-Os systematics on the mafic rocks in the Acasta gneiss complex: Implications for the Hadean basaltic crust

越田 湊子<sup>1\*</sup>; 石川 晃<sup>2</sup>; 鈴木 勝彦<sup>3</sup>; 小宮 剛<sup>2</sup>  
KOSHIDA, Keiko<sup>1\*</sup>; ISHIKAWA, Akira<sup>2</sup>; SUZUKI, Katsuhiko<sup>3</sup>; KOMIYA, Tsuyoshi<sup>2</sup>

<sup>1</sup> 東京大学理学系研究科地球惑星科学専攻, <sup>2</sup> 東京大学広域科学専攻 広域システム科学系, <sup>3</sup> 海洋研究開発機構  
<sup>1</sup>Department of Earth and Planetary Science, The University of Tokyo, <sup>2</sup>Department of Earth Science and Astronomy, Graduate School of Arts and Sciences, University of Tokyo, <sup>3</sup>Japan Agency for Marine-Earth Science and Technology

The first billion years of the Earth history is poorly understood because of scarcity of terrestrial rock records. Especially, the first 500 million years, named Hadean, is literally dark because no terrestrial rocks are preserved on the earth. To unravel the early history of the earth, it is important to study the oldest rocks rarely present in the Eoarchean terranes.

Acasta Gneiss Complex (AGC), located in the western part of the Slave Province, Canada, is one of the Eoarchean terranes, and mainly consists of 3.6-4.0 Ga felsic and layered gneiss suites with minor mafic rocks (e.g. Bowring et al., 1990; Izuka et al., 2007). The mafic rocks are distributed all over the AGC and occur as rounded to elliptical enclaves and inclusions within the felsic and layered gneisses. Although field occurrence of the mafic rocks suggests that they were formed before the formation of the precursors of the orthogneisses, their formation ages have not been determined exactly because of no magmatic zircons preserved in the mafic rocks (e.g. Mojzsis et al., 2014). Therefore, the whole-rock isochron dating is the most suitable to determine the magmatic age of the mafic rocks. However, the AGC is subjected to numerous metamorphism and alteration events so that the mafic rocks suffered from more or less secondary elemental movement (e.g. Moorbath et al., 1997; Sano et al., 1999). Therefore, it is necessary to find the primary signature and reconstruct the original compositions of the mafic rocks. Our previous study revealed that compositional variations of the mafic rocks were mainly formed due to the migmatization and identified a subset of the least altered samples. This study shows the whole-rock Re-Os isotope systematics of the least altered Acasta mafic rocks.

The twenty-seven, least altered samples were analyzed for whole-rock Re-Os isotopes. They have relatively large variations in  $^{187}\text{Os}/^{188}\text{Os}$  and  $^{187}\text{Re}/^{188}\text{Os}$  ratios ranging from 2.0 to 150 and from 28 to 2466, respectively. The measured samples display a roughly positive correlation on a  $^{187}\text{Os}/^{188}\text{Os}$  vs  $^{187}\text{Re}/^{188}\text{Os}$  diagram, corresponding to reference lines between 3.3 and 4.2 Ga. But, they are highly scattered (MSWD = 266), possibly due to post-magmatic Re loss or addition during thermal metamorphism because Re is, generally speaking, more mobile than Os (e.g. Reisberg et al., 2008). The nine samples with the highest Re contents display a correlation with a gentler slope, corresponding to an age of  $3453 \pm 20$  Ma (MSWD=16, initial  $^{187}\text{Os}/^{188}\text{Os} = -0.8 \pm 1.0$ ). The gentle slope and high Re contents suggest Re addition at or after  $3453 \pm 20$  Ma. In the former case, the age possibly indicates the metamorphic age, in agreement within errors with the whole-rock Sm-Nd isochron age of  $3371 \pm 141$  Ma (e.g. Moorbath et al., 1997; Antoine et al., 2014). In the latter case, the age may correspond to geological event; artificial.

On the other hand, eighteen samples, except for the samples with high Re concentrations, form two parallel lines with only small scattering. The line with high  $^{187}\text{Os}/^{188}\text{Os}$  ratios yields an age of  $4273 \pm 200$  Ma (MSWD=15, initial  $^{187}\text{Os}/^{188}\text{Os} = 1.83 \pm 0.74$ ), whereas the other yields an age of  $4081 \pm 320$  Ma (MSWD=11, initial  $^{187}\text{Os}/^{188}\text{Os} = 0.19 \pm 0.70$ ). Those ages are consistent with geological occurrence that the mafic rocks occur as enclaves within the 3.6-4.0 Ga felsic gneisses, suggesting that the Acasta mafic rocks were formed in the Hadean. In addition, their high initial  $^{187}\text{Os}/^{188}\text{Os}$  ratios relative to those of chondrite, 0.097 at 4273 Ma and 0.099 at 4081 Ma, suggest that the Acasta mafic rocks were formed from an enriched source.

キーワード: 苦鉄質岩, Re-Os 同位体, アカスタ  
Keywords: mafic rock, Re-Os isotopes, Acasta