

カナダ・ラブラドル、サグレック岩体の初期太古代 (>3.95 Ga) 堆積岩中の炭質物
と炭酸塩の炭素同位体比
Carbon isotope compositions of carbonaceous materials and carbonate from Saglek Block
(>3.96 Ga), Labrador, Canada

田代 貴志^{1*}; 石田 章純³; 堀 真子³; 伊規須 素子⁴; 佐野 有司³; 小宮 剛²
TASHIRO, Takayuki^{1*}; ISHIDA, Akizumi³; HORI, Masako³; IGISU, Motoko⁴; SANNO, Yuji³;
KOMIYA, Tsuyoshi²

¹ 東京大学大学院理学系研究科地球惑星科学専攻, ² 東京大学大学院総合文化研究科広域科学専攻, ³ 東京大学大気海洋研究所, ⁴ 独立行政法人海洋研究開発機構

¹Department of Earth and Planetary Science Graduate School of Science The University of Tokyo, ²Department of Earth Science & Astronomy Graduate School of Arts and Sciences The University of Tokyo, ³Atmosphere and Ocean Research Institute The University of Tokyo, ⁴Japan Agency for Marine-Earth Science and Technology

Elucidation of origin of life is an everlasting challenge but it provides an important constraint on the origin of life to find evidence for early life. So far, the oldest evidences for biogenic carbonaceous materials were reported from the 3.80 Ga Isua supracrustal belt based on carbon isotope ratio (Rosing, 1999) and morphological features (Ohtomo et al., 2013). But, the origin of carbonaceous materials in the 3.83 Ga Akilia Association (Fedó and Whitehouse, 2002) and 3.75 Ga Nuvvuagittuq Supracrustal Belt is still ambiguous (Papineau et al., 2011).

To understand the origin of organic matter in the Eoarchean and find older organic matter, we investigated occurrence and carbon isotope values of carbonaceous material in the >3.95 Ga metasediment rocks from the Saglek Block, northern Labrador, Canada. The metasediment rocks underwent the amphibolite to granulite facies metamorphism, but some avoid pervasive elemental mobility during the metamorphism. We observed thin sections of pelitic rocks (n = 70), conglomerates (n = 14), carbonate rocks (n = 39), cherts (n = 30), and chert nodules in carbonate rocks (n = 3) from over 2000 samples. Among the metasedimentary rocks (n = 156), 54 specimens including the pelitic rocks (n = 21), conglomerates (n = 4), carbonate rocks (n = 26) and chert nodules in carbonate rocks (n = 3) contain carbonaceous materials. Twenty-nine rock samples with the carbonaceous materials were selected for $\delta^{13}\text{C}_{org}$ analysis: pelitic rocks (n = 20), conglomerates (n = 4), carbonate rocks (n = 3) and chert nodules (n = 2). $\delta^{13}\text{C}_{org}$ values of the pelitic rocks range from -27.5 to -11.6 ‰. The $\delta^{13}\text{C}_{org}$ value increases as increasing in the metamorphic grade from amphibolite to granulite facies, indicating that the minimum $\delta^{13}\text{C}_{org}$ value reflects a primary signature. Raman spectroscopic observation of the carbonaceous materials showed that the matter comprises crystalline graphite, consistent with the intense thermal metamorphism. The $\delta^{13}\text{C}_{carb}$ values of carbonate rocks (n = 3) range from -3.8 to -2.6 ‰. Because it is well-known that the $\delta^{13}\text{C}_{carb}$ value decreases due to secondary alteration and metamorphism, the primary $\delta^{13}\text{C}_{carb}$ value was estimated to be higher than -2.6 ‰.

The minimum fractionation between the $\delta^{13}\text{C}_{org}$ and $\delta^{13}\text{C}_{carb}$ reaches 25 ‰, indicating biologic origin for the carbonaceous materials. This work presents the organism has already existed ca. 3.95 Ga. The large fractionation up to 25 ‰ implies autotrophs utilizing the reductive acetyl-CoA pathway or Calvin cycle in the Eoarchean.

キーワード: 炭質物, 炭酸塩, 初期生命, 炭素同位体, ラブラドル

Keywords: carbonaceous material, carbonate, early life, carbon isotope, Labrador