Japan Geoscience Union Meeting 2015

(May 24th - 28th at Makuhari, Chiba, Japan) ©2015. Japan Geoscience Union. All Rights Reserved.

HCG35-02



時間:5月27日09:30-09:45

黒部川堆積物のESR及びTLによる解析 ESR and TL signals in quartz in the present river sediments along with the Kurove river

吉田 真徳¹; 豊田 新¹*; 高田 将志²; 島田 愛子³; 蜷川 清隆¹ YOSHIDA, Msanori¹; TOYODA, Shin^{1*}; TAKADA, Masashi²; SHIMADA, Aiko³; NINAGAWA, Kiyotaka¹

¹ 岡山理科大学理学部,² 奈良女子大学文学部,³JEOL RESONANCE ¹Faculty of Science, Okayama University of Science, ²Nara Women's University, ³JEOL RESONANCE

The variations in the number of oxygen vacancies, measured as the ESR (electron spin resonance) intensity of the E_1 ' center in quartz, and in the TL (thermoluminescence) colors in quartz of quartz of the sediments are investigated along with the Kurobe river together with those in the bedrocks in the river reaches. The number of oxygen vacancy in quartz indicates the age of the host rocks from which the sediments have been generated by weathering while the red TL color corresponds to volcanic origin and blue to plutonic origin. The quantitative TL color measurements were made possible with the time-resolved 2 dimensional TL spectroscopy system.

The number of oxygen vacancies, higher in the upper reach, gets lower in the middle, and further higher in the lower reaches due to the inflow of the sediments originated from the younger and from the older bedrocks, respectively. The high temperature red to low temperature blue TL ratios, higher in the upper reach, gets lower in the middle and lower reaches due to the sediments having lower values. The change in the values along with the river flow is found to be explained by the inflow of the sediments originated from the bedrocks around the river.

キーワード: 河川堆積物, 供給源, ESR, TL Keywords: river sediment, provenance, ESR, TL