

内部発熱対流におけるセルサイズのRa数依存性に関する現象論モデルについて  
A phenomenological model for convective cell size in a fluid layer with internal heat generation at low Rayleigh numbers

熊谷 一郎<sup>1\*</sup>; 阿部 竜太<sup>2</sup>; 田坂 裕司<sup>2</sup>; Davaille Anne<sup>3</sup>; 村井 祐一<sup>2</sup>; 柳澤 孝寿<sup>4</sup>  
KUMAGAI, Ichiro<sup>1\*</sup>; ABE, Ryuta<sup>2</sup>; TASAKA, Yuji<sup>2</sup>; DAVAILLE, Anne<sup>3</sup>; MURAI, Yuichi<sup>2</sup>;  
YANAGISAWA, Takatoshi<sup>4</sup>

<sup>1</sup> 明星大学 理工学部, <sup>2</sup> 北海道大学 工学研究院, <sup>3</sup> FAST, CNRS, パリ南大学, <sup>4</sup> 海洋研究開発機構 地球深部ダイナミクス研究分野

<sup>1</sup> School of Science and Engineering, Meisei University, <sup>2</sup> Faculty of Engineering, Hokkaido University, <sup>3</sup> Laboratoire FAST, CNRS, Universite Pairs-Sud, <sup>4</sup> Department of Deep Earth Structure and Dynamics Research, JAMSTEC

The convective behavior of a fluid layer with internal heat generation at low Rayleigh number ( $6 \leq RaI/RaIc \leq 12$ ) was experimentally investigated. The horizontal fluid layer of 0.5 wt% KCl water solution was internally heated by Joule heating using the electric current. We quantitatively measured 2-D temperature and velocity fields by seeding the micro-encapsulated thermo-chromic liquid crystals in the fluid layer. We experimentally obtained the fluid dynamic scaling on non-dimensional temperature and the maximum downwelling velocity as a function of the Rayleigh number, and also refined the experimental data obtained by the previous studies. The scaling relations were combined with a phenomenological model based on the stability of the top thermal boundary layer. This phenomenological model consistently explained the increase in convective wavelength with increasing the Rayleigh number.

Keywords: internal heat generation, natural convection, cell size, visualization, experiment, fluid mechanics