

スロバキア Kardolína セクションにおける上部三畳系石灰岩の堆積環境と炭素・酸素安定同位体比変動  
Carbon and oxygen isotopes record of Upper Triassic limestone in the Kardolína section, Slovakia

白水 秀子<sup>1\*</sup>; Michalik Josef<sup>2</sup>; 日下 宗一郎<sup>3</sup>; 山下 美沙<sup>4</sup>; 山下 勝行<sup>4</sup>; 尾上 哲治<sup>1</sup>  
SHIROZU, Hideko<sup>1\*</sup>; MICHALIK, Josef<sup>2</sup>; KUSAKA, Soichiro<sup>3</sup>; YAMASHITA, Misa<sup>4</sup>; YAMASHITA, Katsuyuki<sup>4</sup>;  
ONOUE, Tetsuji<sup>1</sup>

<sup>1</sup> 熊本大学大学院自然科学研究科, <sup>2</sup> スロバキア科学アカデミー, <sup>3</sup> 総合地球環境学研究所, <sup>4</sup> 岡山大学大学院自然科学研究科

<sup>1</sup>Graduate School of Science and Technology, Kumamoto University, <sup>2</sup>Slovak Academy of Sciences, <sup>3</sup>Research Institute for Humanity and Nature, <sup>4</sup>Graduate School of Natural Science and Technology, Okayama University

Triassic / Jurassic (T/J) boundary of approximately 201.3 million years ago is known as a stratigraphic boundary recorded one of the big five Phanerozoic mass extinctions. Catastrophic processes such as widespread eruption of the Central Atlantic Magmatic Province (CAMP) flood basalts and extraterrestrial impacts have been proposed to account for the mass extinction event. This paper reports the results of carbon and oxygen isotopes analysis from the uppermost Triassic limestone in the Kardolína section, Slovakia. The Kardolína section crops out on a steep western slope of the Mt Palenica in the Belianske Tatry Mts as the most continuous section from the uppermost Triassic (Rhaetian) Fatra Formation. The Fatra Formation is shallow marine carbonate sequence and is overlain with a sharp contact by marine shale of the lowermost Jurassic (Hettangian) Kopieniec Formation. The Rhaetian age of the Fatra Formation was determined by foraminifera fossils. Several excursions of carbon and oxygen isotopes exist in the Fatra Formation. The large negative carbon isotope excursions (CIE) were found in at least four stratigraphic levels. Although the Fatra Formation is composed of packstone and grainstone, the CIE occurred during the deposition of non-fossiliferous lime-mudstone or ostracod lime-mudstone. The largest negative CIE occurs just before the T/J boundary. The cause of this negative CIE at the vicinity of T/J boundary remains uncertain. However it is assumed that the Kardolína section was exposed on the ground by sea-level fall, and subjected to diagenesis of terrestrial water. The large CIE may possibly be associated with sea-level change of the Kardolína section.

キーワード: 炭素・酸素安定同位体比, 石灰岩, 三畳紀/ジュラ紀境界, レーティアン, 炭素同位体比異常, 海水準変動  
Keywords: carbon and oxygen isotopes, limestone, Triassic/Jurassic boundary, Rhaetian, carbon isotope excursion, sea-level change