

沈み込みチャネルとマントルウェッジの炭素を含む塩水 Carbon-bearing saline fluids in the subduction channel and mantle wedge

川本 竜彦^{1*}
KAWAMOTO, Tatsuhiko^{1*}

¹ 京都大学理学研究科地球熱学

¹ Inst Geotherm Sci, Grad School Sci, Kyoto Univ

We find C-bearing saline fluids in the subduction channel and mantle wedge. Saline fluids are found with or without methane in jadeitites of serpentinite melanges located in Southwest Japan [Mori, Shigeno, Kawamoto, Nishiyama, in progress]. Carbon dioxide-bearing saline fluid inclusions are also reported from sub-arc mantle peridotite xenoliths: 3.7 wt% NaCl in Ichinomegata lherzolites, Northeast Japan arc [Kumagai et al., 2014] and 5.1 wt% NaCl in Pinatubo harzburgites, Luzon arc [Kawamoto et al., 2013]. These findings indicate that aqueous fluids in the subduction channel and mantle wedge can contain certain amounts of C and Cl.

We suggested that separation of slab-derived supercritical fluids into aqueous fluids and melts plays an important role in elemental transfer from subducting slab to the mantle wedge [Kawamoto et al., 2012]. It is, therefore, important to determine the effect of Cl on the trace element partitioning between aqueous fluids and melts. Synchrotron radiation X-ray fluorescence (XRF) analysis is conducted to know Rb, Sr, and Pb partitioning between aqueous fluids and melts simultaneously at high-temperature and high-pressure conditions. There is a positive correlation between partition coefficients and pressure, as well as salinity [Kawamoto et al., 2014]. Two slab-derived components, melt and fluid components, are suggested to explain trace element characteristics of arc-basalts in the Mariana arc [Pearce et al., 2005]. The fluid component is characterized by enrichment of alkali, alkali earth elements, and Pb. Such features can be explained if the fluid component is a saline fluid, because alkali earth elements and Pb are much less mobile with Cl-free fluids than Cl-rich fluids [Kawamoto et al., 2014].

We suggest that slab-derived components have compositional features consistent with a saline fluid and a melt, which can be formed through a separation of a slab-derived supercritical fluid [Kawamoto et al., 2012, 2014]. Slab-derived supercritical fluids contain Cl, and separated aqueous fluids inherit much of the Cl and some of the large-ion lithophile elements. Dissolution of carbon materials into aqueous fluids is enhanced by the salinity [Newton and Manning 2002] and their species can be controlled by oxygen fugacity.

Reference

Kawamoto T., Kanzaki M., Mibe K., Matsukage K. N., Ono S., 2012, Separation of supercritical slab-fluids to form aqueous fluid and melt components in subduction zone magmatism. *Proceedings of the National Academy of Sciences, U. S. A.*, 109, 18695-18700.

Kawamoto T., Yoshikawa M., Kumagai Y., Mirabueno M. H. T., Okuno M., Kobayashi T., 2013, Mantle wedge infiltrated with saline fluids from dehydration and decarbonation of subducting slab. *Proceedings of the National Academy of Sciences, U. S. A.*, 110, 9663-9668.

Kawamoto T., Mibe K., Bureau H., Reguer S., Mocuta C., Kubsy S., Thiaudiere D., Ono S., Kogiso T., 2014, Large ion lithophile elements delivered by saline fluids to the sub-arc mantle, *Earth, Planets and Space*, 66, 61.

Kumagai Y., Kawamoto T., Yamamoto J., 2014, Evolution of carbon dioxide bearing saline fluids in the mantle wedge beneath the Northeast Japan arc, *Contributions to Mineralogy and Petrology*, 168, 1056.

Newton, R.C. and Manning, C.E., 2002, Experimental determination of calcite solubility in H₂O-NaCl solutions at deep crust/upper mantle pressures and temperatures: implications for metasomatic processes in shear zones. *American Mineralogist*, 87, 1401-1409.

Pearce J. A., Stern R. J., Bloomer S. H., Fryer P., 2005, Geochemical mapping of the Mariana arc-basin system: Implications for the nature and distribution of subduction components. *Geochemistry, Geophysics, Geosystems*, 6, Q07006.

キーワード: 水流体, 流体包有物, 島弧マグマ

Keywords: aqueous fluid, fluid inclusion, arc magma