

PEM25-04

会場:202

時間:5月25日 09:45-10:00

イオンスケール・ホイッスラー乱流のカスケードについて Forward cascade of whistler turbulence at ion scales

齊藤 慎司^{1*}; 成行 泰裕²; 梅田 隆行³
SAITO, Shinji^{1*}; NARIYUKI, Yasuhiro²; UMEDA, Takayuki³

¹名古屋大学大学院理学研究科, ²富山大学人間発達科学部, ³名古屋大学太陽地球環境研究所
¹Nagoya University, ²University of Toyama, ³STE Laboratory, Nagoya University

Two-dimensional electromagnetic particle-in-cell simulations in magnetized, homogeneous, collisionless electron-ion plasma demonstrate the forward cascade of whistler turbulence at ion scales. Saito et al. (2015, submitted) emphasized that the modified two-stream instability could contribute the dissipation of kinetic turbulence at ion scales, by demonstrating rapid damping of a monochromatic ion-scale whistler wave in two-dimensional particle-in-cell simulation. The instability is driven by the electric current fluctuation perpendicular to the mean magnetic field. Through the development of the instability, electrons and ions are scattered in the directions parallel and perpendicular to the mean magnetic field, respectively. We expect that the forward cascade of whistler turbulence and the dissipation related to the modified two-stream instability contribute plasma heating and have key role of variability of power-law index of magnetic spectrum at ion scale in solar wind. Solar wind observations show that larger cascade rates of turbulence lead to steeper power-law magnetic spectra. The instability driven dissipation could explain property of the magnetic spectra at ion scales. Discussion will focus on properties of whistler turbulence, such as the power-law index, wavenumber anisotropy, electron and ion heating, through the forward cascade of decaying whistler turbulence.

キーワード: 運動論的乱流, プラズマ加熱, 波動粒子相互作用
Keywords: kinetic turbulence, plasma heating, wave-particle interaction