

ケルビン・ヘルムホルツ不安定性におけるイオンジャイロ運動効果 Ion kinetic effects to nonlinear processes of the Kelvin-Helmholtz instability

梅田 隆行^{1*}; 上野 悟志¹; 中村 琢磨²

UMEDA, Takayuki^{1*}; UENO, Satoshi¹; NAKAMURA, Takuma²

¹ 名古屋大学太陽地球環境研究所, ² オーストリア科学アカデミー宇宙科学研究所

¹-Terrestrial Environment Laboratory, Nagoya University, ²Space Research Institute, Austrian Academy of Sciences

The nonlinear evolution of the Kelvin-Helmholtz (KH) instability at a transverse velocity shear layer in an inhomogeneous space plasma is investigated by means of a four-dimensional (two spatial and two velocity dimensions) electromagnetic Vlasov simulation. When the rotation direction of the primary KH vortex and the direction of ion gyro motion are the same (i.e. the inner product between the vorticity of the primary velocity shear and the magnetic field vector is negative) there exists a strong ion cyclotron damping. In this case, spatial inhomogeneity inside the primary KH vortex is smoothed and the secondary Rayleigh-Taylor/KH instabilities are suppressed. It is also found that another secondary instability on the electron inertial scale is simultaneously generated at secondary shear layers for both cases, but at different locations. The small-scale secondary instability takes place only when the inner product between the vorticity of the secondary shear layer and the magnetic field vector is positive, suggesting the damping of small-scale processes by ion gyro motion. These results indicate that secondary instabilities occurring in the nonlinear stage of the primary KHI show different evolutions depending on the sign of the inner product between the magnetic field and the vorticity of the velocity shear layer. The difference of the nonlinear evolution depending on the ion-to-electron mass ratio is also discussed.

キーワード: ケルビン・ヘルムホルツ不安定性, 計算機シミュレーション, ヴラソフ方程式, 非 MHD

Keywords: Kelvin-Helmholtz instability, computer simulation, Vlasov equation, non-MHD