

## AE および Dst の出現強度分布と太陽風パラメータ Intensity distribution of AE and Dst and its relation to solar wind parameters

森岡 昭<sup>1\*</sup>; 三好 由純<sup>2</sup>; 能勢 正仁<sup>3</sup>  
MORIOKA, Akira<sup>1\*</sup>; MIYOSHI, Yoshizumi<sup>2</sup>; NOSE, Masahito<sup>3</sup>

<sup>1</sup> 東北大学惑星プラズマ大気研究センター, <sup>2</sup> 名古屋大学太陽地球環境研究所, <sup>3</sup> 京都大学地磁気世界資料解析センター  
<sup>1</sup>PPARC, Tohoku University, <sup>2</sup>STEL, Nagoya University, <sup>3</sup>WDC for Geomag, Kyoto University

Storm-substorm relationship and related solar wind-magnetosphere coupling process are studied on the basis of statistical analyses of AE, Dst, epsilon parameter and Em using OMNI data base from 1995 to 2013 and Wp index data from 2005. The statistical relationship between AE and Dst is examined to clarify the difference between CME storm and CIR storm. The intensity distribution of AE and Dst for a year is compared with that of epsilon and Em parameters in the solar wind.

The obtained major results are,

- 1). Relationship between AE at substorm and Dst is rather linear.
- 2). AE vs Dst relationship at CIR storm is different from that at CME storm.
- 3). Intensity distribution of AE and Dst for a year shows the exponential distribution.
- 4). Intensity distribution of epsilon parameter for a year shows the power law distribution.
- 5). Intensity distribution of Em for a year shows the exponential distribution.
- 6). The results 3) to 5) suggest that magnetospheric disturbances are mainly controlled by the solar wind electric field rather than by solar wind Poynting flux.

キーワード: サブストーム, 磁気嵐, 太陽風相互作用

Keywords: substorm, magnetic storm, solar wind interaction