

Geotailで観測された昼側磁気圏境界面での磁気リコネクション II Geotail observations of dayside magnetopause reconnection II

小田切 修一^{1*}; 吉田 和史¹; 長井 嗣信²
ODAGIRI, Shuichi^{1*}; YOSHIDA, Kazufumi¹; NAGAI, Tsugunobu²

¹ 東工大・地惑, ² 東京工業大学

¹Earth and Planetary Sciences, Titech, ²Tokyo Institute Of Technology

Plasma velocity distributions perpendicular to the magnetic field are generally isotropic by Larmor motion of ions and electrons. In actuality, isotropic velocity distributions are observed by Geotail. However, anisotropic ion velocity distributions were observed in the magnetosheath nearby the magnetopause when Geotail crossed the dayside magnetopause and observed ion flow jets by magnetic reconnection. The Geotail data of ion Energy-Time spectrogram on July 6, 2013 indicate anisotropic velocity distributions of ions energies higher than 20 keV at 0330 UT. The Geotail orbit is from magnetosheath through the magnetopause to the magnetosphere. The spacecraft GSM coordinates at the time of anisotropic ion velocity distribution observation are (9.8, 3.0, -0.2) R_E . This Geotail position is in the magnetosheath nearby the magnetopause. Ion energies are about 1 keV in the distant magnetosheath from the magnetopause. There are no ions with energies higher than 10 keV in the magnetosheath. There are ions with energies higher than 20 keV in magnetosphere. Thus, these ions are considered to go out toward the magnetosheath from the magnetosphere. We explain anisotropic ion velocity distributions by reconnecting magnetic field geometry.

キーワード: 磁気圏, 磁気リコネクション

Keywords: magnetosphere, magnetic reconnection