Japan Geoscience Union Meeting 2015

(May 24th - 28th at Makuhari, Chiba, Japan)

©2015. Japan Geoscience Union. All Rights Reserved.



SEM34-08

会場:102A

時間:5月24日12:00-12:15

多様な蛇紋岩の磁性:中央インド洋海嶺ヨコニワ産の海底試料による制約 Variation in magnetic properties of serpentinized peridotites from Yokoniwa Rise, Central Indian Ridge

藤井 昌和 ^{1*}; 沖野 郷子 ¹; 佐藤 暢 ²; 中村 謙太郎 ³; 山崎 俊嗣 ¹ FUJII, Masakazu^{1*}; OKINO, Kyoko¹; SATO, Hiroshi²; NAKAMURA, Kentaro³; YAMAZAKI, Toshitsugu¹

Serpentinization of ultramafic rocks through hydrothermal alteration changes the physical, rheological, chemical, and magnetic properties of the oceanic lithosphere. Recent discovery of widespread exposures of serpentinized mantle materials on the seafloor in a slow-spreading environment renewed interest for this alteration process. However, we have limited understanding of the serpentinization mechanism because of the lack of data measured from seafloor rocks. Since magnetite is a direct product of serpentinization process, magnetic properties of serpentinized peridotites can be a good indicator to understand the process. We collected 30 peridotite samples of different degrees of serpentinization from the seafloor on the non-transform-offset massif called as the Yokoniwa Rise in the Central Indian Ridge. These 30 samples yielded a wide range (17-100%) of serpentinization degrees and provide us a good data set to evaluate the relationship between serpentinization and magnetic properties. The measured range of magnetic parameters are as follow; natural remanent magnetization (0.2-8.4 A/m), magnetic susceptibility (0.002-0.087 SI), and magnetite amount (0.1-5.5 wt%). The amount of magnetite varies nonlinearly, likely exponentially, as a function of serpentinization degree. Remarkable increase of magnetite amount occurs in samples with high degree of serpentinization (>70%), indicating larger production of magnetite during the late stage of serpentinization process. The results provide key constraints on the serpentinization mechanism, and insights on the potential of serpentinized mantle to contribute to marine magnetic anomalies.

キーワード: 蛇紋岩, 上部マントル, 岩石磁気, 低速拡大海嶺

Keywords: serpentinized peridotite, upper mantle, rock magnetics, slow spreading ridge

¹ 東京大学大気海洋研究所, 2 専修大学, 3 東京大学

¹AORI, UTokyo, ²Senshu University, ³UTokyo