

秋田県一ノ目潟のコア堆積物に保存された残留磁化 Remanent magnetization observed in core sediments from the Ichinomegata Maar, Akita Prefecture

安楽 和央^{1*}; 林田 明²; 山田 和芳³; 篠塚 良嗣⁴; 米延 仁志⁵; 五反田 克也⁶; 原口 強⁷
ANRAKU, Kazuhiro^{1*}; HAYASHIDA, Akira²; YAMADA, Kazuyoshi³; SHINOZUKA, Yoshitsugu⁴;
YONENOBU, Hitoshi⁵; GOTANDA, Katsuya⁶; HARAGUCHI, Tsuyoshi⁷

¹ 同志社大学大学院理工学研究科, ² 同志社大学工学部環境システム学科, ³ 静岡県 文化・観光部 文化学術局 ふじのくに地球環境史ミュージアム整備課, ⁴ 北海道大学大学院地球環境科学研究院, ⁵ 鳴門教育大学大学院学校教育研究科, ⁶ 千葉商科大学政策情報学部, ⁷ 大阪市立大学大学院理学研究科

¹Graduate School of Science and Engineering, Doshisha University, ²Department of Environmental Systems Science, Doshisha University, ³Museum of Natural and Environmental history, Shizuoka, ⁴Faculty of Environmental Earth Science, Hokkaido University, ⁵Graduate School of Education, Naruto University of Education, ⁶Faculty of Polycy Informatics, Chiba University of Commerce, ⁷Department of Geosciences, Graduate School of Science, Osaka City University

The Ichi-no-megata is a maar lake with a maximum water depth about 45 m, located in the Oga Peninsula on the Japan Sea coast. A piston-core sample (IMG13P) obtained in 2013 provided a Holocene paleomagnetic secular variation (PSV) record through measurements of natural remanent magnetization (NRM) of 7cm³ cubic samples. Also, we collected Mackereth core samples (IMG13M-1 and IMG13M-2) at the center of the lake, and measured magnetic susceptibility, anisotropy of magnetic susceptibility (AMS) and NRM of 7cm³ cubic samples.

The core sediments are mostly composed of laminated clay or silt intercalating sandy turbidite layers. According to correlation between the Mackereth cores and the piston cores based on lithological and magnetic susceptibility data, it is revealed that uppermost parts of the piston cores were missed. Inclinations of minimum axis and shape parameters *q* of AMS ellipsoids indicate that turbidite layers of upper parts have not preserved primary sedimentary fabric. Stepwise AF demagnetization of the NRM showed that remanence of the laminated layer is more stable than turbidite layers. Therefore, we argue that the turbidite sediment is not suitable for preservation of NRM.

キーワード: 残留磁化, 初磁化率, 一ノ目潟

Keywords: Remanent magnetization, magnetic susceptibility, Ichi-no-megata Maar