

## 高速拡大海嶺下の下部地殻の進化過程：Hess Deep Rift で採取されたトロクトライトの例 Evolutional process of fast-spreading lower oceanic crust: an example of troctolites at the Hess Deep Rift

秋澤 紀克<sup>1\*</sup>; ゴダー マルゲリット<sup>2</sup>; イルデフォンス ベノア<sup>2</sup>; 荒井 章司<sup>1</sup>  
AKIZAWA, Norikatsu<sup>1\*</sup>; GODARD, Marguerite<sup>2</sup>; ILDEFONSE, Benoit<sup>2</sup>; ARAI, Shoji<sup>1</sup>

<sup>1</sup> 金沢大学 地球学科, <sup>2</sup> モンペリエ大学 地球科学科

<sup>1</sup>Department of Earth Sciences, Kanazawa University, <sup>2</sup>Geosciences Montpellier, Universite Montpellier & CNRS

Troctolites were recovered during IODP Expedition 345 at Hess Deep Rift (Dec 2012 - Feb 2013), which targeted plutonic rocks from fast-spread lower ocean crust. The troctolites are divided into two groups based on textural differences; fine-grained troctolite (including a skeletal olivine-bearing troctolite sample), and coarse-grained troctolite.

The major-element compositions of olivine, plagioclase and clinopyroxene in coarse-grained troctolites are intermediate between those in olivine gabbros/olivine-bearing gabbros and peridotites recovered from the Hess Deep Rift. Fo content and NiO of olivine range from 87 to 89, and 0.2 to 0.3 wt.%, respectively. An content of plagioclase ranges from 85 to 90. Mg# and Cr<sub>2</sub>O<sub>3</sub> of clinopyroxene range from 0.88 to 0.91, and 0.5 to 1.2 wt.%, respectively. In contrast, fine-grained troctolites partly overlap with olivine gabbros/olivine-bearing gabbros in mineral chemistry. Fo content and NiO of olivine range from 83 to 86, and 0.08 to 0.2 wt.%, respectively. An content of plagioclase ranges from 77 to 84. Mg# and Cr<sub>2</sub>O<sub>3</sub> of clinopyroxene range from 0.82 to 0.89, and nearly nil to 1.0 wt.%, respectively. Trace-element analyses of olivine and plagioclase show progressive enrichment in REE from coarse-grained to fine-grained troctolites. In contrast, clinopyroxenes show scattered trace-element compositions in the fine-grained troctolites, even in a single thin section.

The changes in chemical composition of olivine and plagioclase from coarse-grained to fine-grained (and skeletal olivine-bearing) troctolites may be ascribed to variable degrees of reequilibration with crystallizing melts during cooling. Fine-grained troctolites possibly record melt/rock interactions that would be responsible for the variable chemical compositions of clinopyroxenes. At Hess Deep, lower crustal troctolites possibly underwent several stages of evolution, combining fractional crystallization of MORB (mid-ocean ridge basalt) melts, combined with melt/troctolite interaction during migration. Melt migration processes in the lower oceanic crust would result in enhanced regional diversity of MORB chemistry.

キーワード: トロクトライト, 高速拡大海嶺, メルト/トロクトライト相互反応, Hess Deep, 微量元素組成  
Keywords: Troctolite, Fast-spreading ridge, Melt/troctolite interaction, Hess Deep, Trace-element composition