

## Single-crystal Brillouin Spectroscopy with Laser Heating and Variable $q$ : Design and Results on Olivine

ZHANG, Jin <sup>s1</sup> ; BASS, Jay<sup>2\*</sup>

<sup>1</sup>Univ of Illinois & COMPRES, <sup>2</sup>COMPRES & Univ of Illinois

We have developed a novel Brillouin spectroscopy system integrated with CO<sub>2</sub> laser heating and Raman spectroscopic capabilities. High-pressure laser heating experiments on liquid water compressed in a diamond-anvil cell up to 2500 +/- 150 K demonstrate the flexibility and performance of the system. Temperature is determined from the grey-body thermal radiation of the heated samples. New single-crystal laser heating Brillouin measurements were made on San Carlos Olivine in the (111) plane at pressures up to ~13 GPa, and T=1300K. We obtain quantitative values for the thermal pressure in the diamond cell. Using KCl and KBr and pressure-transmitting media, we show that pressure gradients in the sample chamber are small at high P-T conditions based on observations of the olivine-wadsleyite transition. This system is additionally designed for continuously varying scattering angles from near forward scattering (0° scattering angle) up to near back scattering (~141°). Our results on the sound velocities of olivine at high pressure-temperature conditions have implications for the nature of the 410 km discontinuity and the olivine content of the transition zone.

Keywords: Elastic properties, Brillouin scattering, Equations of state, Olivine