

## Studies on variations of climate and ice sheet under the international deep ice core project at the North East Greenland Ice Stream

\*Kumiko Goto-Azuma<sup>1,2</sup>, Kenji Kawamura<sup>1,2</sup>, Shuji Fujita<sup>1,2</sup>, Jun'ichi Okuno<sup>1,2</sup>, Ayako Abe-Ouchi<sup>3</sup>, Ralf Greve<sup>4</sup>, Fuyuki SAITO<sup>5</sup>, Tomoyuki Homma<sup>6</sup>, Nobuhiko Azuma<sup>6</sup>, Hiroyuki Enomoto<sup>1,2</sup>, Hideaki Motoyama<sup>1,2</sup>, Dorthe Dahl-Jensen<sup>7</sup>

1.National Institute of Polar Research, 2.SOKENDAI, 3.University of Tokyo, 4.Hokkaido University, 5.JAMSTEC, 6.Nagoka University of Technology, 7.University of Copenhagen

The Greenland Ice Sheet has recently been experiencing drastic changes, such as extended summer melting and increasing mass losses. There is an urgent need to understand the mechanisms of such changes because they are directly linked to global sea level rise. Greenland ice cores have so far provided valuable information on melt events and changes in the surface mass balance in the past. Moreover, the data obtained from multiple deep ice cores drilled during the last few decades, combined with modeling studies, have recently enabled us to reconstruct the past changes of Greenland Ice Sheet elevation. The previous ice cores were drilled at sites with minimal horizontal ice flow, as the main purpose of the past ice coring projects was to reconstruct the past climate and environment at the drill sites. Information on ice flow dynamics obtained from such ice cores has therefore been limited.

Understanding the mechanisms of basal sliding and ice deformation is a prerequisite for better projections of the future changes of the Greenland Ice Sheet and sea level rise. To understand the Greenland Ice Sheet dynamics, the East Greenland Ice Core Project (EGRIP) was proposed by the University of Copenhagen. Japan, Germany, Norway, U.S.A., France and Switzerland have been invited to participate in this international project. Under EGRIP, a deep ice core to the bed will be drilled at the onset of the North-East Greenland Ice Stream (NEGIS), where horizontal flow velocity is expected to be several tens of meters per year. As NEGIS is the largest ice stream in Greenland, the EGRIP ice core will certainly advance our knowledge on the dynamics and past changes of the Greenland Ice Sheet. The EGRIP core will also give us an ideal opportunity to reconstruct the climate and environment changes during the early Holocene, which was considered to be warmer than today and should be an excellent analogue to the future Greenland affected by global warming. The results from the EGRIP core will fill the gap of our knowledge due to the lack of high-resolution, detailed ice core records from the early Holocene.

Japan will participate in the EGRIP under the ArCS (Arctic Challenge for Sustainability) project, a recently funded national project. The first EGRIP steering committee meeting was held in Copenhagen in late October 2015. At the meeting, drilling and fieldwork plans were presented by the University of Copenhagen. Scientific plans were proposed by all the participant nations. At the JpGU meeting, we will present the scientific purposes and plans, together with the current status of the EGRIP and logistic plans.

Keywords: EGRIP, Greenland, Ice core

## Japanese glaciological activity in Antarctica for the next six years (JARE Phase IX)

\*Kenji Kawamura<sup>1</sup>, Hideaki Motoyama<sup>1</sup>, Kumiko Goto-Azuma<sup>1</sup>, Shuji Fujita<sup>1</sup>, Dome Fuji Ice Core Consortium

1.National Institute of Polar Research, Research Organization of Information and Systems

The Dome Fuji ice core covers the past 720,000 years, and its analyses have been conducted. During the Phase VIII of Japanese Antarctic Research Expedition, we made advancements in glaciological activities such as transportation of the second Dome Fuji ice core, shallow and intermediate drillings, and large-area glaciological surveys.

International Partnership in Ice Core Science (IPICS), to which Japan has participated, has identified the "oldest ice core" project, which aims to collect 1.5 million-years-old ice cores from Antarctica, as a most important and challenging project for the international ice core community. We plan to contribute to this project and started the consideration of next deep drilling site in the vicinity of the current Dome Fuji station.

During the next six years, we aim to conduct glaciological surveys, identify a suitable coring site, and start drilling to the first few hundred meters. In addition to the new drilling, a part of the second Dome Fuji core, stored at the station, must be transported to Japan for the study of this core.

In the presentation, we will also introduce our planned coastal activities and updates on international circumstances about the oldest ice project.

Keywords: Japanese Antarctic Research Expedition, Climate change, Ice core

## Determination of the ice drilling site in the next deep ice coring project of the "oldest ice" at Dome Fuji

\*Shuji Fujita<sup>1,2</sup>, Kenji Kawamura<sup>1,2</sup>, Kumiko Goto-Azuma<sup>1,2</sup>, Hideaki Motoyama<sup>1,2</sup>, . Dome Fuji Ice Core Consortium<sup>3</sup>

1.National Institute of Polar Research, Research Organization of Information and Systems(ROIS),  
2.The Graduate University for Advanced Studies (SOKENDAI) , 3.Dome Fuji Ice Core Consortium

In the next phase of the Antarctic research activities of Japan, National Institute of Polar Research along with Dome Fuji Ice Core Consortium (ICC) in Japan plan to perform various activities related to drilling of the "oldest ice" with ages much older than 800 kyr ago, in the vicinity of Dome Fuji. We call the planned ice core as "the 3<sup>rd</sup> DF deep core". In the near future, we aim to: (i) investigate glaciological conditions (ice sheet surface conditions, englacial conditions and subglacial conditions) of the candidate site area; (ii) determine the exact location of the drilling site and (iii) start to perform pilot hole drilling, casing and shallow/middle range deep drilling. Prerequisites of presence of the very old ice are as follows. (i) The ice/bed boundaries must be frozen. (ii) The ice layers near the bed must be maintained as undisturbed by the ice sheet flow. (iii) During glacial/interglacial cycles, surface mass balance must be generally positive. To satisfy the prerequisite (i), presence of subglacial highland is necessary in the glaciological condition of the inland of Antarctica. We have shown in our earlier observational studies that in the vicinity of Dome Fuji ice sheet bed is frozen when ice thickness is less than ~2850 m. We have confirmed, at area ~50 km of the Dome Fuji summit, there is a wide area with ice thickness of ~2200 m, where the ice sheet bed is frozen. To satisfy the condition (ii), the site must be close to the present summit of Dome Fuji. The area ~50 km of the Dome Fuji summit satisfy this condition. It has been estimated that ice layers with ages older than ~800 kyrs ago are highly compressed within the deepest 5 % of the ice sheet thickness. Thus conditions for presence of such compressed ice and rheology is also a major subject of the ice coring project. As for the condition (iii), we have confirmed the condition is satisfied based on the internal layering observations with ice sounding radars. We already started the preliminary investigation of the radar sounding in 1997, and accumulated data since then. We have identified area for the site, and performed shallow drilling at the base camp of the area. In the near future, we will carry out radar surveys spatially in more detail, to determine the exact location of the deep ice coring.

Keywords: Antarctica, ice sheet, ice core

Ice core drilling and related observations by Jare57 in 2015/2016 austral summer on the coastal region of Antarctic ice sheet

\*Hideaki Motoyama<sup>1,2</sup>, Kenji Kawamura<sup>1,2</sup>, Toshimitsu Sakurai<sup>1</sup>, Kenji Sudo<sup>2</sup>, Miho Arai<sup>3</sup>, Toshitaka Suzuki<sup>3</sup>

1.National Institute of Polar Research, 2.Sokendai, 3.Yamagata University

We were carried out an ice core drilling on the coastal region of Antarctic ice sheet as a part of the 57th of Japanese Antarctic Research Expeditions (2015/2016 of austral summer). We selected the H128 site for drill site (69.40S, 41.55E, 1,380m a.s.l . 100km inland from the coast). Advantages of H128 site were as below; high time resolution, few changes of annual snow accumulation rates, low annual mean temperature (10m depth ice temperature) -24 C. We finally drilled to 260m in depth. We will be able to analyze the climate changes of past 2000 years. Firn air samplings were also carried out. We also conducted the borehole logging, snow pit observation, setup of automatic weather station.

Keywords: Antarctic ice sheet, ice core drilling, firn air sampling

## Organic aerosol tracer records in NGRIP ice core during the period from MCA to LIA

\*Osamu Seki<sup>1</sup>, Yohei Yamashita<sup>2</sup>, Kimitaka Kawamura<sup>1</sup>, Takuro Kobashi<sup>4</sup>, Kumiko Azuma<sup>3</sup>

1.Institute of Low Temperature Science, Hokkaido University, 2.Faculty of Environmental Earth Science, 3.National Institution for Polar Research, 4.University of Bern

Secondary organic aerosol (SOA) formed by the photooxidation of biogenic volatile organic compounds (BVOCs) and biomass burning organic aerosol are major components of organic aerosol in the atmosphere. Those aerosols are thought to play an important role in atmospheric chemistry and physics, as well as regional and global climate via direct and indirect effects on radiative balance. However, substantial impact of those aerosols on the Earth's climate remains unclear. To better understand the substantial role of those aerosols in the climate system, it is vital to investigate long-term variability of those aerosol components and property in a past. Here we for the first time report biogenic SOA- and biomass burning- (levoglucosan) tracers and UV absorption spectra of organic matter in ice cores from Greenland ice core (NGRIP) during the period from Medieval Climate Anomaly (MCA) to Little Ice Age (LIA). We find that the concentrations and composition of biogenic SOA- and biomass burning- tracers relate to climate with increases in the organic aerosol concentrations during MCA. On the other hand, UV absorption spectra of organic matter in NGRIP ice core also drastically changes associated with climate change. These results suggest dramatic changes in source, loading and property of organic aerosol from the warm MCA to cold LIA.

Keywords: ice core, organic aerosol, Medieval Climate Anomaly, Little Ice Age

## CH<sub>4</sub> concentrations during the Holocene reconstructed from the NEEM (Greenland) and Dome Fuji (East Antarctica) ice cores

\*Ikumi Oyabu<sup>1</sup>, Kenji Kawamura<sup>1</sup>, Kyotaro Kitamura<sup>1</sup>, Shuji Aoki<sup>2</sup>, Takakiyo Nakazawa<sup>2</sup>, Edward J Brook<sup>3</sup>, Thomas Blunier<sup>4</sup>

1.National Institute of Polar Research, Research Organization of Information and Systems, 2.Tohoku University, 3.Oregon State University, 4.Centre for Ice and Climate, Niels Bohr Institute, University of Copenhagen

Methane (CH<sub>4</sub>), the second most important anthropogenic greenhouse gas, has increased in the atmosphere by a factor 2.5 since the onset of the Industrial Revolution, which account for ~20% of the total increase in radiative forcing over that time<sup>[1]</sup>. Ice cores from both polar regions preserve the past atmospheric CH<sub>4</sub>, and thus have the potential to constrain the changes in CH<sub>4</sub> concentration difference between the polar regions. The inter polar difference of CH<sub>4</sub> is one of the approaches to understand the evolution of CH<sub>4</sub> budget and its relationship with climate. To reconstruct the CH<sub>4</sub> inter polar difference during the Holocene, we have been measuring CH<sub>4</sub> concentrations in the NEEM (Greenland) and Dome Fuji (Antarctica) ice cores over the period from 200 to 14500 years before present (yr BP), with a mean time resolution of ~50 years. Since most of this time period is overlapping with the brittle zone in the Greenland core, it is challenging to reconstruct accurate CH<sub>4</sub> concentration during the Holocene from the NEEM ice core.

Ice samples without visible cracks were carefully selected from the NEEM and Dome Fuji ice cores. We employed a newly established wet extraction system (an improved version of ref. 2) the National Institute of Polar Research, with a typical sample size of ~80 g (ice). The air released from ice was first collected into a sample tube (electropolished stainless steel tube with a metal-seal valve), and then it was split into two aliquots. One aliquot was measured by a gas chromatograph (Agilent Technologies 7890A) for CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O concentrations, and the other was measured by a mass spectrometer (Thermo DELTA V Plus) for δ<sup>15</sup>N of N<sub>2</sub>, δ<sup>18</sup>O of O<sub>2</sub>, δ(O<sub>2</sub>/N<sub>2</sub>), δ(Ar/N<sub>2</sub>) and total air content. We have measured 181 samples for the NEEM ice core. Analytical precision of CH<sub>4</sub> concentration was estimated to be ±2.4 ppb from the pooled standard deviation from duplicate measurements (n=53).

Before the Holocene, the NEEM CH<sub>4</sub> concentration is relatively high (620-705 ppb) during the Bolling-Allerød, and it rapidly decreases to <500 ppb during the Younger Dryas, and then increases to ~750 ppb at the beginning of the Holocene. During the Holocene, CH<sub>4</sub> concentration first decreases to the minimum of ~610 ppb around 5000 yr BP, and it increases afterwards. Our record agrees well with a high resolution CH<sub>4</sub> concentration record from the GISP2 ice core for last 2000 years<sup>[3]</sup>, and that of the NEEM ice core measured by a CFA system between 9500 and 14500 yr BP<sup>[4]</sup>.

We completed the NEEM measurements and started the Dome Fuji measurements, and the resulting CH<sub>4</sub> inter polar difference will be deduced and discussed in the presentation.

### References

- [1] P. Forster et al., in *Climate Change 2007: The Physical Science Basis*. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, S. Solomon et al., Eds. (Cambridge Univ. Press, Cambridge, 2007), pp. 131-234.
- [2] Kawamura et al. (2003). Atmospheric CO<sub>2</sub> variations over the last three glacial-interglacial climatic cycles deduced from the Dome Fuji deep ice core, Antarctica using a wet extraction technique. *Tellus B*, 55(2), 126-137.
- [3] Mitchell et al. (2013). Constraints on the Late Holocene Anthropogenic Contribution to the Atmospheric Methane Budget. *Science*, 342(6161), 964-966.

[4] Chappellaz et al. (2013). High-resolution glacial and deglacial record of atmospheric methane by continuous-flow and laser spectrometer analysis along the NEEM ice core. *Clim. Past*, 9(6), 2579-2593.

Keywords: Ice core, methane, Holocene, Greenland, Antarctica

## Glacial climate states and abrupt climate change in MIROC AOGCM

\*Ayako Abe-Ouchi<sup>1,2,3</sup>, Wing-Le Chan<sup>1</sup>, Rumi Ohgaito<sup>2</sup>, Kenji Kawamura<sup>3</sup>, Masakazu Yoshimori<sup>4</sup>, Akira Oka<sup>1</sup>, Sam Sherriff-Tadano<sup>1</sup>

1.Atmosphere and Ocean Research Institute, The University of Tokyo, 2.JAMSTEC, 3.NIPR, 4.Hokkaido University

Millennial climate change such as D-O cycles and AIM recorded in ice cores in both Hemispheres is known to show a relatively higher amplitude in the middle-level of a glacial cycle than in the interglacial state or severe glacial state. Here we ran several sensitivity experiments using a coupled atmosphere and ocean GCM (MIROC4m, renamed from MIROC3.2.2) and show that the response to fresh water release to the ocean and bipolar response is highly dependent on the background climate. The experiments were conducted with 500 years water hosing of 0.05 to 0.1 Sv (where 1 Sv is equivalent to the water flux of 10m sea level rise in 100 years) in the North Atlantic 50-70N under different basic states; modern climate state with the pre-industrial condition, middle glacial climate state and full glacial condition, mainly differing in the ice sheet configuration and atmospheric amount of Greenhouse Gases. The results under middle glacial condition show largest cooling/warming response in North Atlantic and a reasonable bipolar warming/cooling signal revealed in the ice core data of the both hemisphere. We discuss the responses under different background climates which involve the strong coupling between atmosphere, ocean and sea ice and their dependence on the configuration of ice sheet.

Keywords: Paleoclimate, Climate Model

## Long-term periodicities in solar activity between 170 and 200 kyr ago reconstructed from $^{10}\text{Be}$ in ice-core and sediment archives

\*Kazuho Horiuchi<sup>1</sup>, Kanae Kamata<sup>1</sup>, Shun Maejima<sup>1</sup>, Sho Sasaki<sup>1</sup>, Nobuyoshi Sasaki<sup>1</sup>, Toshitsugu Yamazaki<sup>2</sup>, Shuji Fujita<sup>3</sup>, Hideaki Motoyama<sup>3</sup>, Hiroyuki Matsuzaki<sup>4</sup>

1.Graduate School of Science and Technology, Hirosaki University, 2.The Atmosphere and Ocean Research Institute, The University of Tokyo, 3.National Institute of Polar Research, 4.The University Museum, The University of Tokyo

Centennial to millennial periodicities of past solar activity in the Holocene epoch have been well investigated by using cosmogenic-nuclides records from tree-ring ( $^{14}\text{C}$ ) and ice-core ( $^{10}\text{Be}$ ) archives. However, those of the older ages are quite unclear because high-resolution records of cosmogenic nuclides are scarce. We obtained multiple high-resolution  $^{10}\text{Be}$  records (atmospheric  $^{10}\text{Be}$  flux and authigenic  $^{10}\text{Be}/^9\text{Be}$  ratio) of 170–200 kyr ago, the period that includes the Iceland Basin geomagnetic excursion interval (Horiuchi et al., in press). Strong consistency among the records enables us to construct a robust  $^{10}\text{Be}$  stack of this interval. A wavelet analysis of the stacked record reveals 4-kyr and 8-kyr periodicities, both of which can be interpreted as intrinsic geomagnetic cycles. We also found a cycle of 1.7 kyr in the  $^{10}\text{Be}$  stack record, but it is significant only near the  $^{10}\text{Be}$  maximum (the minimum of the geomagnetic paleointensity). A relative enhancement of solar modulation in  $^{10}\text{Be}$  production is predicted with reduced paleointensity (Masarik and Beer, 1999; Beer et al., 2012). Therefore, it is possible to attribute this cosmogenic 1.7-kyr cycle to the long-term solar cycle that is identified in this period. The wavelet spectrum of the highest-resolution  $^{10}\text{Be}$  record (obtained from the Dome Fuji ice core) shows more a detailed structure than the stacked record. The 4- and 1.7-kyr cycles show similar tendencies to, and are clearer than, those of the stack. The spectrum also shows somewhat intermingled multi-centennial (0.3, 0.5 and 0.7 kyr) cycles around the maxima of  $^{10}\text{Be}$ , which likely represent solar cycles in this period (note: 200-yr Suess/de Vries cycle is under the detection limit of our record). On the other hand, the 2.2–2.4-kyr Hallstatt cycle and 1-kyr Eddy cycle, which are frequently documented in Holocene cosmogenic records, are not dominant in our records.

## Research of annual cosmic ray events using $^{10}\text{Be}$ in the Dome Fuji ice core

\*Fusa Miyake<sup>1</sup>, Kimiaki Masuda<sup>1</sup>, Kazuho Horiuchi<sup>2</sup>, Hideaki Motoyama<sup>3</sup>, Hiroyuki Matsuzaki<sup>4</sup>, Yuko MOTIZUKI<sup>5</sup>, Kazuya Takahashi<sup>5</sup>, Yoichi Nakai<sup>5</sup>

1.Institute for Space-Earth Environmental Research, Nagoya University, 2.Graduate School of Science and Technology, Hirosaki University, 3.National Institute of Polar Research, 4.MALT, University Museum, University of Tokyo, 5.RIKEN

Cosmogenic nuclides,  $^{14}\text{C}$  and  $^{10}\text{Be}$ , are produced in the atmosphere by cosmic rays from outside the Earth. Carbon-14 is stored in tree-rings, and beryllium-10 is stored in ice sheets. Then, we can investigate past cosmic ray intensities by analyzing concentrations of  $^{14}\text{C}$  or  $^{10}\text{Be}$ . Annual cosmic ray increase events (AD 775 and AD993 or AD994) were found in  $^{14}\text{C}$  data of tree-rings. These events were also shown in quasi-annual  $^{10}\text{Be}$  data in the ice cores from the Antarctica and the Greenlands. We analyzed quasi-annual  $^{10}\text{Be}$  concentration in the Dome Fuji ice core, and detected the cosmic ray events. In this presentaion, we will report the results of quasi-annual  $^{10}\text{Be}$  measurments, and discuss a comparison with  $^{14}\text{C}$  data.

Keywords: cosmogenic nuclide, annual cosmic ray event

## Micro-droplets containing sulfate in the Dome Fuji deep ice core, Antarctica: Findings using micro-Raman spectroscopy

\*Toshimitsu Sakurai<sup>1,2</sup>, Ohno Hiroshi<sup>3</sup>, Hideaki Motoyama<sup>1</sup>, Tsutomu Uchida<sup>4</sup>

1.National Institute of Polar Research, 2.Institute of Low Temperature Science, Hokkaido University, 3.Department of Civil and Environmental Engineering, Kitami Institute of Technology, 4.Graduate School of Engineering, Hokkaido University

Climatic signals in deep ice cores, particularly from ion concentrations, may be affected by the diffusion of liquid solution along grain boundaries of ice. Such solutions include sulfates. Because of the difficulty of detecting sulfate liquids in the ice matrix, we must infer the phase state of the sulfates from the ice temperature and inclusion properties. In this study, we use micro-Raman spectroscopy to determine the phase state of three sulfate micro-inclusions in the Dome Fuji ice core at 2798.5-m depth. Using a temperature-ramp test, we find a peak position at  $984\text{ cm}^{-1}$  and a change in the full width at half maximum (FWHM) of the S-O stretching mode that identifies the sulfate in the micro-droplets. Considering the peak position and FWHM of sulfate inclusions, we argue that the sulfate would have existed as a micro-droplet liquid on an air hydrate in the ice. Additionally, the increase in the low frequencies of the Raman spectrum below  $30\text{ cm}^{-1}$  that we detect can be generally used to identify liquids in natural ice. Our investigation also indicates that the surface of air-hydrates in ice is a preferred location for liquid micro-inclusions. The importance of this finding will be discussed.

Carbonate mineral particles observed in a layer of an ice core drilled from Grigoriev ice cap in Kyrgyz Tianshan, Central Asia

\*Nozomu Takeuchi<sup>1</sup>, Koji Fujita<sup>2</sup>, Vladimir Aizen<sup>3</sup>

1.Chiba University, 2.Nagoya University, 3.University of Idaho

In 2007, an 87 m-deep ice core were successfully drilled on Grigoriev Ice Cap (4600 m a.s.l.) located in the Tien Shan Mountains, Kyrgyzstan. We report a layer rich in carbonate mineral particles found at 53.5 m deep in the ice core. Although a number of dust layers consisting of silicate mineral particles were contained in the ice core, the layer contained less silicate, but abundant carbonate mineral particles. Significant negative stable isotope values and higher concentrations of major chemical solutes were also observed at the layers. Pollen based dating of the ice core showed that the layer corresponded to 1833 A.D. Results suggest that the layer was derived from a huge storm. Although the origin of the carbonate minerals is still mystery, it is probably far distant arid area.

Keywords: ice core, carbonate minerals, Central Asia

## Reconstructions of past cyanobacteria flora from ice core samples on Gregoriev Glacier, Kyrgyz Tianshan

\*Takahiro Segawa<sup>1,2</sup>, Takahiro Yonezawa<sup>3</sup>, Nozomu Takeuchi<sup>4</sup>

1.Research organization of information and systems, Transdisciplinary research integration center,  
2.National Institute of Polar Research, 3.Fudan University, 4.Chiba University

Cryoconite, a microbe-mineral aggregate found on glaciers worldwide is formed by the action of microbial phototrophs, principally Cyanobacteria.

The species composition of cyanobacteria in the ice cores could reflect the environmental condition at that time. Thus, these microorganisms in ice cores could be useful to reconstruct past environments. Despite the ecological importance in glacial environments, the phylogeographic distributions and genetic structures of glacial cyanobacteria are still highly limited.

We report results of cyanobacterial species and their evolution by molecular DNA analysis collected from the ice core samples collected on Gregoriev Glacier, Kyrgyz Tianshan. We reconstruct the organisms and their interactions within the community and with the environment on the sampled sites. We also present detailed pictures of cyanobacterial distribution patterns on glaciers over the Arctic, Antarctic, and Asian high mountains combination of 16S rRNA and 16S-23S internal transcribed spacer (ITS) regions. The results implied cyanobacteria could migrate across Asian glaciers for the last 10,000 years.

Keywords: glacier, ice core, DNA, cyanobacteria