

海洋生態系モデル相互比較計画

MARine Ecosystem Model Intercomparison Project (MAREMIP)

Doney Scott¹、*平田 貴文²Scott Doney¹, *Takafumi Hirata²

1.Woods Hole Oceanographic Institution、2.北海道大学地球環境科学研究所

1.Woods Hole Oceanographic Institution, 2.Faculty of Environmental Earth Science, Hokkaido University

Ocean biogeochemistry is strongly influenced by the specific activity of various types of plankton. In an effort to improve the representation of marine ecosystems, ocean biogeochemistry models have evolved to include a growing number of organisms aggregated according to their functionality into "Plankton Functional Types" (PFTs). Such models open up new and exciting avenues of research to explore interactions between marine ecosystems and climate change on various time scales. The "MARine Ecosystem Model Intercomparison Project" (MAREMIP) aims to foster the development of models based on PFTs in order to progress towards the resolution of important scientific questions; what are the impacts of global environmental changes on marine ecosystems, including climate change, ocean acidification and changes in nutrient input? Are there possible regime shifts associated with future environmental changes? What is the role of ecosystem structure and biodiversity for biogeochemical fluxes, marine resources and climate? In this talk, we show an overview of the MAREMIP activities and science highlights.

キーワード：海洋生態系、生態系モデル、相互比較

Keywords: Marine Ecosystem, Ecosystem Model, Intercomparison

Emergence of multiple ocean ecosystem drivers in a large ensemble suite with an Earth system model

*Keith B Rodgers¹, Jonathan Lin², Thomas Frölicher³

1.AOS Program, Princeton University, Princeton, USA, 2.Princeton University, Princeton, USA, 3.ETH, Zurich, Switzerland

Marine ecosystems are increasingly stressed by human-induced changes. Marine ecosystem drivers that contribute to stressing ecosystems –including warming, acidification, deoxygenation, and perturbations to biological productivity –can co-occur in space and time, but detecting their trends is complicated by the presence of noise associated with natural variability in the climate system. Here we use large initial-condition ensemble simulations with an Earth system model under a historical/RCP8.5 (representative concentration pathway 8.5) scenario over 1950-2100 to consider emergence characteristics for the four individual and combined drivers. Using a 1-standard-deviation (67% confidence) threshold of signal to noise to define emergence with a 30-year trend window, we show that ocean acidification emerges much earlier than other drivers, namely during the 20th century over most of the global ocean. For biological productivity, the anthropogenic signal does not emerge from the noise over most of the global ocean before the end of the 21st century. The early emergence pattern for sea surface temperature in low latitudes is reversed from that of subsurface oxygen inventories, where emergence occurs earlier in the Southern Ocean. For the combined multiple-driver field, 41% of the global ocean exhibits emergence for the 2005-2014 period, and 63% for the 2075-2084 period. The combined multiple-driver field reveals emergence patterns by the end of this century that are relatively high over much of the Southern Ocean, North Pacific, and Atlantic, but relatively low over the tropics and the South Pacific. For the case of two drivers, the tropics including habitats of coral reefs emerges earliest, with this driven by the joint effects of acidification and warming. It is precisely in the regions with pronounced emergence characteristics where marine ecosystems may be expected to be pushed outside of their comfort zone determined by the degree of natural background variability to which they are adapted. The results underscore the importance of sustained multi-decadal observing systems for monitoring multiple ecosystem drivers.

Keywords: Ocean biogeochemistry, Earth system modeling, Large ensemble

CMIP5モデルにおける海洋炭素ポンプの診断

Ocean carbon pumps in CMIP5 earth system models diagnosed by a vector diagram

*岡 顕¹*Akira Oka¹

1. 東京大学大気海洋研究所

1. Atmosphere and Ocean Research Institute, The University of Tokyo

The ocean stores 60 times more carbon than the atmosphere and therefore the ocean carbon cycle has a critical role in controlling the atmospheric CO₂ concentration. The ocean carbon cycle is controlled by several ocean pumps such as soft tissue (organic matter) and hard tissue (calcium carbonate) pumps. In the CMIP5 earth system models, these carbon pumps are explicitly simulated in the model and controls the level of the atmospheric CO₂ concentration. In this study, four types of ocean carbon pumps (organic matter, calcium carbonate, gas exchange, and freshwater flux pumps) are defined here and a method for diagnosing effects of individual four carbon pumps on atmospheric CO₂ concentration is proposed. In my method, the simulated 3-D field of dissolved carbon concentration (DIC), total alkalinity (ALK), phosphate, and salinity are used for diagnosing the strength of each carbon pump. In addition, the contributions of four carbon pump components to atmospheric CO₂ are evaluated in one figure (the vector diagram); each carbon pump component is represented by one vector and its contribution to pCO₂ can be measured from the difference in the contour values between the beginning and the end of the vector. The analysis is applied to the climatology and the CMIP5 earth system model simulations. Although all models reproduce the same level of the atmospheric CO₂ concentration as the climatology, it is shown that contributions from four carbon pumps are not the same among models. This study demonstrates that the vector diagram analysis introduced here is a useful tool for quantifying the individual effects of the ocean carbon pumps on atmospheric CO₂ concentration and also for evaluating the reproducibility of ocean carbon cycle models.

キーワード：炭素循環、海洋炭素ポンプ

Keywords: carbon cycle, ocean carbon pump

Nonlinear Interactions between Climate and Atmospheric Carbon Dioxide Drivers of Terrestrial and Marine Carbon Cycle Changes from 1850 to 2300


*Forrest M Hoffman¹, James Randerson², J Keith Moore², Michael L Goulden², Weiwei Fu², Charles D Koven³, Abigail LS Swann⁴, Natalie M Mahowald⁵, Keith Lindsay⁶, Ernesto Munoz⁶, Gordon B Bonan⁶

1.Oak Ridge National Laboratory, 2.University of California Irvine, 3.Lawrence Berkeley National Laboratory, 4.University of Washington, 5.Cornell University, 6.National Center for Atmospheric Research

Quantifying feedbacks between the global carbon cycle and Earth's climate system is important for predicting future atmospheric CO₂ levels and informing carbon management and energy policies. We applied a feedback analysis framework to three sets of Historical (1850-2005), Representative Concentration Pathway 8.5 (2006-2100), and its extension (2101-2300) simulations from the Community Earth System Model version 1.0 (CESM1(BGC)) to quantify drivers of terrestrial and ocean responses of carbon uptake. In the biogeochemically coupled simulation (BGC), the effects of CO₂ fertilization and nitrogen deposition influenced marine and terrestrial carbon cycling. In the radiatively coupled simulation (RAD), the effects of rising temperature and circulation changes due to radiative forcing from CO₂, other greenhouse gases, and aerosols were the sole drivers of carbon cycle changes. In the third, fully coupled simulation (FC), both the biogeochemical and radiative coupling effects acted simultaneously. We found that climate-carbon sensitivities derived from RAD simulations produced a net ocean carbon storage climate sensitivity that was weaker and a net land carbon storage climate sensitivity that was stronger than those diagnosed from the FC and BGC simulations. For the ocean, this nonlinearity was associated with warming-induced weakening of ocean circulation and mixing that limited exchange of dissolved inorganic carbon between surface and deeper water masses. For the land, this nonlinearity was associated with strong gains in gross primary production in the FC simulation, driven by enhancements in the hydrological cycle and increased nutrient availability. We developed and applied a nonlinearity metric to rank model responses and driver variables. The climate-carbon cycle feedback gain at 2300 was 42% higher when estimated from climate-carbon sensitivities derived from the difference between FC and BGC than when derived from RAD. These differences are important to quantify and understand because different model intercomparison efforts have used different approaches to compute feedbacks, complicating intercomparison of ESMs over time. Underestimating the climate-carbon cycle feedback gain would result in allowable emissions estimates that would be too low to meet climate change targets. We further explored the degree to which these nonlinearities affect climate-carbon cycle feedback gain estimates in CMIP5 models at year 2100.

Keywords: carbon cycle, feedbacks, Earth system model

Drivers of Hydrological and Ecological Changes (1850–2300)



Can we bet on negative emissions to achieve the 2°C target even under strong carbon cycle feedbacks?

Can we bet on negative emissions to achieve the 2°C target even under strong carbon cycle feedbacks?

*田中 克政¹、山形 与志樹¹、横畠 徳太¹、江守 正多¹、花岡 達也¹

*Katsumasa Tanaka¹, Yoshiki Yamagata¹, Tokuta Yokohata¹, Seita Emori¹, Tatsuya Hanaoka¹

1.国立環境研究所

1.National Institute for Environmental Studies

Abstract

Given the narrowing windows of opportunities to stay below 2°C, negative emission technologies such as Bioenergy with Carbon dioxide Capture and Storage (BioCCS) play an ever more crucial role in meeting the 2°C stabilization target (Fuss et al. 2014). Negative emission technologies –if deployable at a sufficiently large scale during the second half of this century –would make the 2°C target more feasible in the midst of the slow political progress. However, such technologies are currently at their infancy and their future penetrations may fall short of the scale required to stabilize the warming (Scott et al. 2013). Furthermore, the overshoot in the mid-century prior to a full realization of negative emissions would give rise to a risk because such a temporal but excessive warming above 2°C might amplify itself by strengthening climate-carbon cycle feedbacks, which are known to be positive albeit with large uncertainties (Friedlingstein et al. 2006). When one considers other classes of carbon cycle feedbacks including those with permafrost thawing and wildfire, such a risk could be even higher. It has not been extensively assessed yet how carbon cycle feedbacks might play out during the overshoot in the context of negative emissions, while the literature on carbon cycle feedbacks has burgeoned in recent years.

This study explores how 2°C stabilization pathways, in particular those which undergo overshoot, can be influenced by carbon cycle feedbacks and asks their climatic and economic consequences. We compute 2°C stabilization emissions scenarios under a cost-effectiveness principle, in which the total abatement costs are minimized such that the global warming is capped at 2°C. We employ a reduced-complexity model, the Aggregated Carbon Cycle, Atmospheric Chemistry, and Climate model (ACC2) (Tanaka et al., 2013), which comprises a box model of the global carbon cycle, simple parameterizations of the atmospheric chemistry, and a land-ocean energy balance model. The total abatement costs are estimated from the Marginal Abatement Cost functions for CO₂, CH₄, N₂O, and BC, which are derived from Azar (2013).

Our preliminary results show that, if carbon cycle feedbacks turn out to be stronger than what is known today, it would incur substantial abatement costs to keep up with the 2°C stabilization goal. Our results also suggest that it would be less expensive in the long run to plan for a 2°C stabilization pathway by considering strong carbon cycle feedbacks because it would cost more if we correct the emission pathway in the mid-century to adjust for unexpectedly large carbon cycle feedbacks during overshoot. Furthermore, our tentative results point to a key policy message: *do not rely on negative emissions to achieve the 2°C target*. It would make more sense to gear climate mitigation actions toward the stabilization target without betting on negative emissions because negative emissions might create large overshoot in case of strong feedbacks. Our simple approach illuminates a need for investigating this issue further by using a range of models including coupled Earth System Model (ESM)-Integrated Assessment Models (IAMs).

References

Azar C., et al. (2013) Meeting global temperature targets—the role of bioenergy with carbon capture

and storage. *Environmental Research Letters* 8 034004

Friedlingstein, P., et al. (2006) Climate-carbon cycle feedback analysis: results from the C4MIP model intercomparison. *Journal of Climate* 19 3337-3353

Fuss S., et al. (2014) Betting on negative emissions. *Nature Climate Change* 4 850-853

Scott V., et al. (2013) Last chance for carbon capture and storage. *Nature Climate Change* 3 105-111

Tanaka K., et al. (2013) Emission metrics under the 2°C climate stabilization target. *Climatic Change* 117 933-941

陸域統合モデル：陸面・水資源・作物・土地利用結合モデルの開発

Development of Integrated Terrestrial Model: a biogeophysical land surface model with human components

*横島 徳太¹、伊藤 昭彦¹、花崎 直太¹、櫻井 玄²、木下 嗣基³、飯泉 仁之直²、眞崎 良光¹、新田 友子⁴、Pokhrel Yadu⁵、江守 正多¹

*Tokuta Yokohata¹, Akihiko Ito¹, Naota Hanasaki¹, Gen Sakurai², Tsuguki Kinoshita³, Toshichika Iizumi², Yoshimitsu Masaki¹, Tomoko Nitta⁴, Yadu Pokhrel⁵, Seita Emori¹

1.国立環境研究所、2.農業技術環境研究所、3.茨城大学、4.東京大学大気海洋研究所、5.ラトガース大学
1.National Institute for Environmental Studies, 2.National Institute for Agro-Environmental Sciences, 3.Ibaraki University, 4.Atmosphere Ocean Research Institute, University of Tokyo, 5.Rutgers University

将来の気候変動は、食料生産・水資源・エネルギー・生態系サービスなどの要素に大きな影響を与えられ、それぞれの要素に与える影響は密接に関連するため、要素間の相互作用を考慮することが必要不可欠である。これまでの研究では、気候変動が食料・水・エネルギー・生態系のそれぞれに対して及ぼす影響の評価は行われてきたが、これら全体を陸域における自然-人間活動をシステムとして総合的にとらえて影響評価を行うことが、重要な課題である。そこで本研究では、気候変動が土地・水・農業・生態系に及ぼす影響を総合的に評価できる「陸域統合モデル」を開発する。モデルでは、全球気候モデルMIROC (Watanabe et al. 2010) の一要素である陸面モデルMATSIRO (Takata et al. 2003, Nitta et al. 2014) に、陸域生態系モデル VISIT (Ito and Inatomi 2012)、水資源モデルH08 (Hanasaki et al. 2008, Pokhrel et al. 2012)、作物モデルPRYSBI2 (Sakurai et al. 2015)、土地利用モデルTELMO (Kinoshita et al., in preparation) が結合されたモデルである。モデルでは、各サブモデルで計算された出力変数が、関係する別のモデルに数時間あるいは一日の時間ステップで渡され、時間発展する。たとえば、作物モデルPRYSBI2で計算された穀物収量は、土地利用モデルTELMOに渡され、翌年の土地利用変化が計算される。予報された土地利用変化は、すべてのサブモデルで利用される。また、水資源モデルH08では灌漑プロセス（河川からの取水）、ダム操作を考慮しており、その結果が陸面モデルの土壌水分や河川流量に影響を与える。発表では、モデル開発の現状と、過去再現実験および将来予測実験の結果を報告する。

キーワード：地球システムモデル、気候変動、人間活動

Keywords: Earth system model, climate change, human activity

全球土粒子輸送モデルを用いた河川内の土砂分布の時空間変動に関する研究

A study on spatial and temporal variability of sediment in rivers using global sediment transport model

向田 清峻¹、*芳村 圭¹

Kiyotaka Mukaida¹, *Kei Yoshimura¹


1.東京大学大気海洋研究所

1.Atmosphere and Ocean Research Institute, University of Tokyo

土砂の輸送量の研究は、個々の粒子の振る舞いの詳細な実験による観察やある対象地域における浸食量、堆積量を見積もるなど様々なスケールで行われてきた。個々の研究においては粒径分布や土地利用、土壌の種類など様々な要因について輸送量の時間変動への影響を示唆している。全球スケールにおいては、流域の大まかな気候区分や地形情報を用いて年々変動を表現する経験式を作成することが現在行われて来ている。それに対して本研究においては大河川において土砂動態の時間的変動、空間的変動の理解、再現、予測へとつなげていくために、現状をうまく説明する経験式を推定する立場とは別の、物理過程に従う物質の体積輸送により把握するという立場に立ったアプローチを取る。本研究においては全球スケールの土砂動態を表現するために、物理的過程に基づいて水の流下を計算する全球河川モデルCaMa-Flood (Yamazaki et al. 2011)に、浸食、運搬、堆積の現象に基づいた土砂の輸送過程を導入し、全球土砂輸送モデルCaMa-SEDを開発した。土砂生産過程においては、降水量と傾斜を用いて土砂生産量を推定し、モデルの中で土砂の運搬に関しては流水と同じように移流方程式を用いて掃流輸送と浮遊輸送の二つの輸送形態を計算過程として用い水平方向の移動を表現し、鉛直方向の移動としては河床への沈降量と河床からの巻き上げ量の変数を与えた。ストレージとして浮遊土砂量と堆積土砂量を逐次計算することによって各構成要素の輸送量への寄与を計算することができる。粒径別の計算結果によって河川の上流部から流下した土砂が輸送され粒径の大きいものから先に堆積していく作用による土砂輸送量の空間分布が計算可能となり、細粒分が河口での土砂量を支配していることを示した。生産、運搬、堆積の物理過程に基づいた計算を用いた感度実験の結果により、土砂輸送量に関して沈降速度の感度が高いことが分かった。河口への輸送量が多い粘土、シルト分において沈降速度が非常に小さいため一度浮遊した土砂が沈降しないことに由来する。定点観測データとの比較により、レジーム則や従来の全球河川モデルでは表現できなかった流量と濃度のピークタイミングのずれであるヒステリシス効果を表現することが可能になった。

キーワード：全球土砂輸送モデル、浮遊輸送、レジーム則

Keywords: global sediment transport model, suspended flow, sediment regime



陸域CO₂交換量推定の現状と課題

Current state of terrestrial CO₂ exchange estimations: progresses and remaining issues

*近藤 雅征¹、市井 和仁¹、Patra Prabir¹、佐伯 田鶴¹

*Masayuki Kondo¹, Kazuhito Ichii¹, Prabir Patra¹, Tazu Saeki¹

1. 海洋研究開発機構 地球表層物質循環研究分野

1. Department of Environmental Geochemical Cycle Research, Japan Agency for Marine-Earth Science and Technology

Terrestrial ecosystems play a critical role in formation of a feedback loop of carbon dioxide (CO₂) in atmosphere with atmospheric reservoir and climate, and thus directing a course of the future projection of climate change. The research community has spent significant efforts to understand behaviors of terrestrial ecosystems under a steady rise in atmospheric CO₂ concentration and temperature during the recent decades and deepen knowledge about the regional and global patterns of terrestrial CO₂ sinks and sources. estimate the terrestrial CO₂ exchange, while seeking consistency between simulated and observed CO₂ concentrations. The bottom-up approach estimates the terrestrial CO₂ exchange using ecosystem models, which simulate the ecosystem-scale carbon cycle by considering the internal biogeochemical mechanisms of carbon flows for each prescribed vegetation type and soil.

However, the current estimates of terrestrial CO₂ exchange by the bottom-up and top-down approaches remain inconsistent. As illustrated in the recent IPCC Assessment Report (AR5), the top-down approach tends to indicate stronger CO₂ sinks in temperate and boreal regions than the bottom-up approach does. Furthermore, the two approaches exhibited contrasting CO₂ sink-source patterns in the tropics; the bottom-up approach indicated CO₂ sinks and the top-down approach CO₂ sources. As illustrated by these inconsistencies, a consensus on the geographic distribution of the terrestrial CO₂ exchange has yet to be established among the research community.

In this study, we elaborate the current status and issues of terrestrial CO₂ flux estimations by the top-down and bottom-up approaches. Specifically, we compare the bottom-up estimate from dynamic global vegetation models that are forced by interannual variations of CO₂ concentration, climate and land use changes, with the top-down estimate from atmospheric CO₂ inversions. We show an improved level of agreement between the two estimates in relation to seasonal variability and, regional and global budgets, since the IPCC AR5. We also discuss the remaining issues causing inconsistency between the two estimates.

Acknowledgments

This research was supported by Environment Research and Technology Development Funds (2-1401) from the Ministry of the Environment of Japan and Asia-Pacific Network for Global Change Research (APN: grant#ARCP2011-11NMY-Patra/Canadell).

キーワード：陸域CO₂交換量、大気CO₂インバージョン、陸域生態系モデル

Keywords: Terrestrial CO₂ exchange, Atmospheric CO₂ inversion, Ecosystem model simulation

THE CARBON BALANCE OF THE TERRESTRIAL BIOSPHERE IN THE TWENTIETH CENTURY

*Sitch Alexander Stephen¹

1.University of Exeter

Sitch, S. (1) and the TRENDY DGVM consortium, (1) University of Exeter, UK.

Each year a consortium of Dynamic Global Vegetation Modelling groups perform a factorial set of global simulations over the historical period, 1901 –present, to investigate the temporal and spatial trends in the land sink, and the contribution of land-use to emissions. This activity contributes the annual global carbon budget updates of the Global Carbon Project. Typical around 10 models are forced with reconstructed observed climate, global atmospheric CO₂, gridded fields of historical land-use and land cover changes (LULCC), and nitrogen deposition for a subset of models which include a fully interactive nitrogen cycle. The TRENDY project will be presented, including process developments through to the latest Trendy-v4 (1901-2014). Results are used to ascertain the individual contribution of CO₂, Climate, Land-Use and N deposition on the regional and global land carbon sink. Increasingly offline land surface simulations and coupled ESM simulations use the same land-surface components and results from each can inform the other. Both TRENDY and C4MIP have increasing interest in evaluation activities. Furthermore, observational datasets including those from remote sensing are used to evaluate model performance and help constrain the global land carbon sink over the past two decades.

Keywords: land-atmosphere interactions, DGVMs, climate-carbon cycle models

地球システムモデルMIROC-ESMを用いた2倍増CO₂濃度下における気候-炭素循環の1000年数値積分
Climate-carbon cycle changes during 1000 years in doubled CO₂ concentration simulated by
MIROC-ESM

*羽島 知洋¹、立入 郁¹、河宮 未知生¹

*Tomohiro Hajima¹, Kaoru Tachiiri¹, Michio Kawamiya¹

1.独立行政法人 海洋研究開発機構

1.Japan Agency for Marine-Earth Science and Technology

Transient climate response to cumulative carbon emission, so called TCRE, is defined as the ratio of global warming to cumulative anthropogenic CO₂ emission evaluated when CO₂ concentration reaches the doubled CO₂ level from pre-industrial state. This metric is useful because it gives us roughly estimates on future global warming induced by CO₂ emission on the basis of current and future emission amounts. Since TCRE just characterizes the transient response of climate-carbon cycle, we cannot know what will happen after CO₂ concentration is stabilized (or reduced) after mitigation policies adopted. To estimate the warming degree in such condition and to understand climate-carbon dynamics in the concentration-stabilized phase, we conducted simulations where CO₂ concentration is abruptly doubled from pre-industrial state and fixed over 1000 years, by using an Earth system model (MIROC-ESM). We confirmed from the simulations that after 1000 years have passed, global warming and land carbon uptake almost ceased but weak carbon uptake by the ocean continues.

キーワード：炭素循環、気候変動、地球システムモデル、人為CO₂排出に対する気候の過渡的応答

Keywords: carbon cycle, climate change, Earth system model, TCRE

C4MIP simulations, plans and evaluation requirements

*Chris Jones¹

1. Met Office Hadley Centre

Climate-carbon cycle feedbacks are potentially large and play a leading order contribution in determining the atmospheric composition in response to human emissions of CO₂ and in the setting of emissions targets to stabilise climate or avoid dangerous climate change. For over a decade The Coupled Climate-Carbon Cycle Model Intercomparison Project (C4MIP) has coordinated coupled climate-carbon cycle simulations and in the coming few years C4MIP will be an endorsed activity of CMIP6. It is hoped that this will encourage widespread adoption of the C4MIP set of simulations and enable increased understanding and predictability of future changes in both terrestrial and marine carbon cycle.

C4MIP has 3 key strands of scientific motivation and the requested simulations are designed to satisfy their needs: (1) pre-industrial and historical simulations (formally part of the common set of CMIP6 experiments) to enable model evaluation; (2) idealised coupled and partially-coupled simulations with 1% per year increases in CO₂ to enable diagnosis of feedback strength and its components; (3) future scenario simulations to project how the Earth System will respond over the 21st century and beyond to anthropogenic activity.

In this talk I will outline some previous C4MIP results and present some key priorities for evaluation. It is clear that in biogeochemical modelling and the drive for increased complexity in ESMs, process-based model evaluation has not kept pace. As a result there is very large quantitative spread between CMIP5 carbon cycle results which hinder their usefulness. It is also the case that we have not been able to show demonstrable progress - as a coherent community - in the quality and process-realism of our modelling. There are no agreed quality criteria or metrics which measure whether our ESMs are fit for purpose or if they have improved since the last generation. It is essential that we focus our efforts in the coming years on addressing this deficiency. It is not enough that under CMIP6 there are more models within C4MIP analyses or more advanced processes. We must be able to demonstrate that we have made real progress since CMIP5 in our modelling skills, analysis techniques and our ability to constrain future projections.

There are multiple ways of evaluating carbon cycle models. Activities such as TRENDY and OCMIP (part of OMIP) will perform evaluation activities of offline land and ocean components respectively. It is the role of C4MIP to evaluate the coupled climate-carbon cycle system. Our primary simulations for this activity will be the coupled historical simulations from 1850 up to 2014. There will be two variants. Within the CMIP "DECK" (the central core of CMIP6) all models will perform a "concentration driven" historical run. This means the atmospheric concentration of CO₂ is prescribed to follow the historical record. The second variant, which is required for all models contributing to C4MIP is a parallel "emissions driven" historical simulation in which CO₂ emissions are prescribed to the model and the models simulate the time evolution of CO₂ concentration.

In order to fully exploit these simulations we need to be prepared with some top-level evaluation criteria (e.g. as presented by Anav et al 2013); some rigorous process-based criteria and metrics (such as sensitivity of stores and fluxes to environmental drivers); carefully assembled and processed observational datasets; carefully defined model diagnostic outputs. Here I will briefly outline these requirements in the hope of stimulating discussion to move our plans forward ahead of

model simulations being started by the end of 2016.

Keywords: Carbon cycle, CMIP, evaluation

温暖化に伴う海洋炭素循環の長期変動

Millennium-scale changes in ocean carbon cycle under global warming

*山本 彬友¹、阿部 彩子¹、山中 康裕²

*Akitomo Yamamoto¹, Ayako Abe-Ouchi¹, Yasuhiro Yamanaka²

1.東京大学大気海洋研究所、2.北海道大学 大学院地球環境科学研究所

1.Atmosphere and Ocean Research Institute, The University of Tokyo , 2.Graduate School of Environmental Science, Hokkaido University

海洋は大気中に放出された人為起源二酸化炭素を1000-2000年かけて吸収する。その為、海洋炭素循環の応答を理解することは、長期的な大気CO₂濃度を予測する上で重要である。一方、温暖化に対するSST、海洋循環と生物ポンプの応答は海洋のCO₂吸収量を減少させると考えられている(気候-炭素循環フィードバック)。簡略化モデルを用いた先行研究ではSSTの上昇と海洋循環の変化が気候-炭素循環フィードバックの主要因とされてきた(Sarmiento et al, 1998; Plattner et al, 2001; Matsumoto et al, 2010)。しかしながら、GCMと近年の比較的複雑なbiogeochemical modelを用いて、これらについて評価した研究はほとんどない。

本研究ではGCM(MIROC3.2)とNPZDモデルをベースとしたoffline biogeochemical model(Yamamoto et al, 2015)を用いて温暖化実験を2000年積分し、海洋のCO₂吸収量と気候-炭素循環フィードバックを計算した。更に感度実験を行い、フィードバックに寄与するメカニズムをそれぞれ切り分けて再評価した。

海洋のCO₂吸収量と、温気候-炭素循環フィードバックの割合は先行研究と同程度の値が得られた。一方、感度実験の結果から、気候-炭素循環フィードバックを引き起こす主要因はSSTの上昇と生物ポンプの変化となり、先行研究とは異なる結果になった。生物ポンプについては、新生産の減少と水温上昇に伴う再無機化速度の上昇により表層200mのDIC濃度を増加させる効果がCO₂吸収の減少に重要であることが分かった。

発表では海盆別のCO₂吸収量の時系列変化や、海洋循環の変化の影響についても発表する予定である。

キーワード：海洋炭素循環、2000年温暖化実験、気候-炭素循環フィードバック

Keywords: ocean carbon cycle, multi-millennium simulation, climate-carbon cycle feedback

Global-scale swamp forest modelling using satellite-based elevation and forest datasets

*Dai Yamazaki¹

1.JAMSTEC Japan Agency for Marine-Earth Science and Technology

Swamp forest (seasonally flooded forest) has different characteristics to usual non-flooded forest in terms of carbon pool and nutrient flux. Swamp forest exists in wide climate regions from boreal to tropical regions, but its global distribution is not well studied because satellite observation of inundated forest floor is disturbed by vegetation canopy. Here, we propose a method to estimate the global distribution of swamp forest by global river model simulations using satellite elevation and forest datasets. Digital Elevation Models (DEMs) based on satellite radar or stereoscopy have elevation biases due to forest canopy that impede the simulation of swamp forest hydrodynamics. We removed the elevation biases by combining ICESat forest height data and Landsat forest density data. The simulation with the bias-corrected DEM shows good agreement to L-band radar observation of swamp forest inundation in the Amazon basin, which suggests the potential of the proposed method for estimating swamp forest distribution.

Keywords: Swamp Forest, Global Model, Flooding

CMIP6へ向けた日本の地球システムおよび気候モデリング研究に関するレビュー
Earth system and climate modeling activities toward CMIP6 in Japan: a review

*河宮 未知生¹

*Michio Kawamiya¹

1. 海洋研究開発機構

1. Japan Agency for Marine-Earth Science and Technology

国内において3つのグループが、第6期結合モデル相互比較プロジェクト(CMIP6)への貢献を予定している。すなわち、気象庁気象研究所単独チーム、東京大学大気海洋研究所(AORI)、国立環境研究所、海洋研究開発機構(JAMSTEC)で構成されるTeam MIROC、AORI、理研、JAMSTECで構成されるNICAM開発チームである。上記のうち初めの2グループは種々のモデルバージョンを用いながらCMIP6で承認されたほとんどのモデル相互比較プロジェクト(MIPs)に参加する一方、NICAM開発チームは要求計算資源の大きさから少数のMIPsを選んで参加することになる。2015年に更新され1.3PFlopsの理論性能を持つ地球シミュレータがCMIP6に向けた実験の多くに用いられるが、NICAMを用いた一部の実験には「京」も活用される。国内におけるCMIP6関連活動の多くは文部科学省による「気候変動リスク情報創生プログラム」によって支援されているが、当該プログラムは2016年度いっぱい終了することが予定されている。それ以降の気候予測研究の展開について、関連省庁も交えながら研究者コミュニティでの検討が始まっている。

キーワード：地球システムモデル、気候予測、CMIP6

Keywords: Earth system model, Climate projection, CMIP6

MIROC-ESMにおける葉面積指数バイアスの気候への影響

Effect of bias in Leaf Area Index to climate in MIROC-ESM

*立入 郁¹、羽島 知洋¹

*Kaoru Tachiiri¹, Tomohiro Hajima¹

1. 海洋研究開発機構

1. Japan Agency for Marine-Earth Science and Technology

Leaf Area Index (LAI) is not only an important factor to determine the amount of photosynthesis, but also a key variable to represent ground surface condition which gives significant impact to atmosphere through biophysical processes such as albedo and transpiration. In this study, we first compared the present-day LAI in Earth System Models (ESMs) with remote sensing data and confirmed the significant overestimation by ESMs, which is pointed out by some existing studies. In order to investigate the effect of the bias, we next compared two pre-industrial control experiments: an ordinary control experiment in which LAI is prognosed by the terrestrial ecosystem component of the ESM, and an experiment in which LAI is replaced with dataset made from remote sensing data (the pre-industrial state was estimated by retuning the cultivated areas to natural vegetation). The result shows a significant temperature difference between the two cases, and by analysis of the related variables we concluded that the difference is caused by albedo change due to the change in snow depth/coverage in May-June and in the sea ice coverage in winter.

キーワード：葉面積指数、バイアス、MIROC-ESM、気候

Keywords: Leaf Area Index, bias, MIROC-ESM, climate

Socio-economic implications of stabilization scenarios from MIROC-ESM

*Diego Silva¹, Kaoru Tachiiri¹

1. Project Team for Risk Information on Climate Change, Japan Agency for Marine-Earth Science and Technology (JAMSTEC)

Setting a target for stabilization of climate in the long-term requires significant reductions in greenhouse gas (GHG) emissions at global scale. This involves important transformations in the society, basically affecting the future patterns of energy consumption and production, as well as the patterns of land use. Moreover, the cost for achieving such climate target may be considerable, thus, requiring an optimal allocation of efforts that minimizes the economic impact. This study presents the socio-economic implications of emission scenarios aiming at long-term climate stabilization, estimated with an integrated assessment model (IAM). Emission scenarios are obtained from the earth system model (ESM) Model for Interdisciplinary Research on Climate (MIROC-ESM). The outcomes on supply and demand of energy, land use, and mitigation costs are presented.

The emissions scenarios considered are consistent with the representative concentration pathway (RCP), and aim at a global radiative forcing by 2100 of around 4.5 W/m^2 (RCP4.5) and 2.6 W/m^2 (RCP2.6). The Global Change Assessment Model (GCAM) is applied to study the developments in energy, land use and emissions throughout the 21st century. GCAM is an IAM based on a partial equilibrium approach, which resolves the balance in supply and demand across the energy, land use and agricultural sectors.

Compared to the standard RCPs, the emission scenarios from MIROC-ESM presented lower levels of allowable anthropogenic CO_2 emissions for the same climate target. This is an outcome of the stronger feedback between the carbon-cycle and the climate, and the higher value for climate sensitivity assumed in MIROC-ESM in contrast to the climate model used in the development of RCPs. As a consequence, the changes in the energy and land systems are more drastic, while the cost of mitigations is higher. These differences are greater in the second half of the century.

Keywords: MIROC-ESM, stabilization scenario, integrated assessment model

南大洋域における海洋生態系へ供給される鉄起源としての鉱物エアロゾル

Mineral aerosol as a source of iron for the marine ecosystems in the Southern Ocean

*伊藤 彰記¹、Kok Jasper²

*Akinori Ito¹, Jasper F. Kok²

1.海洋研究開発機構、2.カリフォルニア大学ロサンゼルス校

1.Japan Agency for Marine-Earth Science and Technology, 2.University of California, Los Angeles

大気からダストにより供給される鉄は、食物連鎖を通して海洋生態系へ影響を与える。特に、南大洋域は溶存鉄が植物プランクトンの生育にとって制限要因となっており、その広大さから全球炭素循環を理解する上で極めて重要である。しかし、南半球におけるダスト発生量の推定値には多大な不確実性がある。本研究では、全球大気化学輸送モデルに最新のダスト発生スキームを導入することによって、観測値（鉄濃度など）を再現可能な鉄発生量を推定する。

キーワード：エアロゾル、生物地球化学的循環、大気化学輸送モデル

Keywords: Aerosols , Global Biogeochemical Cycles, Atmospheric chemistry transport model

On constraining the strength of the terrestrial CO₂ fertilization effect in an Earth system model

*Vivek Arora¹, John Scinocca¹

1.Canadian Centre for Climate Modelling and Analysis, Environment and Climate Change Canada

Earth system models (ESMs) explicitly simulate the interactions between the physical climate system components and biogeochemical cycles. Physical and biogeochemical aspects of ESMs are routinely compared against their observation-based counterparts to assess model performance and to evaluate how this performance is affected by ongoing model development. Here, we assess the performance of version 4.2 of the Canadian Earth system model against four, land carbon cycle focused, observation-based determinants of the global carbon cycle and the historical global carbon budget over the 1850-2005 period. Our objective is to constrain the strength of the terrestrial CO₂ fertilization effect which is known to be the most uncertain of all carbon cycle feedbacks. The observation-based determinants include (1) globally-averaged atmospheric CO₂ concentration, (2) cumulative atmosphere-land CO₂ flux, (3) atmosphere-land CO₂ flux for the decades of 1960s, 1970s, 1980s, 1990s and 2000s and (4) the amplitude of the globally-averaged annual CO₂ cycle and its increase over the 1980 to 2005 period. The optimal simulation that satisfies constraints imposed by the first three determinants yields a net primary productivity (NPP) increase from ~ 58 Pg C yr⁻¹ in 1850 to about ~ 74 Pg C yr⁻¹ in 2005; an increase of ~ 27 % over the 1850-2005 period. The simulated loss in the global soil carbon amount due to anthropogenic land use change over the historical period is also broadly consistent with empirical estimates. Yet, it remains possible that these determinants of the global carbon cycle are insufficient to adequately constrain the historical carbon budget, and consequently the strength of terrestrial CO₂ fertilization effect as it is represented in the model, given the large uncertainty associated with LUC emissions over the historical period.

Keywords: Earth system models, CO₂ fertilization effect, Land carbon uptake

北太平洋中層の海洋酸性化についてのマルチモデル解析

Multi-model analysis of ocean acidification in the subsurface layers of the North Pacific

*渡辺 路生¹、河宮 未知生¹*Michio Watanabe¹, Michio Kawamiya¹

1. 海洋研究開発機構

1. Japan Agency for Marine-Earth Science And Technology

Ocean acidification is one of the major threats for marine calcifying organisms such as precious corals. They live in the mesophotic waters at a depth of 80–300 m near Japan and are known as habitat-forming species with an important structural role in mesophotic habitats of continental slopes, sustaining biodiversity hotspots. Using the outputs of projections under the highest emission scenario of the Representative Concentration Pathways performed by Earth System Models (ESMs), we evaluate the ocean acidification rates in the middle layers of the North Pacific, where the strongest sink of atmospheric CO₂ is found in the mid-latitudes. The mixed layer depth in the Kuroshio Extension region reaches ~200 m during winter due to the strong wind forcing and cooling. Consequently, the low potential vorticity (PV) water mass called the Subtropical Mode Water is formed. This mode water shows large dissolved inorganic carbon (DIC) concentration increase, and is advected southwestward, so that, in the Izu-Ogasawara region, DIC concentration increases and ocean acidification occurs faster than in adjacent regions. The ESMs simulate that pH in the middle layers of the Izu-Ogasawara region decreases by 0.3–0.4 from 2006 through 2100. We find that the ESMs with a deeper mixed layer during winter in the Kuroshio Extension region show the large increase in DIC concentration within the Izu-Ogasawara region. For a reliable projection of the ocean acidification in the middle layers of the Izu-Ogasawara region, an ESM should well reproduce the mixed layer deepening during winter in the Kuroshio Extension region.

キーワード：海洋酸性化、CMIP5、地球システムモデル

Keywords: ocean acidification, CMIP5, Earth System Model

活性汚泥および脱窒菌からの亜硝酸ガス (HONO)の発生測定の発生測定

Measurements of Gaseous Nitrous Acid (HONO) Emission from Activated Sludge and Denitrifying Bacteria

*峰島 知芳¹、宮下 智之¹

*Chika Minejima¹, Tomoyuki Miyashita¹

1.国際基督教大学

1.International Christian University

Introduction

Gaseous nitrous acid (HONO) is known as a precursor of OH radicals, a strong oxidant in the atmosphere. Up to 34 % of OH radical is produced from HONO in a city and a rural area¹⁾. Therefore HONO is an important species to know OH radical behaviors.

The high HONO concentration have been observed during daytime in spite of the HONO photolysis.

There are several known HONO sources: gas phase reactions, heterogeneous reactions and combustion process. Also, HONO emission from soil by the equilibrium between gaseous nitrous acid and aqueous nitrous acid in the soil and the direct emission by nitrifying bacteria have been observed²⁾³⁾. In the soil, there are not only nitrifying bacteria but denitrifying bacteria. However the emission by denitrifying bacteria is not studied.

The research purpose is to determine whether or not denitrifying bacteria in the activated sludge emits HONO directly.

Experimental

HONO emissions from activated sludge in aerobic condition and anaerobic condition were measured. Also HONO emission from the sterilized supernatant solution was measured. HONO emissions from biological process and chemical process were compared. Activated sludge in Duran bottle was purged with air or N₂ for 1 day to 4 days and HONO was captured with a filter pack. The sludge was aerobic with air purge and anaerobic with N₂ purge. Duran bottle and filter packs were covered with tin foil to avoid HONO photodissociation. Dissolved Oxygen was measured to keep the condition of activated sludge and pH was stabilized at 7.8-8.1 by adding 0.1 M HCl solution or 50 g/L NaHCO₃ solution not to decrease the bacteria's activity. The flow speed was controlled at 2 L/min with mass flow controllers.

Also, activated sludge and its sterilized supernatant solution were purged with room air and N₂ at 22 °C. Also the activated sludge was purged with N₂ at 27, 32 °C. The activated sludge in a Duran bottle was put in the water which is controlled with a heat controller to stabilize the temperature of activated sludge. In these experiments, three filter packs were placed for each experiment. Air or N₂ was purged in 4.5 L/min because three filter packs were prepared and the purged gas was controlled to flow through each filter pack in 1.5 L/min. The NO₂⁻ concentrations of the sludge and supernatant were measured before these experiments with the pack test. The purging time was 8 hours in order to keep controlling pH.

Results and Discussion

In this experiment, HONO emissions from the sludge were observed at anaerobic condition. The contribution of biological process was more than 90%. However, calculated activation energy from temperature dependent experiment was much bigger than that of the denitrifying bacteria. Thus, there is a possibility that denitrifying bacteria reduce NO₃⁻ to NO₂⁻ and the increase in NO₂⁻ concentration increased the HONO emission from the chemical process. Also, NO₂⁻ concentration in the sludge should be measured more accurately because it has influence on HONO emission.

Reference

- 1) Wong *et al.*, *Atmos. Chem. Phys.*, **11**, 3595-3609 (2011).
- 2) Su, *et al.*, *Science*, **333**, 1616-1618 (2011).
- 3) Oswald, *et al.* *Science* **341**, 1233-1235 (2013).

キーワード: nitrous acid, atmosphere, soil

Keywords: nitrous acid, atmosphere, soil

IMPACT OF BIOGEOPHYSICAL AND BIOGEOCHEMICAL PROCESSES AND THEIR INTERACTIONS ON PERMAFROST SOIL CARBON STOCKS

IMPACT OF BIOGEOPHYSICAL AND BIOGEOCHEMICAL PROCESSES AND THEIR INTERACTIONS ON PERMAFROST SOIL CARBON STOCKS

*Jain Atul¹, Shu Shijie¹, Barman Rahul¹, El-Masri Bassil¹

*Jain K Atul¹, Shijie Shu¹, Rahul Barman¹, Bassil El-Masri¹

1.Department of Atmospheric Science, University of Illinois

1.Department of Atmospheric Science, University of Illinois

One of the major challenges in more detailed Earth system models (ESMs) is the treatment of the biophysical and biogeochemical processes and feedbacks and their impact on soil organic carbon (SOC), in the Northern high latitudes (NHL) permafrost region. This is because a larger suite of active terrestrial processes coupled with scarcity of observational data introduce many challenges for modelling these processes. Nonetheless, several studies in the recent past have demonstrated improved permafrost modelling capabilities by incorporating soil/snow processes that critically influence the ground energetics in these environments, such as: deep soil layers and organic soils, and the effects of wind compaction and depth hoar formations, and structural properties of vegetation (phenology, assimilated vegetation C allocation to leaves, stems, and roots and root dynamics) However, no study has yet evaluated the combined effects of the improvements of these biogeophysical and biogeochemical processes for the entire NHL.

We use a land surface model, the Integrated Science Assessment Model (ISAM), to investigate the effects of feedbacks between the biogeochemical (C and N) and biogeophysical (water, and energy) processes on the model estimated soil organic carbon (SOC) for the NHL permafrost region. We not only focus on recent model improvements in the biogeophysical processes that are deemed important for the high latitude soils/snow and permafrost SOC; such as deep soil column, modulation of soil thermal and hydrological properties, wind compaction of snow, and depth hoar formation; but also biogeochemical processes that are important for soil biogeochemistry; such as dynamic phenology and root distribution, litter carbon decomposition rates and nitrogen amount remaining. We select multiple sites to evaluate the modeled processes. We then carried out several model simulations to study the effects of feedbacks between biogeochemical and biogeophysical processes on SOC in NHL.

After accounting for dynamic biogeochemical processes, ISAM is able to capture permafrost extent and the carbon stored in NHLs, as well as the seasonal variability in leaf area index (LAI), and root distribution in the soil layers and the root response of soil water uptake and transpiration. The evaluation of the model results suggest that without accounting these processes, modelled growing season length (GSL) for NHL was almost two times higher as compared to measurements. To quantify the implication of these processes on the carbon and water fluxes, we compared the results of two different versions of ISAM, dynamic version which accounts for dynamic processes (ISAM_{DYN}) and static version which do not account for dynamic processes (ISAM_{BC}), with measurements from 12 eddy covariance flux sites. The results show that ISAM_{DYN}, unlike ISAM_{BC}, is better able to capture the seasonal variability in GPP and energy fluxes. Our modelling analysis shows that by including the biophysical processes in addition to biogeochemical processes, the modelled NHL permafrost carbon increased by 30% from 328 to 447 GtC in the top one meter of soil which is in better agreement with observational estimates of 495 GtC (Northern Circumpolar Soil Carbon Database). Even though the inclusion of these processes generally reduced vegetation

productivity and litter production due to a decrease in soil temperature and liquid water content, increased soil carbon stocks highlight the dominance of soil water/temperature stress on decomposition processes. While continued improvements are required in the treatment of biogeochemistry, here we demonstrate the importance of soil/snow biogeophysical and biogeochemical processes in modelling permafrost carbon stocks, as important drivers of soil biogeochemical processes.

キーワード：Permafrost Soil Carbon、Land Surface Model、Biogeophysics、Biogeochemistry
Keywords: Permafrost Soil Carbon, Land Surface Model, Biogeophysics, Biogeochemistry