

## Mechanisms and Predictability for Arctic Sea-Ice Variability with the MIROC Climate Model

\*Jun Ono<sup>1</sup>, Hiroaki Tatebe<sup>1</sup>, Yoshiki Komuro<sup>1</sup>, Masato I Nodzu<sup>2</sup>, Masayoshi Ishii<sup>2</sup>

1.Japan Agency for Marine-Earth Science and Technology, 2.Meteorological Research Institute

The Arctic environment has markedly changed due to the rapid decline of sea ice in summer Arctic Ocean. The retreat of sea-ice cover could be associated with the Arctic amplification and an increase in the frequency of severe cold winters in the Northern Hemisphere mid-latitudes. Accurate predictions of sea-ice variability on seasonal to decadal time-scales and its mechanisms would be useful for further progress in science as well as socio-economic activity. To examine the mechanisms and predictability for Arctic sea-ice variability, we analyze the control simulations from the Arctic Predictability and Prediction On Seasonal to Inter-annual Timescales (APPOSITE) project. The model used for the APPOSITE is the climate model MIROC 5.2 in which external forcing is fixed in 2005. The time evolution in sea-ice extent and volume shows that an extreme reduction event occurs one or two for a century even without the global warming trend. The spatial feature in sea-ice distribution and its reduction mechanisms resemble those in 2007. This suggests that an anomalous sea-ice loss could be caused by only natural variability. We have currently investigated the key factors closely related to the sea-ice variability.

Keywords: Arctic, Sea ice, Climate model

## Interannual variability of sea ice production in the Barrow Coastal Polynya off Barrow, Alaska

\*Daisuke Hirano<sup>1</sup>, Yasushi Fukamachi<sup>2</sup>, Eiji Watanabe<sup>3</sup>, Katsushi Iwamoto<sup>4</sup>, Kay I. Ohshima<sup>2</sup>, Takeshi Tamura<sup>1</sup>

1.National Institute of Polar Research, 2.Institute of Low Temperature Science, Hokkaido University, 3.Japan Agency for Marine-Earth Science and Technology, 4.Mombetsu City

Recently, it has been proposed that the Barrow Coastal Polynya (BCP), previously considered to be a latent heat polynya, is a "wind-driven" hybrid latent/sensible heat polynya, with both features caused by the same northeasterly wind (*Hirano et al.*, 2016). In this study, we examine the interannual variability of sea ice production in the BCP from 2002/03 to 2010/11 during AMSR-E operation period, with focus on the northeasterly wind that characterizes the BCP as a hybrid coastal polynya. Throughout the ice-growth season (November–May), sea ice production was the highest in 2003/04 and the lowest in 2010/11. In 2003/04, amount of the suppressed sea ice production was also highest, when ~30% of the BCP ice production was suppressed by the ocean heat transport associated with warm water upwelling. Wind pattern around the BCP region varies from year to year, and frequency and magnitude of the northeasterly wind correlate well with sea level pressure difference between the Beaufort High and Aleutian Low. Compared with climatology, the northeasterly wind in the BCP was more frequent and stronger in 2003/04 due to strengthening of the Beaufort High. In contrast, it was less frequent and weaker in 2010/11 due to weakening of both the Beaufort High and Aleutian Low. Frequency and magnitude of the northeasterly wind, mainly regulated by variabilities of the Beaufort High and Aleutian Low, are considered to be major factors of the interannual variability of sea ice production in the BCP.

Keywords: hybrid latent/sensible heat polynya, sea ice production, interannual variability

A modeling study on water modification and its interannual variability in the Barents Sea

\*Takao Kawasaki<sup>1,2</sup>

1.National Institute of Polar Research, 2.Atmosphere and Ocean Research Institute, The University of Tokyo

The sea ice is not found in the southern half of the Barent Sea even in winter due to the warm Atlantic Water inflow through the Barents Sea Opening. Recent several studies have shown the significant relationship between sea ice extent in the Barents Sea and winter air temperature in some mid latitude regions of northern hemisphere. Therefore, it is important to investigate the inflow and modification of the Atlantic Water in the Barents Sea for getting a better understanding of climate change in mid-latitude regions. In this study, we utilize a high resolution realistically configured ice-ocean general circulation model to examine the mechanism of water modification and its interannual variability.

The modeled routes of the Atlantic Water are affected by the oceanic bottom topography and consistent with observations and previous modeling studies. The cooling and freshening of the Atlantic Water by the atmosphere and sea ice melting, respectively, are also well simulated. Calculated heat flux at the Barents Sea Opening is  $\sim 87\text{TW}$  ( $1\text{ TW} = 10^{12}\text{ Watt}$ ) is in range of observational estimates ( $73\text{-}103\text{TW}$ ). The sea ice formation at the coastal polynya, which contributes the increasing of salinity in the Barents Sea, is slightly underestimated compared with satellite observation. The interannual variability of Atlantic Water modification and its mechanism are currently under investigation.

Keywords: Barents Sea, Ice-Ocean General Circulation Model, Sea surface heat flux

## Change in persistent extratropical regimes under an Arctic amplified climate

\*Masato Mori<sup>1</sup>, Masahiro Watanabe<sup>1</sup>, Masahide Kimoto<sup>1</sup>

1. Atmosphere and Ocean Research Institute, University of Tokyo

The Arctic surface air temperature has warmed more than twice as fast as the global average (e.g., Cohen et al. 2014), which is known as Arctic Amplification (AA). All the fifth Coupled Model Intercomparison Project (CMIP5) model projects that the warming become more and more conspicuous toward the end of this century, which is one of the most robust climate change signal projected by the models. Therefore, it is important to clarify the extent to which the AA influences the Northern Hemisphere mid-latitudes extreme events, especially recurrent and persistent circulation pattern which causes the heat wave and cold spell.

Here we use a 100-member ensemble of historical simulations and future projections with a hi-resolution atmospheric general circulation model to show that as a result of change in the climatological atmospheric flow induced by the AA, the probability of occurrence of a specific circulation anomaly pattern will increase in future. This circulation pattern is strongly tied to winter cold spell over the Northern Hemisphere mid-latitudes in present climate, but not necessarily in the Arctic amplified future climate. This is because a reduced climatological meridional temperature gradient in lower troposphere acts to weaken the variance of surface temperature.

Keywords: Arctic region, Arctic Amplification, extreme event

## Changes in precipitation over the Arctic projected by global atmospheric models with 20-km and 60-km grid sizes

\*Shoji Kusunoki<sup>1</sup>

1.Meteorological Research Institute

A set of global warming projections was conducted using global atmospheric models with high-horizontal resolution of 20-km (MRI-AGCM3.2S, the 20-km model) and 60-km (MRI-AGCM3.2H, the 60-km mode) grid sizes. For the present-day climate (1983-2003, 21 years), models were forced with observed historical sea surface temperatures (SST). For the future climate (2079-2099, 21 years, RCP8.5), models were forced with future SST distributions projected by the models of the Fifth phase of Couple Model Intercomparison Project (CMIP5). The uncertainty of projection was evaluated by ensemble simulations for four different SST distributions and three different cumulus convection schemes.

The annual mean precipitation (PAVE), the simple daily precipitation intensity index (SDII), and the maximum 5-day precipitation total (R5d) averaged over the Arctic increased in the end of the 21st century. The increases in PAVE, SDII, and R5d can be partly attributed to an increase in water vapor associated with increasing temperatures, and to an increase in the horizontal transport of water vapor from low to high latitudes. These results are consistent with Kusunoki et al. (2015).

Keywords: Arctic, Global warming projection, Global atmospheric model, Precipitation

## Simulation on low pressure systems over the Arctic Ocean using a cloud-resolving model

\*Taro Shinoda<sup>1</sup>, Masaya KATO<sup>1</sup>, Jun Inoue<sup>2,3</sup>, Hirohiko Masunaga<sup>1</sup>, Kazuhisa Tsuboki<sup>1</sup>

1.Institute for Space-Earth Environmental Research, Nagoya University, 2.National Institute of Polar Research, 3.Japan Agency for Marine-Earth Science and Technology

Nagoya University continues to develop Cloud Resolving Storm Simulator (CReSS). CReSS is applied to analyze the structure of precipitation systems in the tropics and mid-latitude regions. However, we have never conducted the simulation around the Arctic region, except for a case of the cloud-streak structure over the Labrador Sea during a cold-air outbreak. The R/V Mirai (MR13-06) of Japan Agency for Marine-Earth Science and Technology (JAMSTEC) made an intensive observation using a Doppler radar, upper-air soundings, and instruments for surface meteorological parameters at a fixed observation point (168.25W, 72.75N) in the Arctic Ocean in 2013. To confirm the performance of CReSS in the Arctic region, we conducted a simulation to reproduce low pressure systems observed by the R/V Mirai.

A numerical experiment using CReSS is examined. The horizontal grid spacing is 2.5 km, and the domain has an area of 2000 km times 2000 km, including the fixed observation point of the R/V Mirai. The vertical grids are consisted of 32 layers and the top of the domain is set at 12.8 km. The numerical experiment is conducted for 72 h, starting at 00Z on September 23, 2013. Data of the Global Spectral Model (GSM: Horizontal grid resolution was 0.5 degree) provided by Japan Meteorological Agency (JMA) are used as the initial and boundary conditions of the simulation. Sea surface temperature (SST) is initialized using the Optimum Interpolation Sea Surface Temperature (OISST) data provided by National Oceanic and Atmospheric Administration (NOAA). Vertical heat transfer below the land and sea surfaces are calculated, as a result, time variation of sea surface temperature (SST), sensible and latent heat fluxes (SHF, LHF) from the surface are calculated. No sea ice parameterization is included.

A synoptic scale low pressure accompanying weak precipitation (snow) less than 1 mm/h is reproduced in the north of Wrangel Island. A mesoscale low pressure around the synoptic scale one is confirmed by a satellite observation. The simulation can reproduce a vortex-like structure related to the mesoscale low pressure. Using the Mirai Doppler radar, deep convective clouds whose echo-top height reaches a height of 4 km are observed around the mesoscale low. However, the cloud-top height in the simulation is restricted only below a height of 1.5 km, thus the deep convective clouds cannot be reproduced. To compare the simulation result with the satellite observation, 3-dimensional distribution of the reflectivity in the frequency of the CloudSat-CPR (95 GHz) is calculated using Satellite Data Simulator Unit (SDSU). The low reproducibility of the depth of convection is confirmed by the composite analysis using 20 flight paths. High reflectivity greater than 0 dBZ with the convective structure is observed by the CloudSat-CPR, however, the simulation cannot reproduce the high reflectivity region.

Time series of SST at the R/V Mirai between the observation and simulation are analyzed. At the initial time, OISST are 0.2 degree Celsius lower than that of the observation. SST is almost constant in the observation during the period, however, that of the simulation decreases gradually by the SHF and LHF. As a result, SST difference increases 1.0 degree Celsius at the end of the simulation. The constant SST would be attributed to the northward horizontal advection of warm water by southerly surface wind and heat loss by the SHF and LHF. CReSS cannot reproduce the wind-driven ocean flow, thus we fail to reproduce time series of SST. The failure of the SST at the R/V Mirai influences on the LHF from the sea surface and unrealistic saturated atmospheric boundary layer (fog) in the last 24 hours. The weak forcing from the sea surface would contribute to the

failure of the development of deep convection. The air-sea interaction and three-dimensional ocean simulation would be important to reproduce the meteorological phenomena in the Arctic region.

Keywords: Arctic region, air-sea interaction, a cloud-resolving model, mesoscale polar low

## A Comparison between the mid-Holocene and the future in the Arctic warming mechanism

\*Marina Suzuki<sup>1</sup>, Masakazu Yoshimori<sup>2,3</sup>

1.Graduate School of Environmental Science, Hokkaido University, 2.Faculty of Environmental Earth Science, Hokkaido University, 3.Arctic Research Center, Hokkaido University

From the climate model simulations, it is projected that the Arctic warming will be strengthened in the future.

However, the degree of projected warming varies with climate models and there are substantial uncertainties in the future projections of Arctic warming.

Shmidt et al.(2013) suggested a possibility of constraining uncertainty in the future climate projections by using information from past periods when the Arctic region was warmer than today. In their study, however, only a statistical correlation between paleoclimate and future climate simulations across models is presented. Therefore, the underlying mechanism causing the correlation is unclear. As they are well aware, the sound physical explanation is necessary in order for us to apply their method to constrain the future climate projections.

The principle aim of this study is to investigate commonality and difference of Arctic warming mechanism in the past and in the future, and to obtain insight into the possibility of constraining uncertainty of future climate projections by using paleo-climate data. We use the outputs of pre-industrial control, the quadruple of atmospheric CO<sub>2</sub> concentration(4xCO<sub>2</sub>) and mid-Holocene experiments from 11 CMIP5/PMIP3 atmosphere-ocean general circulation models. The mid-Holocene here refers to 6000 years ago and the Arctic region was warmer than today because of the difference in earth's orbital parameters.

First, we identified predominant processes of Arctic warming in each experiment based on the surface energy balance, and also investigated sea ice, clouds, water vapor and sea surface temperature response. In both experiments, most of the anomalous energy input into the Arctic region in summer is used for melting of sea ice, absorbed into the ocean. Consequently, the surface warming is moderate during summer. Several months later, the heat is released from the exposed warm sea surface and it causes the Arctic warming. In addition, the role and approximate timing of sea ice, clouds and water vapor changes are common in the two experiments. We found that there are many commonality in the Arctic warming mechanism in the future and in the mid-Holocene even though they are caused by completely different external forcing.

Next, we quantified that contribution of individual processes to the inter-model variance of the surface temperature changes in the Arctic region for each experiment. In the 4xCO<sub>2</sub> experiment the largest contribution to the variance of annual mean (ANN) surface temperature change is the surface albedo feedback. As to the October - December (OND) mean surface temperature change, on the other hand, it is the heat release from the ocean. In the mid-Holocene experiment the largest contributions to the variance of both ANN and OND surface temperature change in the Arctic region are made by the differences in downward clear-sky longwave radiation. Also, the contribution of the surface albedo feedback to the variance of ANN surface temperature changes and the contribution of the oceanic heat release to the variance of OND surface temperature changes are relatively large and statistically significant. As the surface temperature and near-surface air temperature which determines the downward longwave radiation are tightly coupled, it is expected that constraints of other processes would result in a reduction of total uncertainty. In other words, if the Arctic warming in the mid-Holocene is simulated accurately, the reliability of the model's representation of surface albedo feedback, the process related to oceanic heat release, and hence the future projections would increase.



Based on the understanding of Arctic warming mechanism from this study, it is considered that paleo-environment information of Arctic Warming in the mid-Holocene is useful for constraining uncertainty in the future projections of Arctic warming.

Keywords: Climate models, Paleoclimate, Future projections

## Climate changes in Ny-Ålesund and Longyearbyen, Svalbard based on long-term meteorological and terrestrial dataset

\*Tomotaka Saruya<sup>1</sup>, Tetsuo Sueyoshi<sup>1</sup>, Hiroyuki Enomoto<sup>1</sup>

### 1. National Institute of Polar Research

Arctic has been undergoing various climatic and environmental changes in associated with global warming. Temperature increase rate is almost double that the rate of global average, which causes the melting of permafrost, the reduction of sea ice. Since the climatic system in the Arctic consists of the interactions between atmospheric circulation, hydrological process, and sea ice variation, the understanding of atmosphere-land (ice)-ocean feedbacks is required to reveal the environmental changes due to global warming. Therefore long-term and multi-point monitoring of atmospheric, ocean and terrestrial components are required.

Svalbard is representative area for Arctic research where various institutes have been performing long-term monitoring. Longyearbyen and Ny-Ålesund in Spitsbergen that are located in far from human activity are better suited for monitoring of minor variation components and Arctic climate changes. National Institute of Polar Research (NIPR) and Alfred-Wegener Institute (AWI) have monitored climatic components since early 1990s using Automated weather stations. In Longyearbyen, The University Centre in Svalbard (UNIS) has monitored meteorological and permafrost components at Adventdalen with boreholes and weather stations. In addition to research institutes, Norwegian Meteorological Institute (NMI) has monitored snow cover and precipitation at Svalbard Airport and Ny-Ålesund since 1970s. We have compared various components of climatic system such as air temperature, ground temperature, precipitation, snow cover at several locations using NIPR, AWI, UNIS and NMI dataset. As a result of analysis of time-series variations, some notable trends in air temperature and precipitation were found. Winter temperatures at Longyearbyen and Ny-Ålesund are increased during the last two decades, however, temperatures at March remain mostly unchanged or decrease. As for the precipitation, precipitation amount at Ny-Ålesund remain unchanged, while that at Longyearbyen decreases. In this presentation, we discuss the long-term trend and correlation of meteorological and permafrost components.

Keywords: climate change, Svalbard

## Revisiting impacts of spring Eurasian snow cover change on the East Asian summer precipitation

Subaru Fujiwara<sup>1</sup>, \*Toru Nozawa<sup>1</sup>

1. Graduate School of Natural Science and Technology, Okayama University

The Eurasian snow cover extent (SCE) anomaly in spring has been considered as one of the important factors affecting East Asian summer monsoon (e.g. Wu and Kirtman, 2007; Wu et al., 2009; Yim et al., 2010). In these studies, the authors analyzed traditional SCE dataset of National Oceanic and Atmospheric Administration (NOAA). Recently, Japan Aerospace Exploration Agency (JAXA) has developed a new long-term SCE product using Advanced Very High Resolution Radiometer (AVHRR) and Moderate Resolution Imaging Spectroradiometer (MODIS) data spanning from 1980's to 2014. This new product (JAXA/SCE) has higher spatial resolution and smaller commission error compared with NOAA/SCE. Continuity of the algorithm is another strong point in JAXA/SCE. Here, we revisit impacts of spring Eurasian snow cover change on the East Asian summer precipitation by using the new JAXA/SCE dataset. Climatological mean fields of spring SCE is not largely different from each other. On the other hand, interannual variability of spring SCE has somewhat different spatial distribution over the Eurasian region (45°N-70°N, 20°E-140°E); NOAA/SCE shows a dipole pattern but JAXA/SCE shows monopole pattern. We will present analyzed results on relationships between spring SCE anomaly over the Eurasia and the East Asian summer rainfall anomaly.

## Endurance of larch forest ecosystems in eastern Siberia under warming trends

\*Hisashi Sato<sup>1</sup>, Hideki Kobayashi<sup>1</sup>, Go Iwahana<sup>2</sup>, Takeshi Ohta<sup>3</sup>

1.Department of Environmental Geochemical Cycle Research, Japan Agency for Marine-Earth Science and Technology (JAMSTEC), 2.International Arctic Research Center, University of Alaska Fairbanks, 3.Graduate School of Bioagricultural Sciences, Nagoya University

The larch (*Larix* spp.) forest in eastern Siberia is the world's largest coniferous forest. Its existence is considered to be depending on near-surface permafrost, which increases water availability for trees, and the boundary of the forest closely follows the permafrost zone. Therefore, the degradation of near-surface permafrost due to forecasted warming trends during the 21st century is expected to affect the larch forest in Siberia. However, predictions of how warming trends will affect this forest vary greatly, and many uncertainties remain about land-atmospheric interactions within the ecosystem.

We developed an integrated land surface model to analyze how the Siberian larch forest will react to current warming trends. This model analyzed interactions between vegetation dynamics and thermo-hydrology and showed that, under climatic conditions predicted by the IPCC's RCP scenarios 2.6 and 8.5, annual larch net primary production (NPP) increased about 2 and 3 times, respectively, by the end of 21st century compared with that in the 20th century. Soil water content during larch growing season showed no obvious trend, even after decay of surface permafrost and accompanying sub-surface runoff. A sensitivity test showed that the forecasted warming and pluvial trends extended leafing days of larches and reduced water shortages during the growing season, thereby increasing productivity.

Keywords: Siberian Larch Forest, Climatic Change, Vegetation Model

## Tree-ring and modeling estimates for tree response to climate change over circumpolar forest ecosystems

\*Shunsuke Tei<sup>1,2</sup>, Atsuko Sugimoto<sup>2</sup>, Maochang Liang<sup>3</sup>, Yojiro Matsuura<sup>4</sup>, Akira Osawa<sup>5</sup>, Hitoshi Yonenobu<sup>6</sup>, Hisashi Sato<sup>7</sup>, Trofim Maximov<sup>8,9</sup>

1.National Institute of Polar Research, 2.Hokkaido University, 3.Yangtze University, 4.Forestry and Forest Products Research Institute, 5.Kyoto University, 6.Naruto University of Education, 7.Japan Agency for Marine-Earth Science and Technology, 8.Institute for Biological Problem of Cryolithozone, 9.North-Eastern Federal University

Arctic and boreal ecosystems are exposed to rapid and strong increases in temperature and related environmental changes under Arctic amplification. Yet, there is uncertainty how trees in these ecosystems respond to the changes due to an insufficiency of such long term records and this is where tree-rings can provide an advantage. Early dendrochronological studies in the region focused on the positive growth of trees to warmth (D'Arrigo and Jacoby, 1993, *Clim. Change*). However, A number of more recent studies have demonstrated a reduced sensitivity of tree growth to rising temperatures (now referred to as "divergence problem") at least since the 1960s (e.g., Wilson et al., 2007, *J. Geophys. Res*). Although several studies (e.g., Barber et al., 2000, *Nature*) suggested that temperature-induced drought may limit tree growth under the limited availability of soil moisture, the underlying processes for the phenomenon are not well understood.

We here investigated past tree response to climate changes, especially to warming, using retrospective analyses from tree-ring width and carbon isotope ratios ( $\delta^{13}\text{C}$ ) of three genera (*Larix*, *Picea* and *Pinus*) in 6 forest sites with a strong gradient of temperature and precipitation, reaching from northern Europe to northern America; Kalina (59N, 27E), Yakutsk (62N, 129E), Ust'Maya (60N, 133E), Chokurdakh (70N, 148E), Inuvik (68N, 133W) and Fort Smith (60N, 112W). The results suggest that tree response to past climate changes have varied with regions. The tree responses to warming are negative in eastern Siberia forests, resulting in decreasing trend of tree growth over past 60 years. On the other hand, the negative effect of warming is not seen in European and Canadian forests, where no decrease trend of growth is observed. The results then have been used in testing a dynamic global vegetation model (SEIB-DGVM, Sato et al., 2007, *Ecol. Model*). The simulated annual net primary productions (NPP) show no decreasing trend over the study period and discrepancy from tree-ring based long-term (more than half-decadal) growth variations in eastern Siberian forests, although relatively better reproductions of the model for the variations are obtained in European and Canadian forests.

The observed discrepancy in eastern Siberian forest may become more severe for future projection. We developed a climate-driven statistical growth equation that uses regional climate variables to model tree-ring width values for each site and then applied these growth models to predict how tree growth will respond to twenty-first-century climate change (RCP8.5 scenario). Although caution should be taken when extrapolating past relationships with climate into the future, we observed future continues reduction of the growth in central part of eastern Siberia, which is opposite trend from the DGVM based estimate. Our results imply that the negative effect of warming override the expected positive effects i.e., warming-induced lengthened growing season and increase in photosynthetic ratio, in arid region such as eastern Siberia, suggesting further reduction of tree growth by future warming, and no reproduction of the negative effect in the DGVM seems to be a cause for the observed discrepancy between tree-ring and DGVM estimates. The negative effect of warming for tree growth is a key process for accurate future projection of ecosystem functions and therefore further field and modeling investigations are essential to deep understanding of the

underlying processes for the phenomenon.

## Spatial distribution pattern of willows in Indigirka river lowland of northeast Siberian Arctic

\*Tomoki Morozumi<sup>1</sup>, Rong Fan<sup>1</sup>, Ryo Shingubara<sup>1</sup>, Shinya Takano<sup>1</sup>, Ruslan Shakmatov<sup>1</sup>, Shunsuke Tei<sup>2</sup>, Hideki Kobayashi<sup>3</sup>, Rikie Suzuki<sup>3</sup>, Trofim C Maximov<sup>4,5</sup>, Atsuko Sugimoto<sup>6,1</sup>

1.Graduate School of Environmental Science, Hokkaido University, 2.NIPR, 3.JAMSTEC, 4.IBPC SB RAS, 5.BEST center, NEFU , 6.Faculty of Earth Environmental Science, Hokkaido University

A warming climate and longer growing season may lead higher primary production and expansion of deciduous tundra shrubs in the arctic. Willows and alders are dominant plants in riparian ecosystem, which is frequently disturbed by river flooding. Because of their high productivity, carbon assimilation in this ecosystem is expected to be important. However, willows distribution and primary production have not be fully understood yet. Willows dominate large area of the floodplain of Yana-Indigirka-Kolyma river lowland in northeast Siberian Arctic. We investigated distribution patterns and NDVI of willows in the Indigirka River floodplain using satellite image classification and GIS.

High resolution vegetation map and satellite derived NDVI were obtained from a WorldView-2 satellite image (10 x 10 km) based on the field observations in July 2013 near Chokurdakh (70°N, 148°E). The willow distribution pattern corresponding to a distance from the river was analyzed. The willow covers 1/6 of lowland in local scale, and showed large extent along mainstream of the Indigirka River. This implies that large area along the main stream is affected by spring flooding, and willows cover this area. Besides, the area covered by willows showed the highest NDVI among the vegetation classes. These results mean the willow vegetation along the main stream and tributaries may greatly affect the primary production in river lowland of northeast Siberian Arctic.

Keywords: vegetation, floodplain, NDVI, GIS

## Methane Oxidation Potential of Arctic Wetland Soil of a Taiga-Tundra Ecotone in Northeastern Siberia

\*Jun Murase<sup>1</sup>, Atsuko Sugimoto<sup>2,3</sup>, Ryo Shingubara<sup>2</sup>, Trofim C Maximov<sup>4</sup>

1.Graduate School of Bioagricultural Sciences, Nagoya University, 2.Graduate School of Environmental Science, Hokkaido University, 3.Faculty of Environmental Earth Science, Hokkaido University, 4.Institute for Biological Problems of Cryolithozone

Arctic wetlands are significant sources of atmospheric methane and the observed accelerated warming of the arctic causes increased methane formation in water-saturated tundra soil with deepened permafrost thawing. Methane oxidation is the key process to regulate methane emission from wetlands. In this study we determined the potential methane oxidation rate of the wetland soils of a Taiga-Tundra transition zone in Northeastern Siberia. Peat soil samples were collected in the summer from depressions that were covered with tussocks of sedges and *Sphagnum* spp. and from mounds vegetated with moss and larch trees. The potential methane oxidation rate was estimated by a bottle incubation experiment in which homogenized soil samples were incubated with methane at the initial concentration of 0.5-0.8 %v/v. Soil samples collected from depressions in the moss- and sedge-dominated zones exhibited active methane oxidation with no lag. The potential methane oxidation rates at 15 °C ranged from 270 to 190 nmol h<sup>-1</sup> g<sup>-1</sup> dw. Methane oxidation was active over the depths including the water-saturated anoxic layers. The maximum methane oxidation rate was recorded in the layer above the water-saturated layer: the surface (0-2cm) layer in the sedge-dominated zone and in the middle (4-6 cm) layer in the moss-dominated zone. The methane oxidation rate was temperature-dependent and the threshold temperature of methane oxidation was estimated to be -4 to -11 °C, which suggested methane oxidation at subzero temperatures. Soil samples collected from the frozen layer of *Sphagnum* peat also showed immediate methane consumption when incubated at 15 °C. The present results suggest that methane oxidizing bacteria keep their activity in the wetland soils even under anoxic and frozen conditions and immediately utilize methane when the conditions become favorable. On the other hand, difluoromethane, the inhibitor of methane oxidation, did not alter the methane flux from the sedge and moss vegetation, indicating the undetectable levels of methane oxidation associated with the peat plants.

Keywords: Methane Oxidation, Peat, Permafrost soil, Oxygen



Multi-year response of CH<sub>4</sub> efflux to wetting at Indigirka Lowland in Northeastern Siberia

\*Ryo Shingubara<sup>1</sup>, Atsuko Sugimoto<sup>2,1</sup>, Jun Murase<sup>3</sup>, Shunsuke Tei<sup>4,2</sup>, Shinya Takano<sup>1</sup>, Tomoki Morozumi<sup>1</sup>, Maochang Liang<sup>1,7</sup>, Go Iwahana<sup>2,8</sup>, Trofim C. Maximov<sup>5,6</sup>

1.Grad. Sch. Envir. Sci., Hokkaido Univ., 2.Fac. Earth Envir. Sci., Hokkaido Univ., 3.Grad. Sch. Bioagr. Sci., Nagoya Univ., 4.NIPR, 5.IBPC SB RAS, Russia, 6.BEST center, NEFU, Russia, 7.Yangtze Univ., China, 8.IARC, UAF, USA

Under the amplified Arctic warming climatic response of CH<sub>4</sub> emission from the wetlands needs to be understood and predicted because of possible influence to the global climate. Indigirka Lowland in Northeastern Siberia has wetlands in a taiga-tundra boundary on permafrost, whose ecosystem are possibly sensitive to the climate change. Though environmental controls on CH<sub>4</sub> efflux have been found such as water level (soil moisture), soil temperature and vegetation, the quantitative relationship between the controls and CH<sub>4</sub> efflux are still unclear, which depends on region and timescale (Olefeldt et al., 2013, Global Change Biol.; Treat et al., 2007, JGR). One difficulty is that CH<sub>4</sub> emission is composed of 3 processes, i.e. CH<sub>4</sub> production, oxidation and transport; they can respond to environmental controls and affect CH<sub>4</sub> efflux in different ways. These processes are reflected by stable isotope ratios of CH<sub>4</sub> (delta-<sup>13</sup>C-CH<sub>4</sub>, delta-D-CH<sub>4</sub>), which can associate field observation and knowledge from laboratory incubation experiments on CH<sub>4</sub> production and on oxidation.

In this study we assessed interannual variation in chamber CH<sub>4</sub> efflux and in delta-<sup>13</sup>C-, delta-D-CH<sub>4</sub> near Chokurdakh (70.62 N, 147.90 E) over summers of 2009-2013 to understand relationship between CH<sub>4</sub> efflux and environmental factors based on the 3 processes of CH<sub>4</sub>.

CH<sub>4</sub> efflux was around the detection limit at dry tree mounds through the observation period, while large interannual variation was observed at wet areas of sphagnum moss and sedges. Wet event concurrent with the highest precipitation occurred in 2011 and CH<sub>4</sub> efflux increased at wet areas in the same year. Although water level decreased from 2011 to 2013, large CH<sub>4</sub> emission continued. Moreover, dissolved CH<sub>4</sub> concentration in soil pore water (at 10-15 cm depth) increased by 1 order of magnitude from 2011 to 2012 and kept high till 2013. CH<sub>4</sub> isotopes implies that CH<sub>4</sub> oxidation was depressed in 2012 after the wetting in 2011, suggesting soil reduction induced by the wetting proceeded over multiple years, which may have affected dissolved CH<sub>4</sub> concentration and CH<sub>4</sub> efflux. Such variation in CH<sub>4</sub> efflux and in dissolved CH<sub>4</sub> concentration will be discussed in relation to the 3 processes in this presentation.

Keywords: methane flux, interannual variation, isotope ratio, taiga-tundra boundary

## Occurrence frequency of ice and snow accretion in the boreal forest regions, Fairbanks and Yakutsk

\*Konosuke Sugiura<sup>1,2</sup>, Shin Nagai<sup>2</sup>, Rikie Suzuki<sup>2</sup>, Hajo Eicken<sup>3</sup>, Trofim Maximov<sup>4</sup>

1.University of Toyama, 2.Japan Agency for Marine-Earth Science and Technology, 3.International Arctic Research Center, University of Alaska Fairbanks, 4.Institute for Biological Problems of Cryolithozone Siberian Branch of RAS

Generally snow covers the ground, even if the ground type is different in such as tundra and taiga. Erosion/deposition of the snow cover and growth/decay of ice and snow accretion to vegetation occasionally occur, and the surface albedo changes and will affect the atmosphere. Previous research has demonstrated the high variations of the surface albedo in winter/spring in snow-covered forest regions in various global climate models. In this study, we focused on the surface albedo over snow-covered forest regions, and carried out field observations to verify the occurrence frequency of ice and snow accretion in the boreal forest regions. Using interval cameras installed on the observation tower at the site located to the north of Fairbanks (USA) and on the observation tower at the site located to the north of Yakutsk (Russia), ice and snow accretion in the boreal regions were monitored. It was found that the boreal forest at the Yakutsk site is covered with snow in comparison with the boreal forest at the Fairbanks site for a long term. After calculating using a one-dimensional mathematics model about the energy flow including atmospheric multiple scattering, it was shown that the mean surface temperature rises approximately 0.5 [K] when the boreal forest is not covered with snow. In this presentation, the snow albedo parameterization and the one-dimensional mathematics model are discussed to contribute to a better understanding of the role of snow in the climate system.

Keywords: ice accretion, snow accretion, albedo, boreal forest

## Vegetation masking effect on snow albedo feedback in Siberia during future global warming simulated by MIROC-ESM

\*Manabu Abe<sup>1</sup>, Kumiko TAKATA<sup>2</sup>, Michio Kawamiya<sup>1</sup>, Shingo Watanabe<sup>1</sup>

1.JAMSTEC, 2.NIPR

We have investigated future change in snow amount and vegetation masking effect on snow albedo feedback (SAF) in Siberia, boreal forests region of the northern Eurasia, in the future climate change simulation by Earth System Model, MIROC-ESM. Under the future scenario, RCP8.5, snow amount during fall-spring over the northern Eurasia decreases due to global warming. The significant reduction of snow amount is found in western Eurasia during fall-spring. On the other hand, although reduction of snow amount over Siberia in fall and spring occurs, winter snow amount over Siberia increases due to increasing snow fall, which is attributable to more water vapor with higher air temperature. Relating to such snow changes, then, surface air temperature (SAT) changes are enhanced through SAF. During spring, particularly, future SAT increases largely over Siberia, boreal forest region, although snow cover decreases less than that in western Eurasia. The dominant increase in SAT over Siberia is attributed to strong SAF which is caused by both reduced snow-covered surface albedo and reduced snow cover fraction. Further, to evaluate an effect of future LAI change on the surface albedo reduction, we have conducted an additional future climate change simulation, in which change in LAI is not included. The comparison between the future climate changes with and without the LAI changes suggests growing vegetation in the future may be a potential factor of the future strong warming through the vegetation masking effect on snow-covered surface albedo change.

Keywords: snow albedo feedback, vegetation masking effect, Earth system model

## Snow cover properties observed in Indigirka lowland near Chokurdakh, Northeastern Siberia

\*Shinya Takano<sup>1</sup>, Shunsuke Tei<sup>2,3</sup>, Tomoki Morozumi<sup>1</sup>, Atsuko Sugimoto<sup>1,2</sup>, Roman Petrov<sup>4,5</sup>, Trofim Maximov<sup>4,5</sup>

1.Graduate School of Environmental Science, Hokkaido University, 2.Faculty of Environmental Earth Science, Hokkaido University, 3.National Institute of Polar Research, 4.Institute for Biological Problems of Cryolithozone, SB RAS, 5.BEST center, NEFU, Yakutsk, Russia

Quantifying the spatial and temporal variations in snow depth, density, and snow water equivalent (SWE) is essential for many applications in hydrology and ecology. Snow survey including observation on the water isotope ratios of snow was conducted in Indigirka lowland near Chokurdakh (70.62 N, 147.90 E), Northeastern Siberia. Isotopic composition of water is powerful tool for investigation of hydrological processes such as discerning of source water for river discharge, ground ice, etc. The purposes of this study are (1) to know the spatial variations in snow depth, density, SWE and stable isotopic composition in this area, and (2) to estimate SWE in areal or regional scale, by scaling-up based on topographic and vegetative controls on SWE.

Snow survey was conducted in April 2014 and April 2015. Two transects from Chokurdakh to south and southwest, which are approximately 40 km and 20 km in length respectively, were set, and observation and sampling were made at 7 points and 4 points in 2014, respectively, and 12 points on the 40 km transect in 2015. In addition, snow survey was conducted at 25 points in 2014 (24 points in 2015) in total in the area measured approximately 1.2 km east to west at site K where various observations are conducted for taiga-tundra boundary ecosystem. The ranges of snow depth, density, SWE and  $\delta^{18}\text{O}$  in this area observed in 2014 were 30 to 90 cm, 0.137 to 0.318 g/cm<sup>3</sup>, 70 to 200 mm and -36.5 to -22.9‰, respectively, whereas those observed in 2015 were 12 to 83 cm, 0.131 to 0.325 g/cm<sup>3</sup>, 20 to 160 mm and -31.2 to -22.8‰, respectively. Although the values and the ranges were slightly different between 2014 and 2015, observed snow cover properties depended on vegetation type and showed consistencies: snow cover was the deepest at the site covered by dense and tall shrub, while snow density was the highest on ice over a lake. The SWE was the highest at shrub site, whereas that was the lowest at the site of sedge and/or sphagnum wetland. Spatial variation in delta-values of snow was observed, however there was no correlation with vegetation type, snow depth and snow density. Since clear relationship between SWE and vegetation type, SWE was estimated using a data on fraction of each vegetation obtained from a vegetation map drawn with high resolution satellite data (world view 2) and in situ observation (Morozumi et al., in preparation). The local average SWE values in observation area (10 x10 km) were estimated to be 100 mm in 2014 and 78 mm in 2015.

Keywords: snow water equivalent, water isotope ratios, Northeastern Siberia

## Surface elevation change of outlet glaciers in northwestern Greenland

\*Shun Tsutaki<sup>1,2</sup>, Naoki Katayama<sup>2,3</sup>, Daiki Sakakibara<sup>2,3</sup>, Shin Sugiyama<sup>2</sup>, Takanobu Sawagaki<sup>4</sup>

1.Arctic Environment Research Center, National Institute of Polar Research, 2.Institute of Low Temperature Science, Hokkaido University, 3.Graduate School of Environmental Science, Hokkaido University, 4.Faculty of Social Science, Hosei University

Ice discharge from calving glaciers has been increasing in the Greenland ice sheet (GrIS) since 2000s. This increase plays important roles in the volume change of GrIS and its contribution to sea level rise. To investigate the mass loss of GrIS calving glaciers, ice surface elevation change has been studied by differencing digital elevation models (DEMs) derived by satellite remote-sensing. Panchromatic Remote-sensing Instrument for Stereo Mapping (PRISM) images of Advanced Land Observing Satellite (ALOS) have a spatial resolution of 2.5 m, which is fine enough to measure several meters of elevation change on glaciers. The large spatial coverage of the images (1225 km<sup>2</sup>) is suitable for studying glaciers distributed over a large area.

In this study, we measured surface elevation change of 14 outlet glaciers near Inglefield Bredning in northwestern Greenland (77°47'–78°10'N, 65°00'–72°47'W). These glaciers flow into the ocean except for two land-terminating glaciers. We processed stereo pair ALOS PRISM images acquired in summer 2007 and 2010 with a digital map plotting instrument (Leica Photogrammetry Suite) to generate DEMs with a 25 m grid mesh. Elevation data from ALOS DEMs were calibrated on ice-free terrain, and compared to calculate ice surface elevation change between 2007 and 2010.

The surface elevation of all the studied glaciers decreased and the magnitude of the elevation change increases downglacier. The mean elevation change rate ranged from  $-0.4$  to  $-4.9$  m a<sup>-1</sup>. Marine-terminating Tracy and Bowdoin Glaciers thinned at rates of  $-4.9$  and  $-4.1$  m a<sup>-1</sup>, which were larger than those at other glaciers. The rate at Tugto Glacier, a land-terminating glacier located near Bowdoin Glacier, was  $-2.8$  m a<sup>-1</sup>. This result confirms that recent thinning of GrIS outlet glaciers is more significant at marine-terminating glaciers as compared to land-terminating glaciers. Rapid thinning of marine-terminating outlet glaciers observed in this study suggests the importance of ice dynamics and/or ice-ocean interaction in the mass loss of GrIS.

Keywords: Arctic glaciology, Glacier thinning, Glacier dynamics

## Water properties and circulation in front of tidewater glaciers in northwestern Greenland

\*Masahiro Minowa<sup>1,2</sup>, Shin Sugiyama<sup>1</sup>, Yoshihiko Ohashi<sup>1,2</sup>, Takanobu Sawagaki<sup>3</sup>, Shun Tsutaki<sup>4</sup>, Daiki Sakakibara<sup>2,1</sup>, Shigeru Aoki<sup>1</sup>, Evgeny Podolskiy<sup>5</sup>, Yvo Weindmann<sup>6</sup>

1.Institute of Low Temperature Science, Hokkaido University, 2.Graduate School of Environmental Science, Hokkaido University, 3.Faculty of Social Science, Hosei University, 4.Arctic Environment Research Center, National Institute of Polar Research, 5.Arctic Research Center, Hokkaido University, 6.The Laboratory of Hydraulics, Hydrology and Glaciology

Tidewater glaciers in Greenland are rapidly retreating, and it contributed global sea-level rise during the twentieth century. Studies have been carried out to understand the mechanisms of the glacier retreat. Previous studies suggested the importance of submarine melting, but physical processes relevant to submarine melting (e.g. heat source of the melting, water circulation and bathymetry) are not understood well. This is because in-situ observations are difficult in front of a glacier where icebergs usually cover the ocean. In addition to this, there is no hydrographical observation in front of tidewater glacier in northwestern Greenland. To better understand the ice-ocean interaction, we measured ocean temperature, salinity, turbidity, chlorophyll, dissolved oxygen,  $\delta^{18}\text{O}$ , d-excess and bathymetry in front of Bowdoin and Sun Glaciers, tidewater glaciers in northwestern Greenland, in 2014 and 2015 summer. We also performed high spatial (2560x1920 pixel) and temporal (10 s) time-lapse photography in front of Bowdoin Glacier in 2015 July. Below the depths of 280 m (2014) and 250 m (2015) in the Bowdoin Fjord, we observed Atlantic Water (AW), which is believed as the heat source of submarine melting. The mean temperature and salinity within the layer of AW was 1.2°C and 34.4 g kg<sup>-1</sup> in 2014, and 1.1°C and 34.5 g kg<sup>-1</sup> in 2015. The results suggested that warm water flows into the Bowdoin Fjord from the open ocean. Contrasting to these observations at Bowdoin, AW was missing in front of Sun Glacier and relatively fresh and cold water mass was found. It was suggested that relatively shallow bathymetry (~100-m deep) and existence of sill (~10-m deep) inhibited entering warm water from the open ocean. Near the surface in the vicinity of plume of Sun Glacier (~200 m away from the front), water mass properties were completely different from those in the open ocean. Water was highly turbid, fresh and cold, suggesting subglacial discharge of meltwater as the origin of the water. The time-lapse photographs revealed fjord circulation near the ice-ocean interface. It was clear that the circulation was driven by buoyant plume, which was generated subglacial discharge and/or submarine melt. In early July, waters emerged by buoyant plume was always visible along the surface approximately 5 km from the front, but it was only visible near the front in late July. The change from early July to late July may be explained by amount of subglacial discharge and the stability of stratification near the ocean surface as reported by recent modelling studies. Our observations water mass structures and circulation in the fjord in front of the tidewater glaciers in northwestern Greenland, which are important properties to calculate submarine melting rate in two different types of tidewater glaciers.

Keywords: Tidewater glacier, Greenland, Fjord, Submarine melting

## Mass balance fluctuation of Qaanaaq Ice Caps and its vicinity in northeastern Greenland

\*Keiko Konya<sup>1</sup>, Jun Saito<sup>2</sup>, Mihiro Maruyama<sup>2</sup>, Shin Sugiyama<sup>2</sup>

1.DEGCR, Japan Agency for Marine-Earth Science and Technology, 2.Institute of Low Temperature Science, Hokkaido University

Many of glaciers and ice caps (GICs) exist at the margin of the Greenland. The contribution of GICs mass loss to sea level rise under recent warming is large. The northeastern Greenland is one of the areas, which experienced large mass change and has little in-situ mass balance observation. Bolch et al. (2013) estimated that the surface elevation change of ice caps in northeastern Greenland from 2003 to 2008 was  $-0.6 \text{ m a}^{-1}$ . Saito et al. (2015) revealed that the mean surface level change of six ice caps in northeastern Greenland from 2006 to 2010 was  $-1.1 \text{ m a}^{-1}$ .

We have estimated surface mass balance of five ice caps in northern Greenland by a mass balance model constructed by Hock (1999). The model takes temperature index method for calculating ablation. The model computes spatial variation of surface mass balance for the ice caps. The 100m - gridded DEM and surface condition of the ice cap, derived from modified ALOS (Advanced Land Observing Satellite) data, were used for the model calculation. The climate data as input of the model was air temperature and precipitation at Thule climate station (TCS, 77.2N, 68.4W), which is one of the long-term running climate stations in Greenland situated about 100 km south to Qaanaaq. The air temperature at TCS has been increasing after 1990.

The calculation has been done for Qaanaaq Ice Cap (QIC) since 1980s. The mass balance of QIC has been negative since 1980s. The mass balance calculation by the model has been done for other four ice caps for 2006 -2009 and compared with the surface elevation change reported by Saito et al. (2015). Ice caps situated in coastal area show less negative mass balance than those situated inland. The ice caps situated in coastal area show higher albedo (Saito, et al., 2015), which is possibly because higher fraction of precipitation falls as snow.

Keywords: Greenland, mass balance, climate change, ice cap

## New developments of Arctic Data archive System(ADS)

\*Hironori Yabuki<sup>1,2</sup>, Takeshi Sugimura<sup>2</sup>, Takeshi Terui<sup>2</sup>

1.Japan Agency for Marine-Earth Science and Technology, 2.National Institute of Polar Research

Arctic is the region where the global warming is mostly amplified, and the atmosphere/ ocean/ cryosphere/ land system is changing. Active promotion of Arctic environmental research, it is large and responsible for observational data. Promotion of Arctic research in Japan, has not been subjected to independent in their respective fields.

In the National Institute of Polar Research, perform the integration and sharing of data across a multi-disciplinary such as atmosphere, ocean, snow and ice, land, ecosystem, model, for the purpose of cooperation and integration across disciplines, we build a Arctic Data archive System (ADS).

Arctic Data archive System (ADS), to promote the mutual use of the data across a multi-disciplinary to collect and share data sets, such as observational data, satellite data, numerical experiment data. Through these data sets, clarify of actual conditions and processes of climate change on the Arctic region, and further contribute to assessment of the impact of global warming in the Arctic environmental change, to improve the future prediction accuracy.

A new project of the Arctic research (ArCS :Arctic Challenge for Sustainability) has been started in 2015. ArCS is a national flagship project funded by the Ministry of Education, Culture, Sports, Science and Technology. The National Institute of Polar Research (NIPR), Japan Agency for Marine-Earth Science and Technology (JAMSTEC) and Hokkaido University are playing the key roles in this project, and will continue to carry it out for approximately four-and-a-half years from September 2015 to March 2020. ADS is responsible for the data management of this project.

Keywords: Arctic, Environment, Global Warming, ArCS, Data



## Access analysis of data visualization web service for polar region

\*Takeshi Terui<sup>1</sup>, Takeshi Sugimura<sup>1</sup>, Hironori Yabuki<sup>1</sup>

### 1. National Institute of Polar Research

Arctic Data archive System (ADS) has been constructed for archiving or distributing various datasets, which are observed in the Arctic region by Japanese researchers. ADS was developed the following web services during GRENE Arctic Climate Change Research Project.

1. Metadata search service KIWA.
2. Online visualization application VISION.
3. Quasi-real-time polar environment observation monitor VISHOP.

These Web services are accessed from all over the world. In this study, we performed the web access analysis to the period of GRENE project. We will present the analysis results, such as the time-series of access number and viewers' behavior on ADS web site.

Keywords: Web Service, Access Analysis, Visualization, Data