

黒潮大蛇行を引き起こす傾圧不安定の発達に膠州海山が果たす役割

Effects of Koshu Seamount on the Development of Baroclinic Instability Leading to the Kuroshio Large Meander

*田中 祐希¹、日比谷 紀之¹*Yuki Tanaka¹, Toshiyuki Hibiya¹

1. 東京大学大学院理学系研究科

1. Graduate School of Science, The University of Tokyo

It is well known that the Kuroshio south of Japan shows bimodal path fluctuations between the large meander (LM) path and the nonlarge meander (NLM) path. The transition from the NLM path to the LM path is triggered by a small meander which is generated off the southeastern coast of Kyushu and then propagates eastward to Cape Shiono-misaki while being amplified slowly through baroclinic interaction with a lower layer cyclone-anticyclone pair. This small meander thereafter amplifies rapidly over Koshu Seamount located about 200 km to the south of Cape Shiono-misaki, leading to the formation of the LM path. Although it is shown that the existence of Koshu Seamount is essential for the rapid amplification of the small meander, the underlying physical mechanism has not been fully understood yet.

In this study, the effects of Koshu Seamount on the development of baroclinic instability leading to the formation of the LM path of the Kuroshio are investigated using a two-layer quasi-geostrophic model taking into account the effect of bottom topography. Numerical experiments show that the transition processes from the NLM path to the LM path can be successfully reproduced only when the bottom topography mimicking Koshu Seamount is taken into account. In this case, the upper layer meander trough is amplified rapidly together with the lower layer cyclone-anticyclone pair during their passage over the seamount. This suggests that the transition from the NLM path to the LM path is caused by baroclinic instability enhanced over the seamount. A linear stability analysis with the bottom topography mimicking Koshu Seamount shows that baroclinic instability over the seamount is caused by a coupling between the upper layer Rossby wave propagating eastward in the background geostrophic flow and the lower layer topographically trapped wave propagating clockwise around the seamount. These two waves in the upper and lower layers propagate in the same direction with nearly the same speed so that they can resonantly interact with each other over the northern slope of the seamount. The spatial structure of the most unstable mode is shown to be close to that of the rapidly amplifying meander trough over the seamount reproduced in the numerical experiment.

キーワード：黒潮大蛇行、膠州海山、傾圧不安定、地形性捕捉波、二層準地衡流モデル、線形安定性解析

Keywords: Large Meander of Kuroshio, Koshu Seamount, Baroclinic Instability, Topographically Trapped Wave, Two-Layer Quasi-Geostrophic Model, Linear Stability Analysis

Dynamics of the Atlantic meridional overturning circulation and Southern Ocean in an ocean model of intermediate complexity

Dynamics of the Atlantic meridional overturning circulation and Southern Ocean in an ocean model of intermediate complexity

McCreary Julian P.²、*Furue Ryo¹、Schloesser Fabian²、Burkhardt Theodore W.³、野中 正見¹
Julian P McCreary²、*Ryo Furue¹、Fabian Schloesser²、Theodore W Burkhardt³、Masami Nonaka¹

1.APL, JAMSTEC、2.ハワイ大学、3.テンブル大学

1.APL, JAMSTEC, 2.Univ. of Hawaii, 3.Temple Univ.

A steady-state, variable-density, 2-layer, ocean model (VLOM) is used to investigate basic dynamics of the Atlantic meridional overturning circulation and Southern Ocean. The domain consists of idealized (rectangular) representations of the Atlantic, Southern, and Pacific Oceans. The model equations represent the depth-averaged, layer-1 response.

A hierarchy of solutions is obtained in which forcings and processes are individually introduced. The complete solution set includes a wide variety of solution types: with sinking in the northern North Atlantic and with sinking near Antarctica; with and without wind forcing; with, without, and for two parameterizations of northern-boundary sinking that represent cooling external to and within the North Atlantic; for a wide range of mesoscale-eddy mixing strength and wind stress in the Antarctic Circumpolar region; and for different closures for mesoscale-eddy mixing. Novel aspects of the model and solutions include the following: use of VLOM, which allows buoyancy forcing to be introduced realistically; the aforementioned closure, which allows eddy-induced transport to be determined when layer 1 represents *_both_* the surface mixed layer ($h=h_m$) and the depth of subsurface isopycnals ($h>h_m$); latitude where layer 1 outcrops in the Southern Ocean being *_internally_* determined rather than externally specified; and a boundary layer, based on Gill's (1968) solution, that smoothly connects the Southern- and Atlantic-Ocean responses across the latitude of the southern tip of South America. Finally, some solutions in the set are comparable to solutions to idealized, ocean general circulation models (OGCMs); in these cases, our solutions provide insight into the underlying dynamics of the OGCM solutions, for example, pointing toward processes that may be involved in eddy saturation and compensation.

キーワード：海洋深層循環、層モデル、南大洋

Keywords: Oceanic deep circulation, Layer model, Southern Ocean

回転系における深水波の直接数値計算

Direct numerical simulation of deep-water waves in rotating frame

*藤原 泰¹、吉川 裕¹、松村 義正²*Yasushi Fujiwara¹, Yutaka Yoshikawa¹, Yoshimasa Matsumura²

1.京都大学大学院理学研究科、2.北海道大学低温科学研究所

1.Graduate School of Science, Kyoto University, 2.Institute of Low Temperature Science, Hokkaido University

The ratio of the periods of oceanic surface waves (wind waves and swells) and the inertial period is about 10^{-4} , and the earth's rotation does not greatly affect the orbital motion of fluid particles. Waves, however, do modify mean flow under the influence of Coriolis force. This is because the slight tilt of the orbital plane of fluid particle generates Reynolds stress. Hasselmann (1970) and Huang (1979) demonstrated that this Reynolds stress induced by waves can be expressed as the Coriolis force acting on the Stokes drift. The latter expression is called Coriolis-Stokes forcing. In ocean surface layer studies, Coriolis-Stokes forcing has been used as a standard formulation to incorporate the wave-stress effect. However, Coriolis-Stokes forcing is derived under several assumptions, and there has been no research that directly examined the appropriateness of the forcing.

Here we investigated the Coriolis-Stokes forcing, by performing direct numerical simulations of deep-water waves using a recently developed free-surface nonhydrostatic numerical model. The new scheme that this model adopts allows for the accurate simulation of the orbital motion and the dispersion relation of deep water waves, which could not be achieved by the conventional mode-splitting scheme.

Simulations were carried out under idealized conditions of x-z two dimensional domain with periodic horizontal boundaries. Waves were maintained by surface pressure perturbation. Reynolds stress was obtained from the velocity field, and the Coriolis-Stokes forcing was calculated using the Stokes drift, which we obtained from the on-line particle tracking.

Comparison of the two forces tells us that the Coriolis-Stokes forcing is nearly identical to the wave stress under the idealized condition. These forces induce a Eulerian response to the Stokes drift. In the existence of viscosity, this Eulerian flow generates the spiral current throughout the Ekman depth, even there is no net momentum input from the surface. By imposing a uniform stress on the surface, we also find that the mean current profile is described by Ekman-Stokes solution (Polton et al., 2005), which is obtained by considering Coriolis-Stokes forcing, rather than the classical Ekman solution. We will be presenting the results of further simulations under various conditions.

キーワード：波・平均流相互作用、コリオリ・ストークス力、自由表面非静力学モデル

Keywords: wave-mean flow interaction, Coriolis-Stokes forcing, free-surface nonhydrostatic numerical model

流動化現象が定常状態に達するまでの過程

A route to steady state of liquid fluidization

*金 芳隆¹、栗田 敬¹*Yoshitaka Kon¹, Kei Kurita¹

1. 東京大学地震研究所

1. Earthquake Research Institute, The University of Tokyo

When fluid is injected with low flux into the saturated packed bed of particles, fluid flows through interstitial spaces. This is known as a permeable flow. In this situation an empirical relationship, the Darcy law is known to exist in laminar flow, where the fluid flux is proportional to the applied pressure difference. When the applied pressure difference is increased and hence, the fluid flux is increased and both fluid and particles begin to flow together as a suspension. This gross flow called fluidization is important because it can drastically change efficiency of material transport. In the field of hydrology, for an example, this is related to an initiation of debris flow. In the field of volcanology, this process is related to the rejuvenation of a dormant magma chamber where the injection of new magma into the crystal mush causes the replenishment. In the field of chemical engineering, fluidization is well investigated for the engineering applications. The transition between a permeable flow and a gross flow is a kind of phase change; This critical superficial velocity is known as a minimum fluidization velocity (v_{mf}). The dynamics of the change from a permeable flow state to a gross flow state is more important in the initiation of such as debris flow and rejuvenation of magma chamber. The process of the initiation of fluidization is as follows; Water start to flow with high velocity enough to fluidize, both upper and fluidized front are lifted from top and bottom of particles bed, respectively (Slis et al. (1959)). Gibilaro et al. (1984) derived that upper front velocity has constant value until it reaches steady state and depends on water flux, and Thelen and Ramirez (1997) confirmed it experimentally. Upper front velocity is discussed like above, however, fluidized front velocity is not observed experimentally. In this presentation, we focus on the dynamics of the initiation process of the fluidization and present experimental approaches.

To observe the initiation process of the fluidization, we employed a vertical fluidizing bed. Transparent acrylic pipe (inside diameter: 30 mm, length: 40~120 cm) is used, where particles are packed at the bottom of 12.5 cm. From water-saturated state, water is injected from the bottom at constant flux (the superficial velocity of 0.5~4.7 cm/s). The hydrostatic pressures are monitored at 3 different positions at 0, 5, 10 cm high from bottom. Glass beads (diameter: 0.8 mm, density: 2.5 g/cm³) and polystyrene beads (diameter: 0.8 mm, density: 1.03 g/cm³) are used as particles. The all move of beads are filmed, and inside area of this pipe is divided into fixed, fluidized and no particle area by differential of particles density.

We found that top of the particle bed and the fluidization area propagates from bottom to upward when the injected velocity is above the critical value v_{mf} . Because fluidization front velocity is larger than upper front velocity, the thickness of fixed bed becomes gradually small, and after that all beads become fluidized. On checking pressure and movie of the same time, it is revealed that hydrostatic pressure gradient of fixed bed are larger than fluidized bed during rise. Propagation of fluidized bed is divided into two types by flux; In case injected velocity is slightly larger than v_{mf} , the propagation ends when both upper and fluidization front reach same height. In case injected velocity is enough high compared with v_{mf} , rise lasts after a while that. Fluidized front velocity depends on injected flux when it is low, however, fluidized front velocity saturates when it is high. Porosity of propagating fluidized bed is kept nearly equal value to one

of terminal fluidized bed.

We also compare the cases for soft gels are used as elastic particles and discuss the effect of modification of packing and particle shape during flow.

キーワード：流動化、粒子層、物質輸送、サスペンション、室内実験

Keywords: Fluidization, Bed of particles, Substance transportation, Suspension, Laboratory experiment

吸収線輪郭から探る太陽表面对流速度の高さ構造

Height structure of solar surface convective velocity from absorption line profile

*大場 崇義¹、飯田 佑輔²、清水 敏文²

*Takayoshi Oba¹, Yusuke Iida², Toshifumi Shimizu²

1.総合研究大学院大学物理科学研究科、2.宇宙科学研究所/宇宙航空研究開発機構

1.The Graduate University for Advanced Studies, 2.Institute of space and astronomical science / Japan aerospace exploration agency

恒星において流体の対流現象は、内部のエネルギー輸送やひいては恒星大気活動などその全域で重要な役割を果たす。恒星の内部構造は、コアにおける核融合反応によって発生したエネルギーを如何に宇宙空間に運びだすかで決定される。この恒星における対流の働きを理解する上で、我々に最も近くに位置する太陽は最もよい例となる。太陽では、コアで発生したエネルギーは光の放射で外層に運ばれる。ある程度外層(~0.7 太陽半径)まで達すると、流体による対流運動により運搬される。その対流運動は、私たちが可視光で観測できる太陽表面に達し、そのガス運動はさらに上空の大気(コロナ)の加熱や様々な大気ダイナミクスのエネルギー源となる。このように太陽表面对流は内部から運搬されたエネルギーを大気に伝える役割を果たすが、その運動は完全に理解されているわけではない。私達が見ることができる太陽表面は明るい「粒状斑」と暗い「間隙」で覆われている。粒状斑は内部からの対流が吹き出した上昇流領域であり、間隙はそれが下降流に転じた領域である。この上昇流が下降流に転じる過程は、主に数値シミュレーションによって研究されている。しかしながら、実際の太陽においてこの速度を失うプロセスが何の物理過程によって起こっているかは定かではない。このプロセスは高さ方向への運動中に起こるが、観測データから十分な時間・空間分解能を持って対流速度の高さ構造を取得できないという問題があった。本研究では、太陽観測衛星「ひので」の可視光望遠鏡/スペクトロポラリメータ(SOT/SP)の分光データに吸収線輪郭の解析を行うことで、対流速度の高さ変化を調べた。ある原子や分子における吸収線輪郭において異なった輝度はそれぞれ異なった高さを反映する。そのため、輝度毎のドップラー速度は高さ毎の速度に対応する。宇宙空間から安定した観測を実施できる「ひので」は、粒状斑と間隙を空間分解した上で高精度の吸収線を観測できる点から、本解析に最適である。得られた対流速度を粒状斑・間隙に分類して解析を行った。粒状斑においては上昇流が加速したのちに減速に転ずる傾向が見られた。一方で間隙においては沈みこむにつれて下降速度が大きくなる傾向を捉えることができた。間隙において下降速度場が大きくなることは、従来の1次元定常大気の描像では説明できない。講演では加速を引き起こすメカニズムの候補について議論する。

キーワード：対流、太陽

Keywords: convection, the sun